

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 August 2002 (01.08.2002)

PCT

(10) International Publication Number  
**WO 02/059106 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 401/04**,  
401/14, 407/14, 409/14, A61K 31/445, 31/44, 31/495,  
A61P 29/00, 35/00, 37/00

(74) Agents: **INSOGNA, Anthony, M. et al.**; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).

(21) International Application Number: **PCT/US01/50401**

(22) International Filing Date:  
21 December 2001 (21.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/258,372 27 December 2000 (27.12.2000) US  
09/972,487 5 October 2001 (05.10.2001) US

(71) Applicant (for all designated States except US): **CEL-GENE CORPORATION** [US/US]; 7 Powder Horn Drive, Warren, NJ 07059 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ROBARGE, Michael, J.** [US/US]; 190 Sandford Avenue, North Plainfield, NJ 07060 (US). **CHEN, Roger, Shen-Chu** [US/US]; 110 Christie Street, Edison, NJ 08820 (US). **MULLER, George, W.** [US/US]; 250 Windmill Court, Bridgewater, NJ 08807 (US). **MAN, Hon-Wah** [US/US]; 27 Grant Way, Princeton, NJ 08540 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **ISOINDOLE-IMIDE COMPOUNDS, COMPOSITIONS, AND USES THEREOF**



(57) **Abstract:** The invention relates to isoindole-imide compounds and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, or mixtures of stereoisomers thereof, pharmaceutical compositions comprising these isoindole-imide compounds, and methods for reducing the level of cytokines and their precursors in mammals. In particular, the invention pertains to isoindole-imide compounds that are potent inhibitors of the production of the TNF- $\alpha$  in mammals. The isoindole-imides described herein are useful for treating or preventing diseases or disorders in mammals, for example, cancers, such as solid tumors and blood-borne tumors; heart disease, such as congestive heart failure; osteoporosis; and genetic, inflammatory, allergic; and autoimmune diseases.

**ISOINDOLE-IMIDE COMPOUNDS, COMPOSITIONS, AND USES THEREOF**

This application is a continuation-in-part of United States Application No. 09/972,487 filed October 5, 2001, which claims the benefit of U.S. Provisional Application  
5 No. 60/258,372, filed December 27, 2000, and which applications are hereby expressly incorporated by reference herein.

**1. Field of The Invention**

The invention encompasses novel compounds including compounds having an  
10 isoindole-imide moiety, pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, or mixtures of stereoisomers thereof, pharmaceutical compositions of these compounds, and methods of using these compounds and compositions in mammals for treatment or prevention of diseases.

**2. Introduction**

The present invention relates to isoindole-imide compounds and pharmaceutically acceptable salts, hydrates, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof; pharmaceutical compositions comprising these isoindole-imide compounds; and methods for reducing the level of cytokines and their precursors in  
20 mammals. In particular, the invention includes isoindole-imide compounds that have one or more of the following activities: modulation of the production of TNF- $\alpha$ ; modulation of the production of IL-1 $\beta$ ; stimulation of the production of IL-10; or stimulation of the production T-cells.

The isoindole-imides described herein are useful for treating or preventing diseases  
25 or disorders in mammals, for example, cancers, such as solid tumors and blood-born tumors. Specific examples of cancers treatable or preventable by compounds of the invention include, but are not limited to, cancers of the skin, such as melanoma; lymph node; breast; cervix; uterus; gastrointestinal tract; lung; ovary; prostate; colon; rectum; mouth; brain; head and neck; throat; testes; kidney; pancreas; bone; spleen; liver; bladder;  
30 larynx; nasal passages; and AIDS-related cancers. The compounds are particularly useful for treating cancers of the blood and bone marrow, such as multiple myeloma and acute and chronic leukemias, for example, lymphoblastic, myelogenous, lymphocytic, and myelocytic leukemias.

35

The compounds of the invention are also useful to treat or prevent heart disease, such as congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, and myocardial infarction.

The compounds of the invention can also be used to treat or prevent viral, genetic, inflammatory, allergic, and autoimmune diseases. For example, the compounds are useful to treat or prevent diseases including, but not limited to, HIV; hepatitis; adult respiratory distress syndrome; bone-resorption diseases; chronic pulmonary inflammatory diseases; dermatitis; cystic fibrosis; septic shock; sepsis; endotoxic shock; hemodynamic shock; sepsis syndrome; post ischemic reperfusion injury; meningitis; psoriasis; fibrotic disease; cachexia; graft rejection including graft versus host disease; auto-immune disease; rheumatoid spondylitis; arthritic conditions, such as rheumatoid arthritis and osteoarthritis; osteoporosis; Crohn's disease; ulcerative colitis; inflammatory-bowel disease; multiple sclerosis; systemic lupus erythematosus; ENL in leprosy; radiation damage; asthma; and hyperoxic alveolar injury.

The compounds of the invention are also useful for treating or preventing bacterial infections or the symptoms of bacterial infections including, but not limited to, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

### **3. Background of The Invention**

Tumor necrosis factor alpha, (TNF- $\alpha$ ) is a cytokine that is released primarily by mono-nuclear phagocytes in response to immunostimulators. TNF- $\alpha$  is capable of enhancing most cellular processes, such as differentiation, recruitment, proliferation, and proteolytic degradation. At low levels, TNF- $\alpha$  confers protection against infective agents, tumors, and tissue damage. But TNF- $\alpha$  also has role in many disease processes. When administered to mammals or humans, TNF- $\alpha$  causes or aggravates inflammation, fever, cardiovascular effects, hemorrhage, coagulation, and acute phase responses similar to those seen during acute infections and shock states. Enhanced or unregulated TNF- $\alpha$  production has been implicated in a number of diseases and medical conditions, for example, cancers, such as solid tumors and blood-born tumors; heart disease, such as congestive heart failure; and viral, genetic, inflammatory, allergic, and autoimmune diseases.

The interleukins are a subclass of the cytokine family and possess a wide spectrum of biological activities including involvement in cell activation, cell differentiation, cell proliferation, and cell-to-cell interactions. Interleukin 1 beta (IL-1 $\beta$ ) and interleukin 10 (IL-10), in combination with other cytokines, play a central role in mediating inflammatory

processes and IL-1 $\beta$  has been implicated as both a growth factor and growth suppressor in certain tumor cells.

T-cells are a class of white blood cells that play an important role in the immune response, and help protect the body from viral and bacterial infections. Diminished T-cell  
5 levels strongly contribute to the inability of HIV patients to combat infections, and abnormally low T-cell levels are prominent in a number of other immune deficiency syndromes, including DiGeorge Syndrome, and in certain forms of cancer, such as T-cell lymphoma.

Cancer is a particularly devastating disease, and increase in blood TNF- $\alpha$  levels are  
10 implicated in the risk of and the spreading of cancer. Normally, in healthy subjects, cancer cells fail to survive in the circulatory system, one of the reasons being that the lining of blood vessels acts as a barrier to tumor-cell extravasation. But increased levels of cytokines, have been shown to substantially increase the adhesion of cancer cells to endothelium *in vitro*. One explanation is that cytokines, such as TNF- $\alpha$  stimulate the  
15 biosynthesis and expression of a cell surface receptors called ELAM-1 (endothelial leukocyte adhesion molecule). ELAM-1 is a member of a family of calcium-dependent cell adhesion receptors, known as LEC-CAMs, which includes LECAM-1 and GMP-140. During an inflammatory response, ELAM-1 on endothelial cells functions as a "homing receptor" for leukocytes. Recently, ELAM-1 on endothelial cells was shown to mediate the  
20 increased adhesion of colon cancer cells to endothelium treated with cytokines (Rice *et al.*, 1989, *Science* 246:1303-1306). It has been suggested that an uncontrolled synthesis of IL-1 $\beta$  in leukemia blast cells is thought to result in the production of factors which promote proliferation of these malignant cells (Hestdal *et al.*, 1992, *Blood* 80: 2486-94). In addition to this, IL-1 $\beta$ , in combination with other cytokines, appears to stimulate the growth of  
25 human gastric and thyroid carcinoma cells (Ito *et. al.*, 1993, *Cancer Research* 53: 4102-6).

Inflammatory diseases such as arthritis, related arthritic conditions (*e.g.* osteoarthritis and rheumatoid arthritis), inflammatory bowel disease, sepsis, psoriasis, and chronic inflammatory pulmonary diseases are also prevalent and problematic ailments. Both TNF- $\alpha$  and IL-1 $\beta$  play central roles in the inflammatory response and the administration of  
30 their antagonists block chronic and acute responses in animal models of inflammatory disease. Conversely, IL-10 is an anti-inflammatory cytokine and is responsible for down-regulating inflammatory responses and as such possesses anti-inflammatory ability, including the suppression of production of proinflammatory cytokines such as TNF- $\alpha$  and IL-1 $\beta$

35



Heart disease has caused wide-spread death and debilitation. TNF- $\alpha$  has been implicated in a broad variety of cardiac pathophysiological conditions, such as septic shock, acute viral myocarditis, cardiac allograft rejection, myocardial infarction, and congestive heart failure (*see e.g.*, Steadman *et al.*, 1988, *IEEE Trans. Biomed. Eng.* 35:264-272; Tracey *et al.*, 1986, *Science Wash. DC* 234:470-474; for a review see Ferrari, 1998, *Cardiovascular Research* 37:554-559). In one study, it was found that protective TNF- $\alpha$  binding proteins are downregulated in the hearts of patients with advanced congestive heart failure. During the study it was found that a large percentage of the diseased hearts analyzed had elevated TNF- $\alpha$  levels. The authors noted that the results support the proposition that the heart itself is a target of TNF- $\alpha$  and that myocardial TNF- $\alpha$  production may be a maladaptive mechanism that contributes to progressive heart failure (Torre-Amione *et al.*, 1996, *Circulation* 93:704-711). In other studies, it has been demonstrated *in-vitro* and *in-vivo* (feline) that TNF- $\alpha$  is produced in the myocardium portion of the heart upon endotoxin stimulation. These studies provide compelling evidence indicating that a pathogenic level of biologically active TNF- $\alpha$  may be produced in the heart during endotoxin-mediated septic shock. And that such local concentrations of TNF- $\alpha$  may be the primary instigator of myocardial-function depression during systemic sepsis (Kapadia *et al.*, 1995, *J. Clin. Invest.* 96:1042-1052). Thus, inhibitors of TNF- $\alpha$  activity may prevent its deleterious effects on the heart. For example, it has been demonstrated that soluble TNF-binding proteins modulate the negative inotropic effects of TNF- $\alpha$  *in vitro* in isolated contracting cardiac myocytes (Kapadia *et al.*, 1995, *Am. J. Physiol.* 268:H517-H525).

Enhanced or unregulated TNF- $\alpha$  production has been implicated in viral, genetic, inflammatory, allergic, and autoimmune diseases, for example, HIV; hepatitis; adult respiratory distress syndrome; bone-resorption diseases; chronic pulmonary inflammatory diseases; dermatitis; cystic fibrosis; septic shock; sepsis; endotoxic shock; hemodynamic shock; sepsis syndrome; post ischemic reperfusion injury; meningitis; psoriasis; fibrotic disease; cachexia; graft rejection; auto-immune disease; rheumatoid spondylitis; arthritic conditions, such as rheumatoid arthritis and osteoarthritis; osteoporosis, Crohn's disease; ulcerative colitis; inflammatory-bowel disease; multiple sclerosis; systemic lupus erythematosus; ENL in leprosy; radiation damage; asthma; and hyperoxic alveolar injury. For discussions see Tracey *et al.*, 1987, *Nature* 330:662-664 and Hinshaw *et al.*, 1990, *Circ. Shock* 30:279-292 (endotoxic shock); Dezube *et al.*, 1990, *Lancet*, 335:662 (cachexia); Millar *et al.*, 1989, *Lancet* 2:712-714 and Ferrai-Baliviera *et al.*, 1989, *Arch. Surg.* 124:1400-1405 (adult respiratory distress syndrome); Bertolini *et al.*, 1986, *Nature* 319:516-518, Johnson *et al.*, 1989, *Endocrinology* 124:1424-1427, Holler *et al.*, 1990, *Blood*

- 75:1011-1016, and Grau *et al.*, 1989, *N. Engl. J. Med.* 320:1586-1591 (bone resorption diseases); Pignet *et al.*, 1990, *Nature*, 344:245-247, Bissonnette *et al.*, 1989, *Inflammation* 13:329-339 and Baughman *et al.*, 1990, *J. Lab. Clin. Med.* 115:36-42 (chronic pulmonary inflammatory diseases); Elliot *et al.*, 1995, *Int. J. Pharmac.* 17:141-145 (rheumatoid arthritis); von Dullemen *et al.*, 1995, *Gastroenterology*, 109:129-135 (Crohn's disease); Duh *et al.*, 1989, *Proc. Nat. Acad. Sci.* 86:5974-5978, Poll *et al.*, 1990, *Proc. Nat. Acad. Sci.* 87:782-785, Monto *et al.*, 1990, *Blood* 79:2670, Clouse *et al.*, 1989, *J. Immunol.* 142, 431-438, Poll *et al.*, 1992, *AIDS Res. Hum. Retrovirus*, 191-197, Poli *et al.* 1990, *Proc. Natl. Acad. Sci.* 87:782-784, Folks *et al.*, 1989, *PNAS* 86:2365-2368 (HIV and opportunistic infections resulting from HIV).

- Pharmaceutical compounds that can block the activity or inhibit the production of certain cytokines, including TNF- $\alpha$  and IL-1 $\beta$ , may be beneficial therapeutics. Many small-molecule inhibitors have demonstrated an ability to treat or prevent inflammatory diseases implicated by TNF- $\alpha$  (for a review see Lowe, 1998 *Exp. Opin. Ther. Patents* 8:1309-1332).
- 15 In addition, pharmaceutical compounds that can stimulate the activity or increase the production of certain cytokines, including IL-10, and immune response factors such as T-cells, may be beneficial therapeutics.


- Thalidomide is an emerging immunotherapeutic agent and, in addition to utility in treating a variety of inflammatory disorders, it is projected to be useful in treating cancers (see e.g., Marriott *et al.*, 1999, *Immunology Today* 20:537-540). Thalidomide has been shown to inhibit production of both TNF- $\alpha$  and IL-1 $\beta$  while simultaneously increasing the production of IL-10 and T-cells, and has been tested against a variety of autoimmune and inflammatory diseases, see e.g., Gutierrez-Rodriguez, 1984, *Arth. and Rheum* 27:1118; *The Physician's Desk Reference*, 54th edition, 911-916, Medical Economics Company (2000).
- 25 Thalidomide's teratogenic properties, however, have limited its use and driven efforts to discover analogs or derivatives with reduced toxicity and improved therapeutic activity. The design of thalidomide analogs and derivatives attempts to maintain/enhance activity while subverting toxicity (for a discussion of some recent advances in TNF- $\alpha$  inhibitors structurally related to thalidomide see Marriott, 1997, *Exp. Opin. Invest. Drugs* 6:1105-1108). For example, the following references have disclosed alternatives to thalidomide as inhibitors of TNF- $\alpha$  production: U.S. Patent Nos. 5,385,901; 5,635,517; and 5,798,368 and PCT International Application WO 98/54170. Despite these disclosures, there remains a need for non-toxic and high-potency compounds that treat or prevent cancer, inflammatory disorders, and autoimmune diseases.

Citation or identification of any reference in Section 3 of this application is not an admission that such reference is available as prior art to the present invention.

#### 4. Summary of the Invention

The invention encompasses novel isoindole-imide compounds and compositions thereof that are useful to treat or prevent diseases in mammals, including humans. The invention further encompasses the use of these compounds for treating or preventing diseases or disorders including, but not limited to, cancer; viral, genetic, inflammatory, allergic, and autoimmune diseases; and bacterial infections. The compounds of the invention are particularly useful to treat or prevent diseases caused or aggravated by excessive or unregulated levels of TNF- $\alpha$ , or IL-1 $\beta$ ; or diminished or unregulated levels of IL-10 or T-cells.

In one embodiment, the invention relates to compounds encompassed by Formula I :



I

wherein:

one of X and Y is C=O and the other is CH<sub>2</sub> or C=O;

R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(S)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(O)NHR<sup>3</sup>, C(S)NHR<sup>3</sup>, C(O)NR<sup>3</sup>R<sup>3'</sup>, C(S)NR<sup>3</sup>R<sup>3'</sup> or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

R<sup>2</sup> is H, F, benzyl, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, or (C<sub>2</sub>-C<sub>8</sub>)alkynyl;

R<sup>3</sup> and R<sup>3'</sup> are independently (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl,

(C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>,  
 (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;

R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl,  
 5 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

each occurrence of R<sup>6</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl,  
 10 benzyl, aryl, (C<sub>2</sub>-C<sub>5</sub>)heteroaryl, or (C<sub>0</sub>-C<sub>8</sub>)alkyl-C(O)O-R<sup>5</sup> or the R<sup>6</sup> groups can join to  
 form a heterocycloalkyl group;

n is 0 or 1; and

15 \* represents, in formula I and in the following formulas of the invention, a chiral-carbon  
 center; with the proviso that when n is 0 then R<sup>1</sup> is not H.

In a separate embodiment, the compounds of the invention include compounds of  
 formula I, wherein when n is 0 then R<sup>1</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl,  
 (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl,  
 20 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>,  
 (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(S)NHR<sup>3</sup>, or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

R<sup>2</sup> is H or (C<sub>1</sub>-C<sub>8</sub>)alkyl; and


25 R<sup>3</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl,  
 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl,  
 (C<sub>5</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>; (C<sub>0</sub>-C<sub>8</sub>)alkyl-NH-C(O)O-R<sup>5</sup>; (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>,  
 (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>; and the other variables are as  
 described above.

30 In another embodiment of the compounds of formula I, R<sup>2</sup> is H or (C<sub>1</sub>-C<sub>4</sub>)alkyl.

In still another embodiment of the compounds of formula I, R<sup>1</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl or  
 benzyl.

In yet another embodiment of the compounds of formula I, R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl,  
 benzyl, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or


35



5

In another embodiment of the compounds of formula I, R<sup>1</sup> is

10



wherein Q is O or S, and each occurrence of R<sup>7</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, halogen, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>, or adjacent occurrences of R<sup>7</sup> can be taken together to form a bicyclic alkyl or aryl ring.


In still another embodiment of the compounds of formula I, R<sup>1</sup> is C(O)R<sup>3</sup> or C(O)OR<sup>4</sup>.

In another embodiment of the compounds of formula I, R<sup>3</sup> is (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>1</sub>-C<sub>8</sub>)alkyl, aryl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>.

In yet another embodiment of the compounds of formula I, heteroaryl is pyridyl, furyl, or thienyl.

In another embodiment of the compounds of formula I, the H of C(O)NHC(O) can be replaced with (C<sub>1</sub>-C<sub>4</sub>)alkyl, aryl, or benzyl.

In another embodiment, the invention encompasses compounds of Formula II:



35

## II

wherein:

R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl,  
 5 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(S)R<sup>3</sup>,  
 C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(O)NHR<sup>3</sup>,  
 C(S)NHR<sup>3</sup>, C(O)NR<sup>3</sup>R<sup>3'</sup>, C(S)NR<sup>3</sup>R<sup>3'</sup> or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

R<sup>2</sup> is H or (C<sub>1</sub>-C<sub>8</sub>)alkyl;

10

R<sup>3</sup> and R<sup>3'</sup> are independently (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl,  
 (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl,  
 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>,  
 (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;

15

R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl,  
 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;


20

each occurrence of R<sup>6</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl,  
 benzyl, aryl, (C<sub>2</sub>-C<sub>5</sub>)heteroaryl, or (C<sub>0</sub>-C<sub>8</sub>)alkyl-C(O)O-R<sup>5</sup> or the R<sup>6</sup> groups can join to  
 form a heterocycloalkyl group; and

25 the \* represents a chiral-carbon center.


In another embodiment of the compounds of formula II, R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or

30



In yet another embodiment of the compounds of formula II, R<sup>1</sup> is

35




5

wherein Q is O or S, and each occurrence of R<sup>7</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, halogen, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>, or adjacent occurrences of R<sup>7</sup> can be taken together to form a bicyclic alkyl or aryl ring.

In still another embodiment of the compounds of formula II, R<sup>1</sup> is C(O)R<sup>3</sup> or C(O)OR<sup>4</sup>.

15 In a further embodiment, the invention encompasses compounds of Formula III:



20

III

wherein:

25

R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(S)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(O)NHR<sup>3</sup>, C(S)NHR<sup>3</sup>, C(O)NR<sup>3</sup>R<sup>3'</sup>, C(S)NR<sup>3</sup>R<sup>3'</sup> or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

30

R<sup>2</sup> is H or (C<sub>1</sub>-C<sub>8</sub>)alkyl;

R<sup>3</sup> and R<sup>3'</sup> are independently is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl,

35

(C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>,  
 (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;


R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl,  
 5 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

each occurrence of R<sup>6</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl,  
 10 benzyl, aryl, (C<sub>2</sub>-C<sub>5</sub>)heteroaryl, or (C<sub>0</sub>-C<sub>8</sub>)alkyl-C(O)O-R<sup>5</sup> or the R<sup>6</sup> groups can join to  
 form a heterocycloalkyl group; and


the \* represents a chiral-carbon center.

In yet another embodiment of the compounds of formula **III**, R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 15 CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or



20

In another embodiment of the compounds of formula **III**, R<sup>1</sup> is




wherein Q is O or S, and each occurrence of R<sup>7</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl,  
 (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, halogen,  
 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl,  
 30 (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or  
 C(O)OR<sup>5</sup>, or adjacent occurrences of R<sup>7</sup> can be taken together to form a bicyclic alkyl or aryl  
 ring..

In still another embodiment of the compounds of formula **III**, R<sup>1</sup> is C(O)R<sup>3</sup> or  
 C(O)OR<sup>4</sup>.

35 In a further embodiment still, the invention encompasses compounds of Formula **IV**:



5



IV

wherein:

10

R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(S)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(O)NHR<sup>3</sup>, C(S)NHR<sup>3</sup>, C(O)NR<sup>3</sup>R<sup>3'</sup>, C(S)NR<sup>3</sup>R<sup>3'</sup> or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

15

R<sup>2</sup> is H or (C<sub>1</sub>-C<sub>8</sub>)alkyl;

R<sup>3</sup> and R<sup>3'</sup> are independently (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;

20

R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

25

R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

each occurrence of R<sup>6</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>2</sub>-C<sub>5</sub>)heteroaryl, or (C<sub>0</sub>-C<sub>8</sub>)alkyl-C(O)O-R<sup>5</sup> or the R<sup>6</sup> groups can join to


30

form a heterocycloalkyl group; and

the \* represents a chiral-carbon center.

In another embodiment of the compounds of formula IV, R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>4</sub>)alkyl, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or


35



5

In yet another embodiment of the compounds of formula IV, R<sup>1</sup> is

10




wherein Q is O or S, and each occurrence of R<sup>7</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, halogen, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>, or adjacent occurrences of R<sup>7</sup> can be taken together to form a bicyclic alkyl or aryl ring.

In still another embodiment of the compounds of formula IV, R<sup>1</sup> is C(O)R<sup>3</sup> or C(O)OR<sup>4</sup>.

In yet another embodiment, the invention encompasses compounds of Formula V:

25



V

30 wherein:

R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(S)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(O)NHR<sup>3</sup>, C(S)NHR<sup>3</sup>, C(O)NR<sup>3</sup>R<sup>3'</sup>, C(S)NR<sup>3</sup>R<sup>3'</sup> or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

- R<sup>3</sup> and R<sup>3'</sup> are independently (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>,  
 5 (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;

R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

- 10 R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

each occurrence of R<sup>6</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>2</sub>-C<sub>5</sub>)heteroaryl, or (C<sub>0</sub>-C<sub>8</sub>)alkyl-C(O)O-R<sup>5</sup> or the R<sup>6</sup> groups can join to form a heterocycloalkyl group; and

15


the \* represents a chiral-carbon center.

- In a separate embodiment of compounds of formula V, R<sup>1</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>,  
 20 (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>; and

- R<sup>3</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>5</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>; (C<sub>0</sub>-C<sub>8</sub>)alkyl-NH-C(O)O-R<sup>5</sup>; (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>,  
 25 (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>; and the other variables have the definitions above.


In another embodiment of the compounds of formula V, R<sup>1</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl or benzyl, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or

30



In another embodiment of the compounds of formula V, R<sup>1</sup> is

35




- 5 wherein Q is O or S, and each occurrence of  $R^7$  is independently  $H$ ,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl
- 10 ring.

In still another embodiment of the compounds of formula V,  $R^1$  is  $C(O)R^3$  or  $C(O)OR^4$ .

In another embodiment of the compounds of formula V,  $R^3$  is  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_1-C_8)$ alkyl, aryl, or  $(C_0-C_4)$ alkyl- $OR^5$ .

- 15 In yet another embodiment of the compounds of formula V, heteroaryl is pyridyl, furyl, or thienyl.

In another embodiment, the invention further provides compounds of Formula VI:



- 25 VI

wherein:

- $R^1$  is  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(S)NHR^3$ , or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;
- 30

$R^3$  is  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,

35

(C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;

R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl,  
 5 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;


each occurrence of R<sup>6</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl,  
 10 benzyl, aryl, (C<sub>2</sub>-C<sub>5</sub>)heteroaryl, or (C<sub>0</sub>-C<sub>8</sub>)alkyl-C(O)O-R<sup>5</sup> or the R<sup>6</sup> groups can join to form a heterocycloalkyl group; and

the \* represents a chiral-carbon center.

In a separate embodiment of compounds of formula VI, R<sup>1</sup> is (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
 15 (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(S)NHR<sup>3</sup>, or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>; and

R<sup>3</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl,  
 20 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>5</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>; (C<sub>0</sub>-C<sub>8</sub>)alkyl-NH-C(O)O-R<sup>5</sup>; (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>; and the other variables are defined as above.


In another embodiment of the compounds of formula VI, R<sup>1</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, benzyl,  
 25 CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or



30

In another embodiment of the compounds of formula VI, R<sup>1</sup> is

35




wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl ring.

In still another embodiment of the compounds of formula VI,  $R^1$  is  $C(O)R^3$  or  $C(O)OR^4$ .

In another embodiment of the compounds of formula VI,  $R^3$  is  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_1-C_8)$ alkyl, aryl, or  $(C_0-C_4)$ alkyl- $OR^5$ .

In yet another embodiment of the compounds of formula VI, heteroaryl is pyridyl, furyl, or thienyl. In another embodiment of the compounds of formula VI, the H of  $C(O)NHC(O)$  can be replaced with  $(C_1-C_4)$ alkyl, aryl, or benzyl.

In still another embodiment, the invention encompasses compounds of Formula VII:



VII

wherein:

$R^1$  is  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(S)NHR^3$ , or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;

$R^3$  is  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ ;

$R^4$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  $(C_1-C_4)$ alkyl-OR<sup>5</sup>, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl, or  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl;

$R^5$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, or  $(C_2-C_5)$ heteroaryl;

5

each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl-C(O)O-R<sup>5</sup> or the  $R^6$  groups can join to form a heterocycloalkyl group; and

10 the \* represents a chiral-carbon center.

In a separate embodiment of compounds of formula VII,  $R^1$  is  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl, C(O)R<sup>3</sup>, C(O)OR<sup>4</sup>,  $(C_1-C_8)$ alkyl-N(R<sup>6</sup>)<sub>2</sub>,  $(C_1-C_8)$ alkyl-OR<sup>5</sup>,  $(C_1-C_8)$ alkyl-C(O)OR<sup>5</sup>, C(S)NHR<sup>3</sup>, or  $(C_1-C_8)$ alkyl-O(CO)R<sup>5</sup>; and


15

$R^3$  is  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_5-C_8)$ alkyl-N(R<sup>6</sup>)<sub>2</sub>,  $(C_0-C_8)$ alkyl-NH-C(O)O-R<sup>5</sup>;  $(C_1-C_8)$ alkyl-OR<sup>5</sup>,  $(C_1-C_8)$ alkyl-C(O)OR<sup>5</sup>,  $(C_1-C_8)$ alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>; and the other variables are as

20 defined above.


In another embodiment of the compounds of formula VII  $R^1$  is  $(C_1-C_8)$ alkyl, benzyl, CH<sub>2</sub>OCH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, or

25



In another embodiment of the compounds of formula VII,  $R^1$  is

30



wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,

35

(C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>, or adjacent occurrences of R<sup>7</sup> can be taken together to form a bicyclic alkyl or aryl ring.

5 In still another embodiment of the compounds of formula **VII**, R<sup>1</sup> is C(O)R<sup>3</sup> or C(O)OR<sup>4</sup>.

In another embodiment of the compounds of formula **VII**, R<sup>3</sup> is (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>1</sub>-C<sub>8</sub>)alkyl, aryl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>. In yet another embodiment of the compounds of formula **VII**, heteroaryl is pyridyl, furyl, or thienyl.

10 As used herein, the phrase "compounds of the invention" means, collectively, compounds falling within Formulas **I**, **II**, **III**, **IV**, **V**, **VI**, and **VII** and include pharmaceutically acceptable salts, hydrates, solvates, and clathrates thereof.

The compounds of the invention generally exist in solid form and can be recrystallized according to well-known methods affording high-purity crystals, preferably,  
15 in greater than 95% purity, more preferably, in greater than 98% purity. Narrow melting-point range is an indication of purity, thus, compounds of the invention generally have a melting point within a range of 3 °C to 4 °C, more preferably, within a range of 2 °C.

The compounds of the invention can contain one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (*i.e.*,  
20 geometric isomers), enantiomers, or diastereomers. According to the invention, the chemical structures depicted herein, and therefore the compounds of the invention, encompass all of the corresponding enantiomers and stereoisomers, that is, both the stereomerically pure form (*e.g.*, geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric and stereoisomeric mixtures, *e.g.*, racemates.

25 A compound of the invention is considered optically active or enantiomerically pure (*i.e.*, substantially the R-form or substantially the S-form) with respect to a chiral center when the compound is about 90% ee (enantiomeric excess) or greater, preferably, equal to or greater than 95% ee with respect to a particular chiral center. A compound of the invention is considered to be in enantiomerically enriched form when the compound has an  
30 enantiomeric excess of greater than about 1% ee, preferably greater than about 5% ee, more preferably, greater than about 10% ee with respect to a particular chiral center. As used herein, a racemic mixture means about 50% of one enantiomer and about 50% of is corresponding enantiomer relative to all chiral centers in the molecule. Thus, the invention encompasses all enantiomerically pure, enantiomerically enriched, and racemic mixtures of  
35 compounds of Formulas **I** through **VII**.



Enantiomeric and stereoisomeric mixtures of compounds of the invention can be resolved into their component enantiomers or stereoisomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a  
5 chiral solvent. Enantiomers and stereoisomers can also be obtained from stereomerically or enantiomerically pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.

The invention further encompasses prodrugs of compounds falling within Formulas I, II, III, IV, V, VI, and VII. The term "prodrug" refers to a compound that, following  
10 administration in a mammal, converts, *via* a biotransformation, into a compound falling within Formulas I, II, III, IV, V, VI, and VII *in vivo*. Prodrugs of compounds falling within Formulas I, II, III, IV, V, VI, and VII can be synthesized using well-known methods, such as those described by Burger's Medicinal Chemistry and Drug Chemistry, Fifth Ed., Vol. 1, pp. 172-178, 949-982 (1995).

15 The compounds of the invention are defined herein by their chemical structures and/or chemical names. Where a compound is referred to by both a chemical structure and a chemical name, and the chemical structure and chemical name conflict, the chemical structure is determinative of the compound's identity.

In another embodiment, the present invention further provides pharmaceutical  
20 compositions comprising a therapeutically effective or a prophylactically effective amount of one or more compounds of the invention and a pharmaceutically acceptable vehicle or carrier. A pharmaceutically acceptable vehicle or carrier can comprise an excipient, diluent, or a mixture thereof. The term "therapeutically effective amount" means the amount of a compound of the invention that will elicit the biological or medical response in a mammal  
25 that is being treated by the veterinarian or clinician. The term "prophylactically effective" means the amount of a compound of the invention that will prevent or inhibit affliction or mitigate affliction of a mammal with a medical condition that a veterinarian or clinician is trying to prevent, inhibit, or mitigate.

In another embodiment, the invention concerns a method of modulating the  
30 production or lowering the levels of TNF- $\alpha$  in a mammal comprising administering to said mammal an effective amount of a compound of the invention.

In yet another embodiment, the invention concerns a method of modulating the production or lowering the levels of IL-1 $\beta$  in a mammal comprising administering to said mammal an effective amount of a compound of the invention.

35

In a further embodiment, the invention concerns a method of modulating the production or increasing the levels of IL-10 in a mammal comprising administering to said mammal an effective amount of a compound of the invention.

5 In still another embodiment, the invention concerns a method of modulating the production or increasing the levels of T-cells in a mammal comprising administering to said mammal an effective amount of a compound of the invention.

In still another embodiment, the invention concerns a method of treating or preventing cancer in a mammal, comprising administering to said mammal a therapeutically effective amount of a compound of the invention. The compounds of the invention can be  
10 used to treat or prevent any cancer, for example, solid tumors and blood-born tumors. Specific examples of cancers treatable or preventable by compounds of the invention include, but are not limited to, cancers of the skin, such as melanoma; lymph node; breast; cervix; uterus; gastrointestinal tract; lung; ovary; prostate; colon; rectum; mouth; brain; head and neck; throat; testes; kidney; pancreas; bone; spleen; liver; bladder; larynx; nasal  
15 passages; and AIDS-related cancers. The compounds are particularly useful for treating cancers of the blood and bone marrow, such as multiple myeloma and acute and chronic leukemias, for example, lymphoblastic, myelogenous, lymphocytic, and myelocytic leukemias. The compounds of the invention can be used for treating or preventing either primary or metastatic tumors.

20 In yet one more embodiment, the invention provides methods of treating or preventing cancer in a mammal, comprising administering to a mammal in need thereof, a therapeutically effective amount of a compound of the invention and another cancer chemotherapeutic agent.

In yet another embodiment, the invention concerns a method of treating or  
25 preventing inflammatory disorders in a mammal, comprising administering to said mammal a therapeutically effective amount of a compound of the invention. The compounds of the invention are especially effective to treat or prevent inflammatory diseases related to the up-regulation of TNF- $\alpha$  including, but not limited to, arthritic conditions, such as, rheumatoid arthritis, and osteoarthritis; rheumatoid spondylitis; psoriasis; post ischemic perfusion  
30 injury; inflammatory bowel disease; and chronic inflammatory pulmonary disease.

In one more embodiment still, the invention provides methods of treating or preventing inflammatory disorders in a mammal, comprising administering to a mammal in need thereof, a therapeutically effective amount of a compound of the invention and another anti-inflammatory agent.

35

In a further embodiment, the invention concerns a method of treating or preventing heart disease in a mammal comprising administering to said mammal a therapeutically effective amount of a compound of the invention. For example, the compounds of the invention can be used to treat or prevent congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, and myocardial infarction.

In an additional embodiment, the invention concerns a method of treating or preventing osteoporosis in a mammal comprising administering to said mammal a therapeutically effective amount of a compound of the invention.

10 In a further embodiment, the invention relates to a method of treating or preventing viral, genetic, inflammatory, allergic, and autoimmune diseases. For example, the compounds are useful to treat or prevent diseases including, but not limited to, HIV, hepatitis, adult respiratory distress syndrome, bone resorption diseases, chronic pulmonary inflammatory diseases, dermatitis, cystic fibrosis, septic shock, sepsis, endotoxic shock, hemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, meningitis, psoriasis, fibrotic disease, cachexia, graft rejection, auto-immune disease, rheumatoid spondylitis, Crohn's disease, ulcerative colitis, inflammatory-bowel disease, multiple sclerosis, systemic lupus erythematosus, ENL in leprosy, radiation damage, asthma, or hyperoxic alveolar injury in a mammal comprising administering to said mammal a therapeutically effective amount of a compound of the invention.

In still another embodiment, the invention concerns a method of treating or preventing malaria, mycobacterial infection, or an opportunistic infection resulting from HIV in a mammal, comprising administering to said mammal a therapeutically effective amount of a compound of the invention.

25 In still one more embodiment, the invention relates to treating or preventing mammals having more than one of the conditions treatable by a compound of the invention.

In the above embodiments, it is preferable that the mammal be in need of the treatment or prevention, that is, the mammal is actually suffering from a medical condition or at risk of a medical condition for which a compound of the invention can provide treatment or prevention. However, the compounds of the invention can also be administered to test animals that do not necessarily require such treatment or prevention.

In a further embodiment, the invention encompasses a method of modulating the production or lowering the levels of TNF- $\alpha$  in a mammalian cell or tissue comprising contacting an effective amount of a compound of the invention with said mammalian cell or tissue.

In yet another embodiment, the invention encompasses a method of modulating the production or lowering the levels of IL-1 $\beta$  in a mammalian cell or tissue comprising contacting an effective amount of a compound of the invention with said mammalian cell or tissue.

5 In still another embodiment, the invention encompasses a method of modulating the production or lowering the levels of IL-10 in a mammalian cell or tissue comprising contacting an effective amount of a compound of the invention with said mammalian cell or tissue.

10 In still another embodiment, the invention encompasses a method of modulating the production or lowering the levels of T-cells in a mammalian cell or tissue comprising contacting an effective amount of a compound of the invention with said mammalian cell or tissue.

In these embodiments, the term "effective amount" means the amount of the compound that will induce the biological response sought by the researcher, veterinarian, 15 physician, or clinician. It should be understood that the cell can be in a cell culture or a tissue culture (*in vitro*) or in an organism (*in vivo*) including a human.

The present invention may be understood by reference to the detailed description and examples that are intended to exemplify non-limiting embodiments of the invention.

20

## 5. Definitions

The phrase "pharmaceutically acceptable salt(s)," as used herein includes, but is not limited to, salts of acidic or basic groups that may be present in the compounds of the invention. Compounds of the invention that are basic in nature are capable of forming a 25 wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable salts of such basic compounds are those that form salts comprising pharmacologically acceptable anions including, but not limited to, acetate, benzenesulfonate, benzoate, bicarbonate, bitartrate, bromide, calcium edetate, camsylate, carbonate, chloride, bromide, iodide, citrate, dihydrochloride, edetate, edisylate, estolate, 30 esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydroxynaphthoate, isethionate, lactate, lactobionate, malate, maleate, mandelate, mesylate, methylsulfate, muscate, napsylate, nitrate, panthothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, succinate, sulfate, tannate, tartrate, teclate, triethiodide, and pamoate (*i.e.*, 1,1'-methylene-*bis*-(2-hydroxy-3- 35 naphthoate)). Compounds of the invention that include an amino moiety also can form

pharmaceutically acceptable salts with various amino acids, in addition to the acids mentioned above. Compounds of the invention that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include alkali metal or alkaline earth metal salts and, particularly, calcium, magnesium, sodium, lithium, zinc, potassium, and iron salts.

As used herein, the term "solvate" means a compound of the invention or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of a solvent bound by non-covalent intermolecular forces. Preferred solvents are volatile, non-toxic, and/or acceptable for administration to humans in trace amounts.

As used herein, the term "hydrate" means a compound of the invention or a salt thereof, that further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.

The term "clathrate" means a compound of the invention or a salt thereof in the form of a crystal lattice that contains spaces (*e.g.*, channels) that have a guest molecule (*e.g.*, a solvent or water) trapped within.

As used herein, the term "alkyl group" means a saturated, monovalent, unbranched or branched hydrocarbon chain. Examples of alkyl groups include, but are not limited to, (C<sub>1</sub>-C<sub>8</sub>)alkyl groups, such as methyl, ethyl, propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, *t*-butyl, pentyl, isopentyl, neopentyl, and hexyl, heptyl, and octyl. An alkyl group can be unsubstituted or substituted with one or two suitable substituents.

An "alkenyl group" means a monovalent, unbranched or branched hydrocarbon chain having one or more double bonds therein. The double bond of an alkenyl group can be unconjugated or conjugated to another unsaturated group. Suitable alkenyl groups include, but are not limited to (C<sub>2</sub>-C<sub>8</sub>)alkenyl groups, such as vinyl, allyl, butenyl, pentenyl, hexenyl, butadienyl, pentadienyl, hexadienyl, 2-ethylhexenyl, 2-propyl-2-butenyl, 4-(2-methyl-3-butene)-pentenyl. An alkenyl group can be unsubstituted or substituted with one or two suitable substituents.

An "alkynyl group" means monovalent, unbranched or branched hydrocarbon chain having one or more triple bonds therein. The triple bond of an alkynyl group can be unconjugated or conjugated to another unsaturated group. Suitable alkynyl groups include, but are not limited to, (C<sub>2</sub>-C<sub>8</sub>)alkynyl groups, such as ethynyl, propynyl, butynyl, pentynyl,

hexynyl, methylpropynyl, 4-methyl-1-butynyl, 4-propyl-2-pentynyl, and 4-butyl-2-hexynyl. An alkynyl group can be unsubstituted or substituted with one or two suitable substituents.

An "aryl group" means a monocyclic or polycyclic-aromatic group comprising carbon and hydrogen atoms. Examples of suitable aryl groups include, but are not limited to, phenyl, tolyl, anthacenyl, fluorenyl, indenyl, azulenyl, and naphthyl, as well as benzo-fused carbocyclic moieties such as 5,6,7,8-tetrahydronaphthyl. An aryl group can be unsubstituted or substituted with one or more suitable substituents. Preferably, the aryl group is a monocyclic ring, wherein the ring comprises 6 carbon atoms, referred to herein as "(C<sub>6</sub>)aryl".

10 A "heteroaryl group" means a monocyclic or polycyclic aromatic ring comprising carbon atoms and one or more heteroatoms, preferably, 1 to 3 heteroatoms, independently selected from nitrogen, oxygen, and sulfur. Preferred heteroaryl-ring systems include 5 to 6 membered monocyclic, 8 to 11 membered bicyclic, and 11 to 15 membered tricyclic ring systems. As well known to those skilled in the art, heteroaryl rings have less aromatic  
15 character than their all-carbon counter parts. Thus, for the purposes of the invention, a heteroaryl group need only have some degree of aromatic character. Illustrative examples of heteroaryl groups include, but are not limited to, pyridinyl, pyridazinyl, pyrimidyl, pyrazyl, triazinyl, pyrrolyl, pyrazolyl, imidazolyl, (1,2,3,- and (1,2,4)-triazolyl, pyrazinyl, pyrimidinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, phienyl, isoxazolyl, and  
20 oxazolyl. A heteroaryl group can be unsubstituted or substituted with one or two suitable substituents. Preferably, a heteroaryl group is a 5 or 6 membered monocyclic ring, wherein the ring comprises 2 to 5 carbon atoms and 1 to 3 heteroatoms, referred to herein as "(C<sub>2</sub>-C<sub>5</sub>)heteroaryl", which optionally can be fused to one or more other aryl, cycloalkyl, heteroaryl, or heterocyclic ring systems to form 7 to 10 membered bicyclic or 10 to 15  
25 membered tricyclic ring systems.

A "cycloalkyl group" means a non-aromatic, monocyclic or polycyclic ring comprising carbon and hydrogen atoms. A cycloalkyl group can have one or more carbon-carbon double bonds in the ring so long as the ring is not rendered aromatic by their presence. Examples of cycloalkyl groups include, but are not limited to, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl  
30 groups, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, and saturated cyclic and bicyclic terpenes and (C<sub>3</sub>-C<sub>8</sub>)cycloalkenyl groups, such as cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, and cycloheptenyl, and unsaturated cyclic and bicyclic terpenes. A cycloalkyl group can be unsubstituted or substituted by one or two suitable substituents. Preferably, the cycloalkyl group is a  
35 monocyclic ring or bicyclic ring.

A "heterocycloalkyl group" means a non-aromatic monocyclic or polycyclic ring comprising carbon atoms and at least one heteroatom, preferably, 1 to 4 heteroatoms independently selected from nitrogen, oxygen, and sulfur. Preferred heterocyclic-ring systems include 3 to 8 membered monocyclic, 8 to 11 membered bicyclic, and 11 to 15 membered tricyclic ring systems. A heterocycloalkyl group can have one or more carbon-carbon double bonds or carbon-heteroatom double bonds in the ring as long as the ring is not rendered aromatic by their presence. Examples of heterocycloalkyl groups include aziridinyl, pyrrolidinyl, pyrrolidino, piperidinyl, piperidino, piperazinyl, piperazino, morpholinyl, morpholino, thiomorpholinyl, thiomorpholino, tetrahydrofuranyl, tetrahydrothiofuranyl, tetrahydropyranyl, and pyranyl. A heterocycloalkyl group can be unsubstituted or substituted with one or more suitable substituents. Preferably, the heterocycloalkyl group is a monocyclic or bicyclic ring, more preferably, a 3-7 membered monocyclic ring, wherein the ring comprises from 1 to 6 carbon atoms and from 1 to 3 heteroatoms, referred to herein as "(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl", which optionally can be fused to one or more other aryl, cycloalkyl, heteroaryl, or heterocyclic ring systems to form 7 to 10 membered bicyclic or 10 to 15 membered tricyclic ring systems.

The term "alkoxy group" means an -O-alkyl group, wherein alkyl is as defined above. An alkoxy group can be unsubstituted or substituted with one or two suitable substituents. Preferably, the alkyl chain of an alkoxy group is from 1 to 8 carbon atoms in length, referred to herein as "(C<sub>1</sub>-C<sub>8</sub>)alkoxy".

The term "aryloxy group" means an O-aryl group, wherein aryl is as defined above. An aryloxy group can be unsubstituted or substituted with one or two suitable substituents. Preferably, the aryl ring of an aryloxy group is a monocyclic ring, wherein the ring comprises 6 carbon atoms, referred to herein as "(C<sub>6</sub>)aryloxy".

The term "benzyl" means CH<sub>2</sub>-phenyl. A benzyl group can be unsubstituted or substituted with one or more suitable substituents.

The term "phenyl" means C<sub>6</sub>H<sub>5</sub>. A phenyl group can be unsubstituted or substituted with one or more suitable substituents.

A "carbonyl" group is a divalent group of the formula -C(O)-.

An "alkoxycarbonyl" group means a monovalent group of the formula C(O)-alkoxy. Preferably, the hydrocarbon chain of an alkoxycarbonyl group is from 1 to 8 carbon atoms in length, referred to herein as a "lower alkoxycarbonyl" group.

As used herein, "halogen" means fluorine, chlorine, bromine, or iodine. Correspondingly, the meaning of the term "halo" encompass fluoro, chloro, bromo, and iodo.

As used herein, a "suitable substituent" means a group that does not nullify the synthetic or pharmaceutical utility of the compounds of the invention or the intermediates useful for preparing them. Examples of suitable substituents include, but are not limited to:


- (C<sub>1</sub>-C<sub>8</sub>)alkyl; (C<sub>1</sub>-C<sub>8</sub>)alkenyl; (C<sub>1</sub>-C<sub>8</sub>)alkynyl; aryl; (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;  
 5 (C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl; (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl; O-(C<sub>1</sub>-C<sub>8</sub>)alkyl; O-(C<sub>1</sub>-C<sub>8</sub>)alkenyl;  
 O-(C<sub>1</sub>-C<sub>8</sub>)alkynyl; O-aryl; CN; OH; oxo; halo, C(O)OH; COhalo; O(CO)halo; CF<sub>3</sub>, N<sub>3</sub>;  
 NO<sub>2</sub>, NH<sub>2</sub>; NH((C<sub>1</sub>-C<sub>8</sub>)alkyl); N((C<sub>1</sub>-C<sub>8</sub>)alkyl)<sub>2</sub>; NH(aryl); N(aryl)<sub>2</sub>; (CO)NH<sub>2</sub>;  
 (CO)NH((C<sub>1</sub>-C<sub>8</sub>)alkyl); (CO)N((C<sub>1</sub>-C<sub>8</sub>)alkyl)<sub>2</sub>; (CO)NH(aryl); (CO)N(aryl)<sub>2</sub>; O(CO)NH<sub>2</sub>;  
 NHOH; NOH((C<sub>1</sub>-C<sub>8</sub>)alkyl); NOH(aryl); O(CO)NH((C<sub>1</sub>-C<sub>8</sub>)alkyl); O(CO)N((C<sub>1</sub>-C<sub>8</sub>)alkyl)<sub>2</sub>;  
 10 O(CO)NH(aryl); O(CO)N(aryl)<sub>2</sub>; CHO; CO((C<sub>1</sub>-C<sub>8</sub>)alkyl); CO(aryl); C(O)O((C<sub>1</sub>-C<sub>8</sub>)alkyl);  
 C(O)O(aryl); O(CO)((C<sub>1</sub>-C<sub>8</sub>)alkyl); O(CO)(aryl); O(CO)O((C<sub>1</sub>-C<sub>8</sub>)alkyl); O(CO)O(aryl);  
 S-(C<sub>1</sub>-C<sub>8</sub>)alkyl; S-(C<sub>1</sub>-C<sub>8</sub>)alkenyl; S-(C<sub>1</sub>-C<sub>8</sub>)alkynyl; and S-aryl. One of skill in art can  
 readily choose a suitable substituent based on the stability and pharmacological and  
 synthetic activity of the compound of the invention.

15

## 6. Detailed Description of The Invention

In one embodiment, the invention encompasses compounds of the formula:

20



- 25 or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH<sub>2</sub> or C=O;

- 30 R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(S)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(O)NHR<sup>3</sup>, C(S)NHR<sup>3</sup>, C(O)NR<sup>3</sup>R<sup>3'</sup>, C(S)NR<sup>3</sup>R<sup>3'</sup> or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

- 35 R<sup>2</sup> is H, F, benzyl, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, or (C<sub>2</sub>-C<sub>8</sub>)alkynyl;



- R<sup>3</sup> and R<sup>3'</sup> are independently (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>,  
 5 (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;

R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

- 10 R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

each occurrence of R<sup>6</sup> is independently H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>2</sub>-C<sub>5</sub>)heteroaryl, or (C<sub>0</sub>-C<sub>8</sub>)alkyl-C(O)O-R<sup>5</sup> or the R<sup>6</sup> groups can join to form a heterocycloalkyl group;

- 15 n is 0 or 1; and

the \* represents a chiral-carbon center;

with the proviso that when n is 0 then R<sup>1</sup> is not H.

In a separate embodiment of compounds of formula I, when n is 0 then R<sup>1</sup> is

- (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl,  
 20 (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

R<sup>2</sup> is H or (C<sub>1</sub>-C<sub>8</sub>)alkyl; and

25





R<sup>3</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>5</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>; (C<sub>0</sub>-C<sub>8</sub>)alkyl-NH-C(O)O-R<sup>5</sup>; (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>; and the other variables have


- 30 the same definitions. Further, the compounds encompassed by Formulas II, III, IV, V, VI, and VII as described above are also included within the invention.

A few examples of compounds of the invention which are illustrative and non-limiting are depicted in Table 1 below.

35


TABLE 1: Examples of Compounds of the Invention

	Structure	Name
5		[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl-methyl]-carbamic acid <i>tert</i> -butyl ester
10	<b>I-1</b>	
15		4-(Aminomethyl)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione
	<b>I-2</b>	
20		<i>N</i> -[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl-methyl]-acetamide
25	<b>I-3</b>	
30		<i>N</i> -[[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]cyclopropyl-carboxamide
	<b>I-4</b>	
35		




[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-indol-4-ylmethyl]-carbamic acid ethyl ester

I-5




2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-indol-4-ylmethyl]-carbamic acid benzyl ester

I-6



2-Chloro-N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}acetamide

I-7




2-(Dimethylamino)-N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-acetamide

I-8


35

5

**I-9**


1-*tert*-Butyl-3-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-ylmethyl]-urea

10

**I-10**

*N*-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-3,3-dimethylbutanamide


15

**I-11**

*N*-[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-3-pyridylcarboxamide

20


25

**I-12**


3-{1-Oxo-4-[benzylamino]isoindolin-2-yl}piperidine-2,6-dione

30


35

**I-13**


2-(2,6-Dioxo(3-piperidyl))-4-[benzylamino]isoindoline-1,3-dione

**I-14**

*N*-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}propanamide

**I-15**

*N*-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-3-pyridylcarboxamide


**I-16**

*N*-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}heptanamide

30


35

5

**I-17**


N-([2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl)-2-furlylcarboxamide

10

**I-18**


2-Azido-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isindol-4-ylmethyl]-acetamide

15

**I-19**

2-Amino-N-([2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl)acetamide

20

**I-20**


Ethyl 6-(N-([2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl)carbamoyl)hexanoate

25

30


35

5

**I-21**


3-[(*tert*-Butoxy)carbonylamino]-*N*-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]propanamide

10

**I-22**


3-Amino-*N*-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]propanamide

15

**I-23**

*N*-[[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]-2-thienylcarboxamide

20

**I-24**


*N*-[[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]-2-methoxyacetamide

25

30

35


5



**I-25**

(*N*-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)methyl acetate


10



**I-26**

Ethyl 2-[(*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)amino]acetate

15




**I-27**

*N*-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(ethylamino)carboxamide

20

25




**I-28**

2-(2,6-Dioxo(3-piperidyl))-4-[(2-furylmethyl)amino]isoindoline-1,3-dione

30


35






*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]-2-methoxyacetamide

**I-29**



*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]heptanamide

**I-30**




{*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]carbamoyl}methyl acetate

**I-31**

30


35



*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]pentanamide

**I-32**


10



*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]-2-thienylcarboxamide

**I-33**

20




Methyl {*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]carbamoyl} formate

**I-34**


30

35




*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-furylcarboxamide

**I-35**



*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]benzamide


**I-36**



*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]propanamide

**I-37**


5



Methyl 3-{*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]carbamoyl}propanoate

**I-38**

10




*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]-2-phenylacetamide

15

**I-39**

20



*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]-2-pyridylcarboxamide


25

**I-40**

30

35


5



*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]-2-chloroacetamide

**I-41**

10




2-azido-*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]acetamide

15

**I-42**

20



2-Amino-*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]acetamide


25

**I-43**

30

35


5



*N*-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]-2-chloroacetamide

**I-44**


10



2-azido-*N*-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]acetamide

**I-45**

20




2-Amino-*N*-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]acetamide

**I-46**

25

30


35



3-{4-[(2-Furylmethyl)amino]-1-oxoisindolin-2-yl}piperidine-2,6-dione

I-47


10



3-[1-Oxo-4-(pentylamino)isoindolin-2-yl]piperidine-2,6-dione

I-48

20




2-(2,6-Dioxo-piperidin-3-yl)-4-(2-methoxyethylamino)-isoindole-1,3-dione

I-49


30

35




2-Benzyloxy-N-[2-(2,6-dioxo-piperidin-3-yl)-  
1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-  
acetamide

**I-50**



2-(2,6-Dioxo-piperidin-3-yl)-4-pentylamino-  
isoindole-1,3-dione


**I-51**



3-Chloro-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-  
dioxo-2,3-dihydro-1H-isoindol-4-yl]-  
benzamide

**I-52**

25




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-  
dihydro-1H-isoindol-4-yl]-2-phenoxy-  
acetamide

**I-53**


35





4-(2-Benzyloxy-ethylamino)-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione


**I-54**



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-fluoro-benzamide

15


**I-55**



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methyl-benzamide

**I-56**


25



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methoxy-benzamide


**I-57**

35



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-trifluoromethyl-benzamide


**I-58**



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-nitro-benzamide

15


**I-59**



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-butyramide

25



**I-60**




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methylamino-acetamide

35

**I-61**

**I-62**

15


**I-63**

25

**I-64**


30

35



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-fluoro-benzamide


**I-65**



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-trifluoromethyl-benzamide

15

**I-66**




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-methyl-benzamide

25

**I-67**

30


35



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-nitro-benzamide

**I-68**


10



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-ethoxy-acetamide

**I-69**

20




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methylsulfanyl-acetamide

**I-70**

30


35



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methoxy-benzamide

**I-71**


10



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-fluoro-benzamide


**I-72**

20




7-Amino-N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl} heptanamide

30




N- {[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl} butanamide



N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}benzamide


**I-75**

10



N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}phenylacetamide

20




N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-2-pyridylcarboxamide

25

**I-77**

30




N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}undecamide

**I-78**


35

5



N-{{2-[(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-2-methylpropanamide

10




N-{{2-[(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}cyclopentylcarboxamide

15

**I-80**

20




N-{{2-[(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}cyclohexylcarboxamide

25

**I-81**

30







N-{{2-[(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(phenylamino)carboxamide



**I-82**

35





TABLE 1 Cont.





	Structure	Name
5		N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(butylamino)carboxamide
10	<b>I-83</b>	
15		N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(propylamino)carboxamide
20	<b>I-84</b>	
25		N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(cyclohexylamino)carboxamide
30		N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}[(methylethylamino)]carboxamide
35	<b>I-86</b>	

**I-87**


15

**I-88****I-89****I-90**

35


**I-91****I-92****I-93****I-94**

35




N-[2-(2,6-Dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-nicotinamide

**I-95**




2-(2,6-Dioxo-piperidin-3-yl)-4-[[[(furan-2-ylmethyl)-amino]-methyl]-isoindole-1,3-dione

**I-96**



N-[2-(2,6-Dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide


**I-97**



2-Dimethylamino-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-acetamide


**I-98**

35



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methyl-benzamide


**I-99**



Heptanoic acid  
[2-(2,6-dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-amide


**I-100**

15



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3,3-dimethyl-butyramide

**I-101**




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-isobutyramide

**I-102**

35


5



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-phenyl-propionamide

**I-103**


10



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-methoxy-benzamide

**I-104**

20




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-trifluoromethyl-benzamide

**I-105**

25

30




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-malonamic acid methyl ester

**I-106**

35


5



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methoxy-propionamide

**I-107**

10




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-hydroxy-acetamide

15

**I-108**

20




4-[(Furan-2-ylmethyl)-amino]-2-(1-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione

25

**I-109**

30




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-isonicotinamide

35

**I-110**

5




N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-acetamide

I-111

10

15




{5-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylcarbamoyl]-pentyl}-carbamate benzyl ester

20

I-112


25



2-(2,6-Dioxo(3-piperidyl))-4-(((cyclohexylamino)thioxomethyl)amino)methylisoindole-1,3-dione

I-113

30




2-(2,6-Dioxo(3-piperidyl))-4-(((ethylamino)thioxomethyl)amino)methylisoindole-1,3-dione

I-114


35





2-(2,6-Dioxo(3-piperidyl))-4-  
({{[(propylamino)thioxomethyl]amino}meth  
yl)isoindole-1,3-dione


I-115



N-[2-(2,6-dioxo(3-piperidyl))-1,3-  
dioxoisindolin-4-yl]-2-chloro-benzylamine

15


I-116



{5-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-  
2,3-dihydro-1H-isoindol-4-yl]carbamoyl]-  
pentyl}-carbamic acid benzyl ester

25


I-117




2-Methoxy-N-[2-(3-methyl-2,6-dioxo-  
piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-  
isoindol-4-yl]-acetamide

35

I-118




Pentanoic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide

**I-119**


Heptanoic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide

15

**I-120**

3-Chloro-N-[2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide


25

**I-121**

N-[2-(3-Methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-propionamide


35

**I-122**




Thiophene-2-carboxylic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide

I-123




2-(2,6-Dioxo-piperidin-3-yl)-4-[(5-methyl-furan-2-ylmethyl)-amino]-isoindole-1,3-dione

I-124



2-(2,6-Dioxo-piperidin-3-yl)-4-[(5-hydroxymethyl-furan-2-ylmethyl)-amino]-isoindole-1,3-dione


I-125



2-(2,6-Dioxo-piperidin-3-yl)-4-[(thiophen-2-ylmethyl)-amino]-isoindole-1,3-dione


I-126

35




4-(3-Chloro-benzylamino)-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione

**I-127**




2-(2,6-Dioxo-piperidin-3-yl)-4-[(pyridin-3-ylmethyl)-amino]-isoindole-1,3-dione

**I-128**



5-{{2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylamino}-methyl}-furan-2-carboxylic acid

**I-129**




4-[(4,5-Dimethyl-furan-2-ylmethyl)-amino]-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione

**I-130**

30

35


5



4-[(Benzofuran-2-ylmethyl)-amino]-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione

**I-131**

10




4-(3-Chloro-benzylamino)-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione

15

**I-132**


20



3-[4-(3-Chloro-benzylamino)-1-oxo-1,3-dihydro-isoindol-2-yl]-piperidine-2,6-dione

**I-133**

25




N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(cyclopentylamino)carboxamide

30


**I-134**

35



N-{{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(3-pyridylamino)carboxamide Hydrochloride


**I-135**



N-{{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}piperidylcarboxamide

15


**I-136**



Tert-Butyl 4-(N-{{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(carbamoyl)piperazinecarboxylate

25

**I-137**




N-{{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(diethylamino)carboxamide

**I-138**

35


5



Cyclopropyl-N-{{2-[(3-methyl-2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carboxamide

**I-139**

10




15

N-{{2-[(2,6-Dioxo(3-piperidyl))-1-oxoisindolin-4-yl]methyl}cyclopropylcarboxamide

**I-140**

20




25

N-{{2-[(2,6-Dioxo(3-piperidyl))-1-oxoisindolin-4-yl]methyl}(ethylamino)carboxamide

**I-141**

30

35



Piperazine-1-carboxylic acid  
[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-amide

# I-142

10

Selected compounds of Table 1 were tested using the *in vitro* assays described below and found to be active for modulating the production of TNF- $\alpha$ .





Examples of optically or enantiomerically pure stereoisomers of the invention are depicted in Table 2 below.



15

TABLE 2: Examples of Stereoisomers of the Invention



No.	Structure	No.	Structure
20			
(R)-I-1		(S)-I-1	
25			
30			
(R)-I-3		(S)-I-3	
35			











10



20



30




### 6.1 Synthesis of the Compounds of the Invention


The compounds of the invention can be obtained via standard, synthetic methodology. Some convenient methods are illustrated in Schemes 1-8. Starting materials useful for preparing the compounds of the invention and intermediates therefor, are  
 5 commercially available or can be prepared from commercially available materials using known synthetic methods and reagents. Such starting materials include, but are not limited to, methyl-2-(methoxycarbonyl)-3-nitrobenzoate; methyl-3-aminomethyl-2-(methoxycarbonyl)benzoate; substituted and unsubstituted aminoglutarimide hydrochloride; di-*t*-butyl dicarbonate; and cyclopropylcarbonyl chloride.

10


Scheme 1: Synthesis of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (Formula I, wherein R<sup>1</sup> is H and n is 1)



20




30



Scheme 1 outlines one method to synthesize 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (**I**, wherein  $R^1$  is H and  $n$  is 1) from compound **10**. In the first step, reduction of **10** (commercially available), for example, with palladium on charcoal and 50 psi of hydrogen, followed by standard isolation and purification gives arylamine **11**. Arylamine **11** converts to nitrile **12** by diazonium salt formation effected by treatment with sodium nitrate then displacement of nitrogen with cyanide according to the classic Sandmeyer procedure. Reduction of nitrile **12**, for example, with palladium on carbon in methanol/aqueous hydrochloric acid under an atmosphere of hydrogen, gives the hydrochloride salt of compound **13**. Treatment of **13** with triethylamine liberates the free base, which in turn reacts with di-*t*-butyl dicarbonate (**14**) (commercially available, for example, from Aldrich Chemical Co. Milwaukee, WI) giving carbamate **15**. Treatment of carbamate **15** with **16**, where  $R^2$  is as defined above, and a base, such as diisopropylethyl amine gives compound **17** that converts to **I**, wherein  $R^1$  is H and  $n$  is 1 upon standard hydrolysis, for example, with aqueous hydrochloric acid/dioxane. Compounds **16** can be obtained by cyclizing the appropriately substituted, amino-protected glutamine by well-known methods (*e.g.*, see WO 98/54170, incorporated herein by reference).

Scheme 2: Synthesis of 4-Amino-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione  
 (Formula **I**, wherein  $R^1$  is H and  $n$  is 0)

20




30

Scheme 2 outlines a convenient method to synthesize 4-amino-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione (**I**, wherein  $R^1$  is H and  $n$  is 0) from 4-nitrophthalic

35

anhydride (18). In the first step, a mixture of 18 and 16 in an acidic medium (*e.g.*, sodium acetate in glacial acetic acid) is heated at about 60 °C to about 150 °C for a time of about 1 hour to about 24 hours, until the reaction is substantially complete. After an aqueous workup, 19 is isolated and characterized according to standard methods (*see e.g.*, United States Patent No. 5,635,517, incorporated herein by reference). Alternately, the reaction may be carried out in other solvents, including pyridine. Conversion of 19 to I, wherein R<sup>1</sup> is H and n is 0, is accomplished by standard hydrogen reduction, for example, with palladium on carbon under about 50 psi to 200 psi of hydrogen at about room temperature to about 100 °C (*see e.g.*, the procedure recited in United States Patent 5,635,517).

Scheme 3: Synthesis of Compounds of Formula I, wherein R<sup>1</sup> is C(O)R<sup>3</sup> or C(O)OR<sup>4</sup>




Scheme 3 outlines a convenient synthesis of compounds of Formula I, wherein R<sup>1</sup> is C(O)R<sup>3</sup> or C(O)OR<sup>4</sup>. In the first step, compound 13, as prepared as in Scheme 1 above, reacts with a compounds 20 or 21, depending on whether R<sup>1</sup> of C(O)R<sup>3</sup> or C(O)OR<sup>4</sup> is desired, to give compounds 22. According to scheme 3, E is a suitable leaving group, for example, but not limited to halides, such as chloride, bromide, and iodide; azido (N<sub>3</sub>); arylsulfonyloxy or alkylsulfonyloxy (*e.g.*, tosyloxy or mesyloxy); phenoxy; alkoxy; and oxycarbonyl groups. Preferably, E is a halide, more preferably, chloride. Preferably,

compounds **20** are acid chlorides, such as acetyl chloride and cyclopropylcarbonyl chloride and compounds **21** are chloroformates, such as ethylchloroformate or benzylchloroformate. The reaction is carried out according to standard, well-known procedures, *e.g.* see March, J. *Advanced Organic Chemistry; Reactions Mechanisms, and Structure*, 4th ed., 1992, pp.417-419, incorporated herein by reference. Treatment of **22** with **16** using the same procedure as outlined in Scheme 1 gives compounds of Formula I, wherein  $R^1$  is  $C(O)R^3$  or  $C(O)OR^4$ .


**Scheme 4: Alternative Synthesis of Compounds of Formula I, wherein**

**$R^1$  is  $C(O)R^3$ , or  $C(O)OR^4$**




Scheme 4 outlines an alternative synthesis of compounds of Formula I, wherein  $R^1$  is  $C(O)R^3$  or  $C(O)OR^4$  and a convenient synthesis of compounds of Formula I, wherein  $R^1$  is  $CH_2R^3$ . In the first step, compounds I, wherein  $R^1$  is H, as prepared as in Scheme 1 ( $n = 1$ ) or Scheme 2 ( $n = 0$ ) above, reacts with a compounds **20**, **21**, or **23** depending on whether  $R^1$  of  $C(O)R^3$ ,  $C(O)OR^4$ , or  $CH_2R^3$  is desired, to give compounds I where  $R^1$  is  $C(O)R^3$ ,  $C(O)OR^4$ , or  $CH_2R^3$ . As defined above for Scheme 3, E is a suitable leaving group. Preferably, E is a halide, more preferably, chloride. Preferably, compounds **20** are acid chlorides, such as chloroacetyl chloride and *t*-butylacetyl chloride. Aldehydes **23** are readily available commercially or synthesized by well-known methods. The reaction of **20** or **21** with I, where  $R^1$  is H is carried out according to standard, well-known procedures for nucleophilic displacement, *e.g.* see March, J. *Advanced Organic Chemistry; Reactions Mechanisms, and Structure*, 4th ed., 1992, pp.417-419. The reaction of **23** with I, where  $R^1$  is H is accomplished according to the well-known reductive-amination procedure between an aldehyde and a primary amine, *e.g.* see March, J. *Advanced Organic Chemistry; Reactions Mechanisms, and Structure*, 4th ed., 1992, pp.898-902, incorporated herein by reference.

Scheme 5: Synthesis of Compounds of Formula I, wherein  $R^1$  is  $C(O)CH_2N(R^6)_2$



Scheme 5 outlines one method for the synthesis of compounds of Formula I, wherein  $R^1$  is  $C(O)CH_2N(R^6)_2$ . A compound of Formula I, wherein  $R^1$  is  $C(O)R^3$  and  $R^3$  is  $(CH_2)_nE$ , where E is a suitable leaving group as defined for Scheme 3, reacts with amines 24 to give the desired Formula I compound, where  $R^1$  is  $C(O)CH_2N(R^6)_2$ . Preferably, E is chloro and  $R^5$  is  $(C_1-C_8)$ alkyl, such as methyl. The reaction is performed according to standard, well-known procedures, e.g. see March, J. *Advanced Organic Chemistry; Reactions Mechanisms, and Structure*, 4th ed., 1992, pp. 411-413, incorporated herein by reference.


20 Scheme 6: Synthesis of Compounds of Formula I, wherein  $R^1$  is  $C(O)NHR^5$



30 Scheme 6 shows one method to synthesize compounds of Formula I, wherein  $R^1$  is  $C(O)NHR^5$ . A compound of Formula I, where  $R^1$  is H reacts with isocyanates 25 under routine conditions to give a compounds I, where  $R^1$  is  $C(O)NHR^5$ . The reaction is performed according to standard, well-known procedures, e.g. see March, J. *Advanced Organic Chemistry; Reactions Mechanisms, and Structure*, 4th ed., 1992, p. 903, incorporated herein by reference.

35

Scheme 7: Synthesis of Compounds of Formula I, wherein R<sup>2</sup> is F



20 Scheme 7 shows one method to synthesize compounds of Formula I, wherein R<sup>2</sup> is F. A similar method is described in U.S. Patent No. 5,874,448, incorporated herein by reference. In the first step, compounds of Formula I, where R<sup>1</sup> and R<sup>2</sup> are H are first protected with suitable nitrogen-protecting groups (PG<sup>1</sup>, PG<sup>2</sup>, and PG<sup>3</sup>) at the methyleneamino and glutarimide nitrogens, respectively, to give compounds 26. As used

25 herein, a "nitrogen protecting group" means a group that is reversibly attached to the nitrogen that renders the nitrogen moiety unreactive during a subsequent reaction(s) and that can be selectively cleaved to regenerate the original nitrogen moiety once its protecting purpose has been served. Examples of suitable protecting groups are found in Greene, T.W., *Protective Groups in Organic Synthesis*, 3rd edition 494-654 (1999), incorporated


30 herein by reference, and United States Patent No. 5,874,448. Preferably, the nitrogen-protecting group is stable in a basic reaction medium but can be cleaved by acid. Preferably, all of PG<sup>1</sup>, PG<sup>2</sup>, and PG<sup>3</sup> protecting groups are *tert*-butoxycarbonyl, attached by treatment of compounds of Formula I, where R<sup>1</sup> and R<sup>2</sup> are H with in excess of 3 equivalents of di-*tert*-butyl carbonate as described in United States Patent No. 5,874,448.

35 The fluorination-reaction procedure, to give compounds 27, is described in detail in United



States Patent No. 5,874,448 and can be effected by treating deprotonated 26 with a variety of reagents, such as *N*-fluorobenzenesulfonimide, perchloryl fluoride, or *N*-fluorobenzenedisulfonimide. Deprotonated 26 can be prepared by treatment of 26 with strong base, such as *n*-butyl lithium, sodium *bis*(trimethylsilyl)amide, sodium hydride, or lithium diisopropylamide. Deprotection of compounds 27 to give compounds I, where R<sup>1</sup> is H and R<sup>2</sup> is F is effected by standard procedures such as those described in Greene, T.W., *Protective Groups in Organic Synthesis*, 3rd edition 494-654 (1999) and United States Patent No. 5,874,448.

**Scheme 8: Synthesis of Compounds of Formula I, wherein one of one of X and Y is C=O and the other is CH<sub>2</sub>**




compounds I, where n is 0 or 1 and one of X and Y is C=O and the other is CH<sub>2</sub>

Scheme 8 depicts convenient general synthetic methodology to prepare compounds of Formula I, wherein one of X and Y is C=O and the other is CH<sub>2</sub> (*i.e.*, compounds IA and

**IB**). In compounds **IA**, the isoindolin-ring carbonyl is *cis* with respect to the methyleneamino ( $n = 1$ ) or the amino ( $n = 0$ ) group, conversely, in compounds **IB**, the isoindolin-ring carbonyl is *trans*. In one convenient methods, compounds **IA** and **IB** can be prepared starting from compounds **28** or **29** respectively, for example, using the methodology described in WO 98/54170, incorporated herein by reference. Compounds **28** and compounds **29** are available commercially or readily available through well-known synthetic methodology. For example, methyl-2-methyl-3-nitrobenzoate (**29**, where alkyl is methyl) is commercially available from Aldrich Chemical Co., Milwaukee, WI. Compounds **28** and **29** are first brominated at the activated benzylic position with a brominating agent such as *N*-bromosuccinimide under the influence of light or other radical initiator to yield methylbromo-compounds **30**. Exemplary brominating procedures are reviewed in March, J. *Advanced Organic Chemistry; Reactions Mechanisms, and Structure*, 4th ed., 1992, pp. 694-697, incorporated herein by reference. Compounds **30** are then converted into compounds **31** or **32** and thereafter to **IA** or **IB** by adapting the synthetic methods presented in Schemes 1 to 5 above, including standard cyclization with compounds **16**.

Scheme 9: Synthesis of Compounds of Formula I, wherein  $R^1$  is  $C(S)NHR^3$



Scheme 9 shows one method to synthesize compounds of Formula **I**, wherein  $R^1$  is  $C(S)NHR^3$ . A compound of Formula **I**, where  $R^1$  is H reacts with isothiocyanates **28** under routine conditions to give a compounds **I**, where  $R^1$  is  $C(O)NHR^5$ . The reaction is performed according to standard, well-known procedures, *e.g.* see March, J. *Advanced Organic Chemistry; Reactions Mechanisms, and Structure*, 4th ed., 1992, p. 903, incorporated herein by reference.

## 7. Therapeutic Uses of Compounds or Compositions of the Invention

In accordance with the invention, a compound or composition of the invention is administered to a mammal, preferably, a human, with or at risk of a disease or medical condition, for example, cancer, such as solid tumors and blood-born tumors. Specific  
5 examples of cancers treatable or preventable by administering compounds of the invention include, but are not limited to, cancers of the skin, such as melanoma; lymph node; breast; cervix; uterus; gastrointestinal tract; lung; ovary; prostate; colon; rectal; mouth; brain; head and neck; throat; testes; kidney; pancreas; bone; spleen; liver; bladder; larynx; nasal passages; and AIDS-related cancers. The compounds are particularly useful for treating  
10 cancers of the blood and bone marrow, such as multiple myeloma and acute and chronic leukemias, for example, lymphoblastic, myelogenous, lymphocytic, and myelocytic leukemias.

The compounds of the invention are also useful to treat or prevent heart disease, such as congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated  
15 septic shock, acute viral myocarditis, cardiac allograft rejection, and myocardial infarction.

The compounds of the invention can also be used to treat or prevent viral, genetic, inflammatory, allergic, and autoimmune diseases. For example, the compounds are useful to treat or prevent diseases including, but not limited to, HIV; hepatitis; adult respiratory distress syndrome; bone-resorption diseases; chronic pulmonary inflammatory diseases;  
20 dermatitis; cystic fibrosis; septic shock; sepsis; endotoxic shock; hemodynamic shock; sepsis syndrome; post ischemic reperfusion injury; meningitis; psoriasis; fibrotic disease; cachexia; graft rejection; auto-immune disease; rheumatoid spondylitis; arthritic conditions, such as rheumatoid arthritis and osteoarthritis; osteoporosis, Crohn's disease; ulcerative colitis; inflammatory-bowel disease; multiple sclerosis; systemic lupus erythematosus;  
25 ENL in leprosy; radiation damage; asthma; and hyperoxic alveolar injury.

The compounds of the invention are also useful for treating or preventing bacterial infections including, but not limited to, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

In one embodiment, "treatment" or "treating" refers to an amelioration of a disease  
30 or disorder, or at least one discernible symptom thereof. In another embodiment, "treatment" or "treating" refers to an amelioration of at least one measurable physical parameter, not necessarily discernible by the mammal. In yet another embodiment, "treatment" or "treating" refers to inhibiting the progression of a disease or disorder, either physically, e.g., stabilization of a discernible symptom, physiologically, e.g., stabilization of  
35

a physical parameter, or both. In yet another embodiment, "treatment" or "treating" refers to delaying the onset of a disease or disorder.

In certain embodiments, the compounds of the invention or the compositions of the invention are administered to a mammal, preferably, a human, as a prophylactic. As used  
5 herein, "prevention" or "preventing" refers to a reduction of the risk of acquiring a given disease or disorder. In a preferred mode of the embodiment, the compounds and compositions of the present invention are administered as a preventative measure to a mammal, preferably, a human, having a genetic or non-genetic predisposition to a medical condition, for example, cancers, such as solid tumors and blood-born tumors. Specific  
10 examples of cancers preventable by compounds of the invention include, but are not limited to, cancers of the skin, such as melanoma; lymph node; breast; cervix; uterus; gastrointestinal tract; lung; ovary; prostate; colon; rectal; mouth; brain; head and neck; throat; testes; kidney; pancreas; bone; spleen; liver; bladder; larynx; nasal passages; and AIDS-related cancers. The compounds are particularly useful for treating cancers of the  
15 blood and bone marrow, such as multiple myeloma and acute and chronic leukemias, for example, lymphoblastic, myelogenous, lymphocytic, and myelocytic leukemias.

The compounds of the invention are also useful for preventing heart disease, such as congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, and myocardial infarction.

20 The compounds of the invention can also be used to prevent viral, genetic, inflammatory, allergic, and autoimmune diseases. For example, the compounds are useful to prevent diseases including, but not limited to, HIV; hepatitis; adult respiratory distress syndrome; bone-resorption diseases; chronic pulmonary inflammatory diseases; dermatitis; cystic fibrosis; septic shock; sepsis; endotoxic shock; hemodynamic shock; sepsis  
25 syndrome; post ischemic reperfusion injury; meningitis; psoriasis; fibrotic disease; cachexia; graft rejection; auto-immune disease; rheumatoid spondylitis; arthritic conditions, such as rheumatoid arthritis and osteoarthritis; osteoporosis, Crohn's disease; ulcerative colitis; inflammatory-bowel disease; multiple sclerosis; systemic lupus erythematosus; ENL in leprosy; radiation damage; asthma; and hyperoxic alveolar injury.

30 The compounds of the invention are also useful for preventing bacterial infections or symptoms including, but not limited to, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

## 8. Therapeutic/Prophylactic Administration of The Compounds and Compositions of The Invention

Due to the activity of the compounds and compositions of the invention, they are useful in veterinary and human medicine. The invention provides methods of treatment and prevention by administration of a therapeutically effective amount of a compound or a composition of the invention to a mammal, preferably, a human. The term "mammal" as used herein, encompasses any mammal. Preferably a mammal is in need of such treatment or prevention. Examples of mammals include, but are not limited to, cows, horses, sheep, pigs, cats, dogs, mice, rats, rabbits, guinea pigs, monkeys, *etc.*, more preferably, a human.

Administration of compounds of the invention can be systemic or local. In most instances, administration to a mammal will result in systemic release of the compounds of the invention (*i.e.*, into the bloodstream). Methods of administration include enteral routes, such as oral, buccal, sublingual, and rectal; topical administration, such as transdermal and intradermal; and parenteral administration. Suitable parenteral routes include injection via a hypodermic needle or catheter, for example, intravenous, intramuscular, subcutaneous, intradermal, intraperitoneal, intraarterial, intraventricular, intrathecal, and intracameral injection and non-injection routes, such as intravaginal rectal, or nasal administration. Preferably, the compounds and compositions of the invention are administered orally. In specific embodiments, it may be desirable to administer one or more compounds of the invention locally to the area in need of treatment. This may be achieved, for example, by local infusion during surgery, topical application, *e.g.*, in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers.

The compounds of the invention can be administered via typical as well as non-standard delivery systems, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, capsules, *etc.* For example, the compounds and compositions of the invention can be delivered in a vesicle, in particular a liposome (*see* Langer, 1990, *Science* 249:1527-1533; Treat *et al.*, in *Liposomes in Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; *see generally ibid.*). In another example, the compounds and compositions of the invention can be delivered in a controlled release system. In one embodiment, a pump may be used (*see* Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:201; Buchwald *et al.*, 1980, *Surgery* 88:507 Saudek *et al.*, 1989, *N. Engl. J. Med.* 321:574). In another example, polymeric materials can be used (*see* Medical Applications of Controlled Release, Langer

and Wise (eds.), CRC Press., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61; *see also* Levy *et al.*, 1985, *Science* 228:190; During *et al.*, 1989, *Ann. Neurol.* 25:351; Howard *et al.*, 1989, *J. Neurosurg.* 71:105). In still another example, a controlled-release system can be placed in proximity of the target area to be treated, *e.g.*, the liver, thus requiring only a fraction of the systemic dose (*see, e.g.*, Goodson, in Medical Applications of Controlled Release, *supra*, vol. 2, pp. 115-138 (1984)). Other controlled-release systems discussed in the review by Langer, 1990, *Science* 249:1527-1533) can be used.

When administered as a composition, a compound of the invention will be formulated with a suitable amount of a pharmaceutically acceptable vehicle or carrier so as to provide the form for proper administration to the mammal. The term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in mammals, and more particularly in humans. The term "vehicle" refers to a diluent, adjuvant, excipient, or carrier with which a compound of the invention is formulated for administration to a mammal. Such pharmaceutical vehicles can be liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. The pharmaceutical vehicles can be saline, gum acacia, gelatin, starch paste, talc, keratin, colloidal silica, urea, and the like. In addition, auxiliary, stabilizing, thickening, lubricating and coloring agents may be used. Preferably, when administered to a mammal, the compounds and compositions of the invention and pharmaceutically acceptable vehicles, excipients, or diluents are sterile. An aqueous medium is a preferred vehicle when the compound of the invention is administered intravenously, such as water, saline solutions, and aqueous dextrose and glycerol solutions.

The present compounds and compositions can take the form of capsules, tablets, pills, pellets, lozenges, powders, granules, syrups, elixirs, solutions, suspensions, emulsions, suppositories, or sustained-release formulations thereof, or any other form suitable for administration to a mammal. In a preferred embodiment, the compounds and compositions of the invention are formulated for administration in accordance with routine procedures as a pharmaceutical composition adapted for oral or intravenous administration to humans. In one embodiment, the pharmaceutically acceptable vehicle is a hard gelatin capsule. Examples of suitable pharmaceutical vehicles and methods for formulation thereof are described in *Remington: The Science and Practice of Pharmacy*, Alfonso R. Gennaro ed.,

Mack Publishing Co. Easton, PA, 19th ed., 1995, Chapters 86, 87, 88, 91, and 92, incorporated herein by reference.

Compounds and compositions of the invention formulated for oral delivery, are preferably in the form of capsules, tablets, pills, or any compressed pharmaceutical form.

5 Moreover, where in tablet or pill form, the compounds and compositions may be coated to delay disintegration and absorption in the gastrointestinal tract thereby providing a sustained action over an extended period of time. Selectively permeable membranes surrounding an osmotically active driving compound are also suitable for orally administered compounds and compositions of the invention. In these later platforms, fluid from the environment  
10 surrounding the capsule is imbibed by the driving compound that swells to displace the agent or agent composition through an aperture. These delivery platforms can provide an essentially zero order delivery profile as opposed to the spiked profiles of immediate release formulations. A time delay material such as glycerol monostearate or glycerol stearate may also be used. Oral compositions can include standard vehicles, excipients, and diluents,  
15 such as magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, lactose, dextrose, sucrose, sorbitol, mannitol, starch, gum acacia, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidinone, water, syrup, and methyl cellulose, the formulations can additionally include lubricating agents, such as talc, magnesium stearate, mineral oil, wetting agents, emulsifying and suspending agents, preserving agents such as methyl- and  
20 propylhydroxybenzoates. Such vehicles are preferably of pharmaceutical grade. Orally administered compounds and compositions of the invention can optionally include one or more sweetening agents, such as fructose, aspartame or saccharin; one or more flavoring agents such as peppermint, oil of wintergreen, or cherry; or one or more coloring agents to provide a pharmaceutically palatable preparation.

25 A therapeutically effective dosage regimen for the treatment of a particular disorder or condition will depend on its nature and severity, and can be determined by standard clinical techniques according to the judgment of a medical practitioner. In addition, *in vitro* or *in vivo* assays can be used to help identify optimal dosages. Of course, the amount of a compound of the invention that constitutes a therapeutically effective dose also depends on  
30 the administration route. In general, suitable dosage ranges for oral administration are about 0.001 milligrams to about 20 milligrams of a compound of the invention per kilogram body weight per day, preferably, about 0.7 milligrams to about 6 milligrams, more preferably, about 1.5 milligrams to about 4.5 milligrams. In a preferred embodiment, a mammal, preferably, a human is orally administered about 0.01 mg to about 1000 mg of a compound  
35 of the invention per day, more preferably, about 0.1 mg to about 300 mg per day, or about 1

mg to about 250 mg in single or divided doses. The dosage amounts described herein refer to total amounts administered; that is, if more than one compound of the invention is administered, the preferred dosages correspond to the total amount of the compounds of the invention administered. Oral compositions preferably contain 10% to 95% of a compound of the invention by weight. Preferred unit oral-dosage forms include pills, tablets, and capsules, more preferably capsules. Typically such unit-dosage forms will contain about 0.01 mg, 0.1 mg, 1 mg, 5 mg, 10 mg, 15 mg, 20 mg, 50 mg, 100 mg, 250 mg, or 500 mg of a compound of the invention, preferably, from about 5 mg to about 200 mg of compound per unit dosage.

In another embodiment, the compounds and compositions of the invention can be administered parenterally (e.g., by intramuscular, intrathecal, intravenous, and intraarterial routes), preferably, intravenously. Typically, compounds and compositions of the invention for intravenous administration are solutions in sterile isotonic aqueous vehicles, such as water, saline, Ringer's solution, or dextrose solution. Where necessary, the compositions may also include a solubilizing agent. Compositions for intravenous administration may optionally include a local anesthetic such as lignocaine to ease pain at the site of the injection. For intravenous administration, the compounds and compositions of the invention can be supplied as a sterile, dry lyophilized powder or water-free concentrate in a hermetically sealed container, such as an ampule or sachette, the container indicating the quantity of active agent. Such a powder or concentrate is then diluted with an appropriate aqueous medium prior to intravenous administration. An ampule of sterile water, saline solution, or other appropriate aqueous medium can be provided with the powder or concentrate for dilution prior to administration. Or the compositions can be supplied in premixed form, ready for administration. Where a compound or composition of the invention is to be administered by intravenous infusion, it can be dispensed, for example, with an infusion bottle containing sterile pharmaceutical-grade water, saline, or other suitable medium.

Rectal administration can be effected through the use of suppositories formulated from conventional carriers such as cocoa butter, modified vegetable oils, and other fatty bases. Suppositories can be formulated by well-known methods using well-known formulations, for example see *Remington: The Science and Practice of Pharmacy*, Alfonso R. Gennaro ed., Mack Publishing Co. Easton, PA, 19th ed., 1995, pp. 1591-1597, incorporated herein by reference

To formulate and administer topical dosage forms, well-known transdermal and intradermal delivery mediums such as lotions, creams, and ointments and transdermal



delivery devices such as patches can be used (Ghosh, T.K.; Pfister, W.R.; Yum, S.I. *Transdermal and Topical Drug Delivery Systems*, Interpharm Press, Inc. p. 249-297, incorporated herein by reference). For example, a reservoir type patch design can comprise a backing film coated with an adhesive, and a reservoir compartment comprising a  
5 compound or composition of the invention, that is separated from the skin by a semipermeable membrane (*e.g.*, U.S. Patent 4,615,699, incorporated herein by reference). The adhesive coated backing layer extends around the reservoir's boundaries to provide a concentric seal with the skin and hold the reservoir adjacent to the skin.

The invention also provides pharmaceutical packs or kits comprising one or more  
10 containers filled with one or more compounds of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In one embodiment, the kit contains more than one compound of the invention. In another  
15 embodiment, the kit comprises a compound of the invention and another biologically active agent.

The compounds of the invention are preferably assayed *in vitro* and *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro* assays can be used to determine whether administration of a specific compound of the  
20 invention or a combination of compounds of the invention is preferred. The compounds and compositions of the invention may also be demonstrated to be effective and safe using animal model systems. Other methods will be known to the skilled artisan and are within the scope of the invention.

## 25       8.1    Combination Therapy

In certain embodiments, a compound of the invention is administered to a mammal, preferably, a human concurrently with one or more other biologically active agents, or with one or more other compounds of the invention, or with both. By "concurrently" it is meant that a compound of the invention and the other agent are administered to a mammal in a  
30 sequence and within a time interval such that the compound of the invention can act together with the other agent to provide an increased or synergistic benefit than if they were administered otherwise. For example, each component may be administered at the same time or sequentially in any order at different points in time; however, if not administered at the same time, they should be administered sufficiently closely in time so as to provide the  
35 desired treatment effect. Preferably, all components are administered at the same time, and

if not administered at the same time, preferably, they are all administered from about 6 hours to about 12 hours apart from one another.

When used in combination with other therapeutic agents, the compounds of the invention and the therapeutic agent can act additively or, more preferably, synergistically.

5 In one embodiment, a compound or a composition of the invention is administered concurrently with another therapeutic agent in the same pharmaceutical composition. In another embodiment, a compound or a composition of the invention is administered concurrently with another therapeutic agent in separate pharmaceutical compositions. In still another embodiment, a compound or a composition of the invention is administered  
10 prior or subsequent to administration of another therapeutic agent. As many of the disorders for which the compounds and compositions of the invention are useful in treating are chronic disorders, in one embodiment combination therapy involves alternating between administering a compound or a composition of the invention and a pharmaceutical composition comprising another therapeutic agent, *e.g.*, to minimize the toxicity associated  
15 with a particular drug. In certain embodiments, when a composition of the invention is administered concurrently with another therapeutic agent that potentially produces adverse side effects including, but not limited to toxicity, the therapeutic agent can advantageously be administered at a dose that falls below the threshold that the adverse side effect is elicited.

20 The present compounds and compositions can be administered together with hormonal and steroidal anti-inflammatory agents, such as, estradiol, conjugated estrogens (*e.g.*, PREMARIN, PREMPRO, AND PREMPHASE), 17 beta estradiol, calcitonin-salmon, levothyroxine, dexamethasone, medroxyprogesterone, prednisone, cortisone, flunisolide, and hydrocortisone; non-steroidal anti-inflammatory agents, such as tramadol, fentanyl,  
25 metamizole, ketoprofen, naproxen, nabumetone, ketoralac, tromethamine, loxoprofen, ibuprofen, aspirin, and acetaminophen; anti-TNF- $\alpha$  antibodies, such as infliximab (REMICADE™) and etanercept (ENBREL™); AIDS and AIDS-related therapies, such as lamivudine, zidovudine, indinavir sulfate, stavudine, and lamivudine; chemotherapeutics and cancer-related therapies, such as anti-angiogenesis agents, topoisomerase inhibitors,  
30 alkylating agents, nitrogen mustards, antibiotics such as doxorubicin, and paclitaxel, cisplatin, tamoxifen, docetaxel, irinotecan, temozolomide, thalidomide, amino-thalidomide, amino-EM-12, epirubicin, leuprolide, bicalutamide, goserelin implant, gemcitabine, sargramostim anti-cancer antibodies, such as Rituxan, and anti-cancer vaccines, such as Theratope and HSPPC-96; antibiotics, such as amoxicillin, ampicillin sodium, cefaclor, and  
35 ciprofloxacin; dermatological therapeutics, such as isotretinoin, clindamycin phosphate

topical; antiarthritic therapies, such as diclofenac sodium, nabumetone, misoprostol, and rofecoxib; immunosuppressive therapies, such as cyclosporine, FK506, mycophenolate mofetil, and methylprednisolone; multiple sclerosis therapies, such as interferon beta-1a, interferon beta-1b, and glatiramer; osteoporosis therapies, such as vitamin K<sub>2</sub>; cystic fibrosis  
5 therapies, such as dornase alpha and tobramycin; and Alzheimer's disease therapies, such as dolasetron mesylate, and donepezil hydrochloride.

In one embodiment of the invention, the compounds of the invention can be used, not only to directly treat the disorder, but also to reduce the dose or toxicity of another chemotherapeutic. For example, the compounds of the invention can be administered to  
10 reduce gastrointestinal toxicity associated with a topoisomerase inhibitor, such as irinotecan.

## 8.2 Assays

The compounds of the invention can be assayed for their ability to modulate the  
15 production of TNF- $\alpha$  by well-known methods in the art, *see e.g.*, Corral *et al.*, 1999, *J. Immun.* 163:380-386 and Muller *et al.*, 1996, *J. Med. Chem.* 39:3238 (assay for the inhibition of production of TNF- $\alpha$ ) and Muller *et al.*, 1998, *Bioorg. Med. Chem. Lett.* 8:2669, all three of which citations are incorporated herein by reference.

### 20 8.2.1 Assay For The Ability of a Compound of The Invention to Modulate the Production of TNF- $\alpha$

PBMC cells —normal human donors— were obtained by Ficoll-Hypaque density centrifugation (Pharmacia Fine Chemicals, Piscataway, NJ). The cells (about  $2 \times 10^5$  to  $10^6$  cells/ml) are cultured with RPMI (commercially available, *e.g.*, from Gibco Laboratories,  
25 Grand Island, NY) supplemented with 10AB+ serum (commercially available, *e.g.*, from Biocell, Rancho Dominguez, CA), about 2 mM L-glutamine, about 100 U/ml penicillin, and about 100  $\mu$ g/ml streptomycin (Gibco). The test compounds are dissolved in DMSO at 20 mg/ml, further dilution can be done with culture medium. The final DMSO concentration in all samples including the controls should be about 0.25% by weight. Test compounds  
30 were added to cells 1 hour prior to the addition of LPS. The PBMC cells, in triplicate, are stimulated by 1  $\mu$ g/ml LPS from *Salmonella* Minnesota R595 (List Biological Labs, Campbell, CA) and incubated for about 18 to about 20 hours at 37 °C (5% CO<sub>2</sub>) in 96-well flat-bottom polystyrene Costar tissue culture plates (Corning, Corning, NY) for the induction of TNF- $\alpha$ . Cells are incubated with or without compounds of the invention  
35 (negative controls). The supernatants are collected for the determination of cytokine levels by ELISA (Endogen, Cambridge, MA). Percent inhibition can be determined as  $100 \times [1 -$

(TNF- $\alpha$  EXPERIMENTAL/ TNF- $\alpha$  CONTROL)]. Assays are performed, in accordance with the assay kit's manufacturer, in 96-well plates (Nunc Immunoplates, Roskilde, Denmark) coated with the affinity-purified rabbit anti-TNF- $\alpha$  antibody (0.5  $\mu$ g/ml; 12-16 hours; 4 °C) and blocked for 2 hours at room temperature with PBS/0.05% Tween 20 (Sigma Chemical Co., St. Louis, Mo.) containing 5 mg/ml BSA. After washing, 100  $\mu$ l of TNF- $\alpha$  standards, samples and controls are applied to the wells, and the plates are incubated for 12-24 hours at 4 °C. After the incubation, plates are washed and a second antibody, horseradish peroxidase (HRP)-conjugated mouse monoclonal anti-TNF- $\alpha$ , diluted 1:2,000 in PBS/BSA/Tween, is applied to the wells, after which they incubated for 2 hours at room temperature. The color reaction is developed with the OPD substrate (0.4 mg/ml *o*-phenylenediamine [Sigma Chemical Co.] in 24 mM citric acid, 51 mM sodium phosphate, pH 5.0 [phosphate-citrate buffer: Sigma Chemical Co.] containing 0.012% hydrogen peroxide [Fisher Scientific Co., Pittsburgh, PA.]) and absorbance read at 492 nm in an automated ELISA reader (Dynatech Laboratories, Inc., Alexandria, Va.).

15

### 8.2.2 Assay For T-Cell Stimulation and IL-2 Stimulation:

#### PMC Stimulation by Anti-CD3 Ab

PBMC ( $1 \times 10^6$  cells) are stimulated by cross-linking of the TCR by immobilized monoclonal mouse anti-human CD3 (Orthoclone OKT3) as described in Haslett *et al.*, 1998, *J. Exp. Med.* 187:1885, incorporated herein by reference. The anti-CD3 Ab is diluted to 10  $\mu$ g/ml in 100  $\mu$ l PBS and coated onto 48-well flat-bottom polystyrene Falcon tissue culture plates (Becton Dickinson, Franklin Lakes, NJ) by overnight incubation at 4 °C. Appropriate dilutions of compounds of the invention are added at the start of the cell culture. Supernatants are collected at 24, 43, and 72 hours and assayed for TNF- $\alpha$  levels. Cells are collected at 48 hours for evaluation of CD40 ligand (CD40L)<sup>3</sup> and CD3 surface expression by two-color flow cytometry (anti-CD40L, PharMingen, San Diego, CA; anti-CD3, Becton Dickinson, San Jose, CA).

### 8.2.3 Assay For the Modulation of Production of IL-1 $\beta$ and IL-10

This assay can be performed according to the procedure outlined in Muller *et al.*, 1999, *J. Immunol.* 176, 380, hereby expressly incorporated herein by reference. PMBC ( $2 \times 10^5$  cells) incubated in 96-well flat-bottom polystyrene *Costar* tissue culture plates (Corning, Corning, NY) were stimulated by 1 mg/ml LPS from *Salmonella minnesota* R595 (List Biological Labs, Campbell, CA) for the induction of IL-1 $\beta$ , and IL-10. Cells were incubated with or without thalidomide or analogues for 20 h, and culture supernatants

35

were collected and frozen immediately at - 70 °C until assayed in triplicate or duplicate. IL-1 $\beta$  and IL-10 levels were measured by ELISA (Endogen, Cambridge, MA) as described by the manufacturer.

5


## 9. Examples of Syntheses of Compounds of the Invention

The following Examples further illustrate methods for synthesizing compounds and intermediates of the invention. It is to be understood that the invention is not limited to the specific details of the Examples set forth below.

10

### Methyl-3-amino-2-(methoxycarbonyl)benzoate

15




To a solution of methyl-2-(methoxycarbonyl)-3-nitrobenzoate (23.8 g, 99.51 mmol) in ethyl acetate (200 ml) was added 10% Pd/C (1.8 g). The mixture was hydrogenated under 50 psi of hydrogen for 3 hours in a Parr Type Shaker. The mixture was filtered through Celite and the filtrate was concentrated *in vacuo* to yield an oil. The crude product was purified by flash chromatography (dichloromethane/ethyl acetate 95 to 5) to afford 18.1 g (87%) of the product as a brown oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.22 (t, J=7.6 Hz, 1H), 6.90 (d, J=7.2 Hz, 1H), 6.79 (d, J=8.7 Hz, 1H), 5.07 (b, 2H), 3.85 (s, 3H), 3.83 (s, 3H).

25

### Methyl-3-cyano-2-(methoxycarbonyl)benzoate


30



To a stirred suspension of methyl-3-amino-2-(methoxycarbonyl)benzoate (17.0 g, 81 mmol) in a mixture of concentrated HCl (44 ml) and water (440 ml) at 4 °C was added a

solution of  $\text{NaNO}_2$  (6.73 g, 97 mmol) in water (25 ml) dropwise at 4-5 °C. Stirring was continued for 30 min at 4 °C. The mixture was then carefully neutralized with sat. sodium carbonate to pH 6. A stirred solution of  $\text{CuCN}$  (9.46 g, 105 mmol) and  $\text{KCN}$  (6.38 g, 105 mmol) in water (150 ml) was warmed to 60 °C. The cold neutralized diazonium solution was then added in small portions at a time with vigorous stirring. The mixture was stirred at 60 °C for 1 hour and then cooled to room temperature. The mixture was extracted with dichloromethane (4 × 150 ml) and the combined dichloromethane extracts were washed with water (2 × 100 ml), brine (100 ml) and dried. The solvent was removed *in vacuo* and the product was purified by chromatography (dichloromethane) to afford 12.36 g (65%) of the product as a light yellow solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  8.19 (d,  $J=7.9$  Hz, 1H), 7.89 (d,  $J=7.4$  Hz, 1H), 7.64 (t,  $J=8.0$  Hz, 1H), 4.03 (s, 3H), 3.94 (s, 3H).

Methyl-3-aminomethyl-2-(methoxycarbonyl)benzoate hydrochloride



20 To a solution of methyl-3-cyano-2-(methoxycarbonyl)benzoate (12.3 g, 57 mmol) in methanol (250 ml) and 4N HCl (40 ml) was added 10% Pd/C (1.2 g). The mixture was hydrogenated under 50 psi of hydrogen in a Parr Type Shaker for 17 hours. The mixture was filtered through Celite and the filtrate was concentrated *in vacuo*. The residue was further evaporated with ethanol (2 × 25 ml) and toluene (25 ml) and dried under vacuum.


25 The resulting solid was slurried in ether (50 ml) for 1 hour. The slurry was then filtered and dried to give 13.46 g (90%) of the product as a white solid:  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  8.79 (b, 2H), 7.94 (d,  $J=7.8$  Hz, 1H), 7.88 (d,  $J=7.7$  Hz, 1H), 7.72 (t,  $J=7.7$  Hz, 1H), 4.03 (s, 2H), 3.86 (s, 3H), 3.85 (s, 3H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  167.58, 166.12, 133.41, 133.18, 132.28, 130.62, 129.49, 52.99, 52.92, 39.25.

30

35

Methyl-3-[(*t*-butoxycarbonylamino)methyl]-2-(methoxycarbonyl)benzoate

5




10 Triethylamine (3.89 g, 38 mmol) was added dropwise to a stirred suspension of methyl-3-aminomethyl-2-(methoxycarbonyl)benzoate hydrochloride (4.0 g, 15 mmol) in dichloromethane (100 ml). The mixture was cooled in an ice bath to 8 °C. A solution of di-  
 15 *t*-butyl dicarbonate (3.7 g, 16 mmol) in dichloromethane (20 ml) was added dropwise at 8 °C. After the addition was complete, the cooled mixture was stirred for an additional 30 minutes, and then warmed to room temperature for 1 hour. The mixture was washed with water (2 × 40 ml), brine (40 ml) and dried. Solvent was removed *in vacuo* and the product was purified by chromatography (hexane/ethyl acetate 7 to 3) to afford 4.66 g (93%) of the product as an oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.87 (d, J=7.4 Hz, 1H), 7.62 (d, J=7.4 Hz, 1H), 7.46 (t, J=7.8 Hz, 1H), 5.16 (b, 1H), 4.30 (d, J=6.1 Hz, 2H), 3.93 (s, 3H), 3.89 (s, 3H), 1.42 (s, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 169.41, 166.18, 155.57, 137.09, 134.04, 133.32, 129.76, 128.95, 128.64, 79.52, 52.72, 52.50, 42.23.

20

[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-ylmethyl]-carbamic acid *tert*-butyl ester I-1

25




30

Diisopropylethylamine (3.20 g, 25 mmol) was added to a stirred suspension of methyl-3-[(*t*-butoxycarbonylamino)methyl]-2-(methoxycarbonyl)benzoate (8.00 g, 25 mmol) and aminoglutarimide hydrochloride (4.07 g, 25 mmol) in DMF (60 ml). The  
 35 mixture was heated to 120 °C for 24 hours and then cooled to room temperature. The

mixture was poured into cold water (300 ml) and extracted with ethyl acetate (4x100 ml each). The combined ethyl acetate extracts were washed with water (2 x 50 ml), brine (50 ml) and dried. Solvent was removed *in vacuo* and the product purified by flash chromatography (dichloromethane/ethyl acetate 8 to 2) to yield 4.66 g of recovered starting material and 3.31 g (82%) of the product as a white solid: mp 180-182 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.51 (s, 1H), 7.81-7.67 (m, 3H), 5.54 (b, 1H), 5.03-4.96 (dd, J=5.2 and 11.2 Hz, 1H), 4.66 (d, J=6.3 Hz, 2H), 2.95-2.74 (m, 3H), 2.18-2.14 (m, 1H), 1.43 (s, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 170.99, 168.08, 167.95, 167.07, 155.86, 139.17, 135.00, 134.61, 132.15, 128.22, 122.82, 79.81, 49.21, 40.53, 31.33, 28.32, 22.58; Anal. Calcd for C<sub>19</sub>H<sub>21</sub>N<sub>3</sub>O<sub>6</sub>: C, 58.91; H, 5.46; N, 10.85. Found: C, 59.08; H, 5.51; N, 10.69.

4-(Aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione I-2, hydrochloride




A solution of 4 N HCl in dioxane (10 ml) was added to a stirred solution of (*t*-butoxy)-*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl}carboxamide (3.3 g, 8.5 mmol) in dichloromethane (50 ml). The mixture was stirred at room temperature overnight. The resulting slurry was filtered and dried to afford 2.4 g (87%) of the product as a white solid: mp 291-293 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.18 (s, 1H), 8.77 (b, 2H), 8.06-7.93 (m, 3H), 5.22-5.15 (dd, J=5.1 and 12.6 Hz, 1H), 4.49 (s, 2H), 2.97-2.85 (m, 1H), 2.65-2.51 (m, 2H), 2.08-2.04 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.86, 169.76, 167.35, 166.71, 135.62, 134.98, 132.80, 131.46, 128.62, 123.57, 49.00, 37.00, 30.95, 22.07; Anal. Calcd for C<sub>14</sub>H<sub>14</sub>N<sub>3</sub>O<sub>4</sub>Cl + 0.22water: C, 51.31; H, 4.44; N, 12.82; Cl, 10.82. Found: C, 51.08; H, 4.36; N, 12.47; Cl, 10.61.



3-(Acetylamino-methyl)-phthalic acid dimethyl ester

5




Triethylamine (1.87 g, 18 mmol) was added slowly to a stirred suspension of  
 10 methyl-3-(aminomethyl)-2-(methoxycarbonyl)benzoate hydrochloride (2.0 g, 8 mmol) in  
 dichloromethane (30 ml). The resulting mixture was cooled in an ice bath to 4 °C. Acetyl  
 chloride (0.73 g, 9 mmol) was added dropwise at a rate such that the temperature stayed  
 between 4-7 °C. After addition was complete, the mixture was stirred in the ice bath for an  
 15 hours. The reaction mixture was washed with water (2 × 30 ml), brine (30 ml) and dried.  
 Solvent was removed *in vacuo* and the product was purified by chromatography  
 (dichloromethane/ethyl acetate 6 to 4) to afford 1.65 g (80%) of the product as an oil: <sup>1</sup>H  
 NMR (CDCl<sub>3</sub>) δ 7.86 (d, J=7.7 Hz, 1H), 7.62 (d, J=7.6 Hz); 7.46 (t, J=7.7 Hz); 6.29 (b,  
 1H), 4.39 (d, J=6.1 Hz, 2H), 3.93 (s, 3H), 3.90 (s, 3H), 1.96 (s, 3H).

20

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-acetamide  
I-3

25



30


1,8-Diazabicyclo[5,4,0]undec-7-ene (0.93 g, 6 mmol) was added dropwise to a  
 stirred suspension of 3-(acetylamino-methyl)-phthalic acid dimethyl ester (1.61 g, 6.0  
 mmol) and aminoglutarimide hydrochloride (1.0 g, 6.0 mmol) in DMF (15 ml). The  
 35 mixture was then heated to 120 °C for 24 hours. The cooled mixture was concentrated *in*

*vacuo* and the residue was stirred with water (25 ml) and dichloromethane (20 ml). The resulting slurry was filtered to give 0.45 g (22%) of the product as a gray solid.

Recrystallization from methanol gave a white solid: mp 177-179 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ( 11.02 (s, 1H), 8.36 (t, J=5.8 Hz, 1H), 7.74-7.55 (m, 3H), 5.05-4.98 (dd, J=5.3 and 12.5 Hz, 1H), 4.57 9D, j=5.9 Hz, 2H), 2.84-2.70 (m, 1H), 2.51-2.34 (m, 2H), 1.95-1.91 (m, 1H), 1.79 (s, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ ( 172.85, 169.89, 169.81, 167.56, 167.05, 139.37, 134.83, 133.35, 131.58, 127.14, 121.94, 48.91, 37.84, 30.98, 22.54, 22.05; Anal. Calcd. For C<sub>16</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub> + 0.96 water: C, 55.45; H, 4.92; N, 12.12. Found: C, 55.27; H, 4.82; N, 12.00.

10 Methyl-3-[(cyclopropylcarbonylamino)methyl]-2-(methoxycarbonyl)benzoate

15



Triethylamine (1.87 g, 18 mmol) was added dropwise to a stirred suspension of methyl-3-(aminomethyl)-2-(methoxycarbonyl)benzoate hydrochloride (2.0 g, 7 mmol) in dichloromethane (40 ml). The mixture was cooled in an ice bath to 4 °C.

Cyclopropylcarbonyl chloride (0.99 g, 9 mmol) was added slowly at 4-8 °C. After addition, the mixture was stirred in ice bath for 30 min and then warmed to room temperature for 2 hours. The mixture was washed with water (2 × 30 ml), brine (30 ml) and dried. Solvent was removed *in vacuo* and the product was purified by flash chromatography

(dichloromethane/ethyl acetate 9 to 1) to give 2.1 g (93%) of the product as a white solid : <sup>1</sup>H NMR ( CDCl<sub>3</sub>) ( 7.87 (d, J=7.8 Hz, 1H), 7.63 (d, J=7.7 Hz, 1H), 7.46 (t, J=7.77 Hz, 1H), 6.31 (m, 1H), 4.43 (d, J=6.0 Hz, 2H), 3.95 (s, 3H), 3.90 (s, 3H), 1.36-1.29 (m, 1H), 0.99-0.93 (m, 2H), 0.76-0.69 (m, 2H).


30

35

N-{[2-(2,6-Dioxo(3-piperidyl)-1,3-dioxoisindolin-4-yl)methyl]cyclopropyl-carboxamide  
I-4

5

10



15

20


25

Diisopropylethylamine (0.92 g, 7 mmol) was added to a stirred suspension of methyl-3-[(cyclopropylcarbonylamino)methyl]-2-(methoxycarbonyl)benzoate (2.08 g, 7 mmol) and aminoglutarimide hydrochloride (1.17 g, 7 mmol) in DMF (15 ml). The mixture was heated to 120 °C for 24 hours. The mixture was concentrated *in vacuo* and the residue was stirred with water (40 ml) and ethyl acetate (15 ml). The resulting slurry was filtered to give 0.7 g (27%) of the product as a gray solid. Recrystallization for methanol gave a white solid: mp 240-242 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.06 (s, 1H), 8.62 (t, J=5.8 Hz, 1H), 7.79-7.70 (m, 2H), 7.60 (d, J=7.2 Hz, 1H), 5.09-5.02 (dd, J=5.3 and 12.5 Hz, 1H), 4.65 (d, J=5.8 Hz, 2H), 2.83-2.73 (m, 1H), 2.54-2.41 (m, 2H), 1.99-1.94 (m, 1H), 1.57 (m, 1H), 0.62-0.60 (m, 4H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 173.21, 172.85, 169.90, 167.52, 167.01, 139.44, 134.85, 133.37, 131.57, 127.13, 121.94, 48.89, 37.82, 30.97, 22.03, 13.58, 6.52; Anal. Calcd. For C<sub>18</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>: C, 60.84; H, 4.82; N, 11.82. Found: C, 60.46; H, 4.84; N, 11.65.

30


35

Methyl-3-[(Ethoxycarbonylamino)methyl]-2-(methoxycarbonyl)benzoate



Triethylamine (1.57 g, 18.5 mmol) was added to a stirred suspension of methyl-3-(aminomethyl)-2-(methoxycarbonyl)benzoate hydrochloride (2.0 g, 8 mmol) in dichloromethane (30 ml). The mixture was cooled in an ice bath to 4 °C. Ethyl chloroformate (1.0 g, 9 mmol) was added slowly keeping the mixture at 4-6 °C. After addition was complete, the mixture was stirred in an ice bath for 30 minutes and then warmed to room temperature for 2 hours. The mixture was washed with water (2 × 30 ml), brine (30 ml) and dried. Solvent was removed *in vacuo* and the residue was purified by flash chromatography (dichloromethane/ethyl acetate 95 to 5) to give 1.59 g (70%) of the product as an oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.87 (d, J=7.8 Hz, 1H), 7.36 (d, J=7.6 Hz, 1H), 7.46 (t, J=7.7 Hz, 1H), 5.30 (b, 1H), 4.32 (d, J=6.3 Hz, 2H), 4.12 (q, J=7.0 Hz, 2H), 3.93 (s, 3H), 3.89 (s, 3H), 1.22 (t, J=7.2 Hz, 3H).

[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-carbamic acid ethyl ester I-5




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.8 g, 5 mmol) was added to a stirred suspension of methyl-3-[(ethoxycarbonylamino)methyl]-2-(methoxycarbonyl)benzoate (1.54 g, 5 mmol) and aminoglutarimide hydrochloride (0.86 g, 5 mmol) in DMF (15 ml). The mixture was heated to 120 °C for 24 hours. The mixture was cooled to room temperature and poured into water (150 ml). The mixture was extracted with ethyl acetate (3 × 30 ml) and the ethyl acetate solution was washed with water (30 ml), brine (30 ml) and dried. Solvent was removed and the residue was purified by flash chromatography (dichloromethane/ethyl acetate 7 to 3) to give 0.84 g (45%) of the product as a white solid : mp 187-189 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.03 (s, 1H), 7.76-7.58 (m, 4H), 5.06-4.99 (dd, J=5.4 and 12.6 Hz, 1H), 4.52 (d, J=5.9 Hz, 2H), 3.91 (q, J=7.0 Hz, 2H), 2.79-2.70 (m, 1H), 2.51-2.38 (m, 2H), 1.96-1.86 (m, 1H), 1.06 (t, J=7.0 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.89, 169.93, 167.55, 167.05, 156.72, 139.69, 134.91, 132.90, 131.58, 127.04, 121.97,

60.15, 48.90, 39.34, 30.98, 22.04, 14.69; Anal. Calcd. For  $C_{17}H_{17}N_3O_6$ : C, 56.82; H, 4.77; N, 11.69. Found: C, 56.94; H, 4.81; N, 11.37.

Methyl-3-[(benzyloxycarbonylamino)methyl]-2-(methoxycarbonyl)benzoate

5

10



15


20

Triethylamine (1.87 g, 18.5 mmol) was added to a stirred suspension of methyl-3-(aminomethyl)-2-(methoxycarbonyl)benzoate hydrochloride (2.0 g, 8 mmol) in dichloromethane (30 ml). The mixture was cooled in an ice bath to 4 °C. Benzyl chloroformate (1.66 g, 10 mmol) was added slowly keeping the temperature between 4-7 °C. After the addition was complete, the cooled mixture was stirred an additional 30 minutes and then warmed to room temperature for 4 hours. The mixture was washed with water (2 × 30 ml), brine (30 ml) and dried. Solvent was removed and the residue was purified by chromatography (dichloromethane/ethyl acetate 95 to 5) to give 2.1 g (76%) of the product as a solid;  $^1H$  NMR ( $CDCl_3$ )  $\delta$  7.87 (d,  $J=7.8$  Hz, 1H), 7.63 (d,  $J=7.6$  Hz, 1H), 7.46 (t,  $J=7.7$  Hz, 1H), 7.36-7.26 (m, 5H), 5.41 (b, 1H), 5.09 (s, 2H), 4.36 (d,  $J=6.3$  Hz, 2H), 3.91 (s, 3H), 3.88 (s, 3H).

25

[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-carbamic acid benzyl ester I-6


30



35

1,8-diazabicyclo[5,4,0]undec-7-ene (0.88 g, 6 mmol) was added to a stirred suspension of methyl-3-[(benzyloxycarbonylamino)methyl]-2-(methoxycarbonyl)benzoate (2.07 g, 6 mmol) and aminoglutarimide hydrochloride (0.95 g, 6 mmol) in DMF (15 ml). The mixture was heated to 120 °C for 24 hours. The mixture was cooled to room temperature and poured into water (150 ml). The mixture was extracted with ethyl acetate (3 × 30 ml) and the combined ethyl acetate extracts were washed with water (2 × 30 ml), brine (30 ml) and dried. Solvent was removed *in vacuo* and the residue was purified by chromatography (dichloromethane/ethyl acetate 8 to 2) to give 0.58 g (24%) of the product as a white solid: mp 166-168 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.15 (s, 1H), 7.99-7.71 (m, 4H), 7.37 (m, 5H), 5.19-5.12 (dd, J=5.1 and 17.4 Hz, 1H), 5.07 (s, 2H), 4.70 (d, J=5.7 Hz, 2H), 2.97-2.83 (m, 1H), 2.64-2.51 (m, 2H), 2.08-2.04 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.85, 169.89, 167.51, 167.01, 156.55, 139.46, 137.02, 134.87, 132.89, 131.57, 128.44, 127.92, 127.83, 127.06, 121.99, 65.69, 48.89, 30.97, 22.02; Anal. Calcd. For C<sub>22</sub>H<sub>19</sub>N<sub>3</sub>O<sub>6</sub>: C, 62.70; H, 4.54; N, 9.97. Found: C, 62.53; H, 4.57; N, 9.89.


2-Chloro-N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}acetamide I-7



Triethylamine (0.6 g, 6 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.8 g, 2.5 mmol) in THF (70 ml). After stirring for 5 min, chloroacetyl chloride (0.34 g, 3 mmol) was added and the resulting mixture was heated at reflux for 3 hours. The solvent was removed *in vacuo* and the residue was dissolved in dichloromethane (70 ml), washed with water (20 ml), 2N HCl (30 ml), water (2 × 30 ml), brine (30 ml) and dried. Solvent was removed *in vacuo* and the resulting solid was slurried in dichloromethane (10 ml) and ether (10 ml) and filtered to give 0.76 g (84%) of the product: <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.15 (s, 1H), 8.87 (t, J=5.9 Hz, 1H), 7.88-7.68 (m, 3H), 5.19-5.12 (dd, J=5.3 and 12.6 Hz, 1H), 4.77 (d, J=5.9 Hz, 2H), 4.19 (s, 2H), 2.96-2.83 (m, 1H), 2.65-2.51 (m, 2H), 2.11-2.04 (m, 1H); <sup>13</sup>C NMR

(DMSO-*d*<sub>6</sub>)  $\delta$  172.88, 169.90, 167.53, 167.01, 166.67, 138.46, 134.90, 133.22, 131.63, 127.26, 122.14, 48.94, 42.61, 38.27, 30.99, 22.05.

2-(Dimethylamino)-*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-  
 5 acetamide I-8, hydrochloride




Dimethylamine (2M in THF, 5 ml, 10 mmol) was added to a stirred suspension of 2-chloro-*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}acetamide (1.2 g, 3.3 mmol) in acetonitrile (120 ml). The mixture was stirred at room temperature overnight.

20 Solvent was removed *in vacuo* and the residue was dissolved in dichloromethane (75 ml), washed with water (30 ml), brine (30 ml) and dried. Solvent was removed *in vacuo* and the residue was purified by flash chromatography (dichloromethane/methanol 95 to 5) to give 0.96 g (78%) of the free base. The free base was dissolved in ethyl acetate (20 ml) and treated with 1N HCl (5 ml) to afford 0.9 g (86%) of the hydrochloride salt: mp 185-187 °C;

25 <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  11.15 (s, 1H), 10.05 (b, 1H), 9.40 (s, 1H), 7.84 (m, 3H), 5.14 (m, 1H), 4.81 (s, 2H), 4.07 (s, 2H), 2.84 (s, 6H), 2.65-2.52 (m, 3H), 2.09 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>)  $\delta$  172.81, 169.82, 167.40, 166.89, 164.77, 137.91, 134.88, 133.53, 131.55, 127.25, 122.23, 57.21, 48.88, 43.20, 37.91, 30.93, 21.89; Anal. Calcd. For C<sub>18</sub>H<sub>21</sub>N<sub>4</sub>O<sub>5</sub>Cl + 0.65 water : C, 51.41; H, 5.34; N, 13.32; Cl, 8.43. Found: C, 51.12; H, 5.20; N, 12.67; Cl,

30 8.45.


1-tert-Butyl-3-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-urea I-9



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.28 g, 1.9 mmol) was added to stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.9 mmol) in acetonitrile (50 ml). After stirring for 1 hour, *t*-butylisocyanate (0.21 g, 2 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in dichloromethane (70 ml) and washed with 0.1 N HCl (20 ml), water (20 ml), brine (20 ml) and dried. The solvent was removed *in vacuo* and the resulting solid was recrystallized from ethanol/isopropyl ether to give 0.36 g (51%) of the product: mp 186-188 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.04 (s, 1H), 7.77-7.59 (m, 3H), 6.17 (t, J=6.2 Hz, 1H), 5.86 (s, 1H), 5.08-5.01 (dd, J=5.4 and 12.4 Hz, 1H), 4.49 (d, J=6.0 Hz, 2H), 2.82-2.73 (m, 1H), 2.54-2.40 (m, 2H), 1.98-1.94 (m, 1H), 1.12 (s, 9H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.33, 169.39, 167.11, 166.59, 156.85, 140.72, 134.20, 132.99, 131.06, 126.54, 121.20, 48.69, 48.37, 37.89, 30.46, 28.88, 21.53; Anal. Calcd. For C<sub>19</sub>H<sub>22</sub>N<sub>4</sub>O<sub>5</sub> + 0.2 water : C, 58.51; H, 5.79; N, 14.37. Found: C, 58.86; H, 6.15; N, 14.24.




N-([2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl)-3,3-dimethylbutanamide  
I-10



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62 g, 4 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.9 mmol) in acetonitrile (50 ml). After stirring for 20 min, *t*-butylacetyl chloride (0.25 g, 1.9 mmol) was added. The mixture was stirred at room temperature for 17 hours. Solvent was removed *in vacuo* and the residue was dissolved in dichloromethane (90 ml) and washed with 0.1N HCl (30 ml), water (30 ml), brine (30 ml) and then dried. Solvent was removed *in vacuo* and the solid residue was slurried in ethanol (10 ml) to give after filtration 0.55 g (77%) of the product: mp 145-147 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.14 (s, 1H), 8.39 (t, J=5.7 Hz, 1H), 7.87-7.69 (m, 3H), 5.19-5.12 (dd, J=5.3 and 12.4 Hz, 1H), 4.72 (d, J=5.8 Hz, 2H), 2.92-2.83 (m, 1H), 2.63-2.51 (m, 2H), 2.08 (s, 2H), 2.08-2.04 (m, 1H), 0.97 (s, 9H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.69, 171.35, 169.74, 167.51, 166.96, 139.61, 134.69, 133.41, 131.52, 127.11, 121.87, 48.86, 48.62, 37.53, 30.93, 30.52, 29.71, 22.00; Anal. Calcd. For C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O<sub>5</sub> + 0.28 water : C, 61.52; H, 6.08; N, 10.76. Found: C, 61.23; H, 6.18; N, 10.57.

N-[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-3-pyridylcarboxamide I-11




A stirred mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.09 g, 4.0 mmol) and nicotinoyl chloride hydrochloride (1.42 g, 8.0 mmol) in tetrahydrofuran (60 ml) was heated to reflux for 22 h. The suspension was filtered and washed with  
5 tetrahydrofuran (20 ml) and ether (10 ml) to yield a white solid. The solid was slurried in pH 7 buffer (40 ml) and ether (30 ml) for 1h. The suspension was filtered and washed with water (20 ml) and ether (20 ml) to N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]-3-pyridylcarboxamide as a white solid (1.2 g, 79% yield): mp, 176-178 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 2.06-2.10 (m, 1H, CHH), 2.49-2.65 (m, 2H, CH<sub>2</sub>), 2.83-2.97 (m, 1H, CHH),  
10 5.18 (dd, *J* = 5.3, 12.5 Hz, 1H, NCH), 7.64 (dd, *J* = 4.9, 7.9 Hz, 1H, Ar), 7.71 (d, *J* = 7.3 Hz, 1H, Ar), 7.92 (t, *J* = 7.8 Hz, 1H, Ar), 8.29-8.34 (m, 1H, Ar), 8.45 (d, *J* = 8.2 Hz, 1H, Ar), 8.83 (dd, *J* = 1.2, 4.8 Hz, 1H, Ar), 9.15 (d, *J* = 1.7 Hz, 1H, Ar), 10.55 (s, 1H, NH), 11.17 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 21.99, 30.93, 49.00, 119.06, 119.34, 123.91, 127.36, 129.17, 131.55, 135.24, 135.94, 136.20, 148.47, 152.95, 163.85, 166.59, 167.54,  
15 169.69, 172.72; Anal Calcd for C<sub>19</sub>H<sub>14</sub>N<sub>4</sub>O<sub>5</sub> + 0.13 H<sub>2</sub>O: C, 59.95; H, 3.78; N, 14.72; H<sub>2</sub>O, 0.62. Found: C, 59.83; H, 3.66; N, 14.68; H<sub>2</sub>O, 0.64.

3-{1-Oxo-4-[benzylamino]isoindolin-2-yl}piperidine-2,6-dione I-12

20


25



A stirred mixture of 3-(4-amino-1-oxoisoindolin-2-yl)piperidine-2,6-dione (518 mg, 2.0 mmol) and benzaldehyde (0.21 ml, 2.0 mmol) in methanol (20 ml) was heated to reflux for 5 h. The solvent was removed *in vacuo* to give a solid. The solid was re-dissolved in  
30 acetic acid (20 ml). The stirred solution was heated to reflux for 1h, and was then allowed to cool to room temperature. To the stirred solution was added sodium borohydride (90 mg, 2.3 mmol) and stirring continued at room temperature for 18h. The resulting suspension was filtered and washed with acetic acid (10 ml) and ether (20 ml) to give 3-{1-oxo-4-[benzylamino]isoindolin-2-yl}piperidine-2,6-dione as an off-white solid (420 mg,  
35 60% yield): mp, 257-259 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 2.01-2.07 (m, 1H, CHH), 2.23-2.30

(m, 1H, CHH), 2.49-2.65 (m, 1H, CHH), 2.85-3.00 (m, 1H, CHH), 4.19 (d,  $J = 17$  Hz, 1H, CHH), 4.31 (d,  $J = 17$  Hz, 1H, CHH), 4.39 (d,  $J = 5.7$  Hz, 2H, CH<sub>2</sub>), 5.12 (dd,  $J = 5.1$ , 13 Hz, 1H, NCH), 6.37 (t,  $J = 5.9$  Hz, 1H, NH), 6.62 (d,  $J = 8.0$  Hz, 1H, Ar), 6.91 (d,  $J = 7.3$  Hz, 1H, Ar), 7.19-7.40 (m, 6H, Ar), 11.02 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>)  $\delta$  22.79, 31.25, 45.79, 46.09, 51.56, 110.32, 112.34, 126.72, 127.05, 128.32, 129.07, 132.08, 139.72, 143.33, 168.82, 171.22, 172.90; Anal Calcd for C<sub>20</sub>H<sub>19</sub>N<sub>3</sub>O<sub>3</sub>: C, 68.75; H, 5.48; N, 12.03. Found C, 68.73; H, 5.41; N, 12.04.

2-(2,6-Dioxo(3-piperidyl))-4-[benzylamino]isoindoline-1,3-dione I-13




A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.0 g, 3.7 mmol) and benzaldehyde (0.4 ml, 3.9 mmol) in acetic acid (20 ml) was stirred at room temperature for 17h, then was heated to reflux for 3h. The mixture was cooled to room temperature. To the stirred mixture was added sodium borohydride (140 mg, 3.7 mmol) and kept at room temperature for 18h. The mixture was then heated to reflux for 2h. To the mixture was added additional benzaldehyde (0.4 ml, 3.9 mmol) during reflux. After 30 min of reflux the reaction was allowed to cool to room temperature. To the mixture was added sodium borohydride (180 mg, 4.8 mmol) and the mixture stirred at room temperature for 3 days. The solvent was removed *in vacuo* to yield an oil. The oil was diluted with ethyl acetate (90 ml) and aqueous sodium hydrogen carbonate (sat, 100 ml). The organic layer was separated and was washed with aqueous sodium hydrogen carbonate (sat, 2 × 100 ml), brine (100 ml) and dried over MgSO<sub>4</sub>. The solvent was removed *in vacuo* to give a solid. The solid was purified by column chromatography (Silca Gel, 50% EtOAc:CH<sub>2</sub>Cl<sub>2</sub>) to give a yellow solid. The solid was further purified by column chromatography (KP-C18-HS, 35:65 CH<sub>3</sub>CN:0.1% CF<sub>3</sub>COOH in water) to give 2-(2,6-dioxo(3-piperidyl))-4-[benzylamino]isoindoline-1,3-dione as a yellow solid (210 mg, 16% yield): mp, 209-211 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  2.02-2.08 (m, 1H, CHH), 2.46-2.63 (m, 2H, CH<sub>2</sub>), 2.82-2.97 (m, 1H, CHH), 4.56 (d,  $J = 6.2$  Hz, 2H, CH<sub>2</sub>), 5.07 (dd,  $J = 5.3$ , 12.4 Hz, 1H, NCH), 6.96 (d,  $J = 8.6$  Hz, 1H, Ar), 7.02 (d,  $J = 7.0$  Hz, 1H, Ar), 7.19-7.40 (m, 6H, Ar, NH), 7.51 (dd,  $J$

= 7.5, 8.4 Hz, 1H, Ar), 11.11 (s, 1H, NH);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  22.15, 30.99, 45.44, 48.59, 109.58, 110.74, 117.63, 126.95, 126.99, 128.53, 132.21, 136.09, 138.95, 146.09, 167.27, 168.78, 170.07, 172.79; Anal Calcd for  $\text{C}_{20}\text{H}_{17}\text{N}_3\text{O}_4$ : C, 66.11; H, 4.72; N, 11.56. Found: C, 65.96; H, 4.60; N, 11.49.

5

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}propanamide I-14

10



15


1,8-Diazabicyclo[5,4,0]undec-7-ene (0.65 g, 4.25 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 ml). After stirring for 20 min, propionyl chloride (0.2 g, 2.13 mmol) was added. The mixture was stirred at room temperature for 17 hours. Solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (60 ml) and washed with 1N HCl (30 ml),  $\text{H}_2\text{O}$  (30 ml), brine (30 ml) and dried ( $\text{MgSO}_4$ ). The solvent was removed and the resulting solid was slurried in hot  $\text{C}_2\text{H}_5\text{OH}$  (10 ml) to give after filtration N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}propanamide (0.41 g, 64%) as a white solid: mp 219-221°C;  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  d 11.15 (s, 1H), 8.42 (t,  $J=5.8$  Hz, 1H), 7.87-7.67 (m, 3H), 5.19-5.12 (dd,  $J=5.3$  and 12.5 Hz, 1H), 4.72 (d,  $J=5.8$  Hz, 2H), 2.98-2.84 (m, 1H), 2.65-2.48 (m, 2H), 2.26-2.17 (m, 2H), 2.09-2.04 (m, 1H), 1.05 (t,  $J=7.8$  Hz, 3H);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  d 173.52, 172.81, 169.79, 167.53, 167.01, 139.52, 134.79, 133.20, 131.55, 127.10, 121.86, 48.89, 37.69, 30.95, 28.43, 22.03, 9.90; Anal. Calcd. For  $\text{C}_{17}\text{H}_{17}\text{N}_3\text{O}_5 + 0.19 \text{H}_2\text{O}$ : C, 58.88; H, 5.05; N, 12.12. Found: C, 58.77; H, 4.97; N, 12.12.

35

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-3-pyridylcarboxamide I-15

5


10



15

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.98 g, 6.48 mmol) was added to stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 ml). After stirring for 20 min, nicotinoyl chloride (0.41 g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 ml). The  $\text{CH}_2\text{Cl}_2$  solution was washed with  $\text{H}_2\text{O}$  (30 ml), brine (30 ml) and dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* and the residue was purified by flash chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ :  $\text{CH}_3\text{OH}$  97.5:2.5) to give N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-3-pyridylcarboxamide (0.47 g, 64%) as a white solid: mp 148-151°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  d 11.16 (s, 1H), 9.36 (t,  $J=5.6$  Hz, 1H), 9.09 (d,  $J=1.25$  Hz, 1H), 8.75-8.73 (m, 1H), 8.25 (d,  $J=8.0$  Hz, 1H), 7.84-7.76 (m, 3H), 7.57-7.52 (m, 1H), 5.22-5.15 (dd,  $J=5.4$  and 12.7 Hz, 1H), 4.96 (d,  $J=5.6$  Hz, 2H), 2.92-2.85 (m, 1H), 2.65-2.50 (m, 2H), 2.11-2.06 (m, 1H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  d 172.72, 169.80, 167.51, 166.94, 165.23, 152.02, 148.44, 138.86, 135.10, 134.83, 133.20, 131.55, 129.48, 127.20, 123.49, 121.95, 48.89, 38.33, 30.93, 21.98; Anal. Calcd. For  $\text{C}_{20}\text{H}_{16}\text{N}_4\text{O}_5 + 0.28 \text{H}_2\text{O}$ : C, 60.45; H, 4.20; N, 14.10. Found: C, 60.29; H, 4.28; N, 13.82.

35


N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}heptanamide I-16

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.65 g, 2.22 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in CH<sub>3</sub>CN (50 ml). After stirring for 20 min, heptanoyl chloride (0.33 g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 ml). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 ml), H<sub>2</sub>O (30 ml), brine (30 ml) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: EtOAc 7: 3) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}heptanamide (0.49 g, 66%) as a white solid: mp 130-132°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ d 11.14 (s, 1H), 8.44 (t, J=5.7 Hz, 1H), 7.86-7.78 (m, 2H), 7.71-7.65 (m, 1H), 5.19-5.12 (dd, J=5.2 and 12.4 Hz, 1H), 4.69 (d, J=5.7 Hz, 2H), 2.98-2.83 (m, 1H), 2.64-2.50 (m, 2H), 2.18 (t, J=7.3 Hz, 2H), 2.08-2.04 (m, 1H), 1.53 (t, J=6.0 Hz, 2H), 1.25 (s, 6H), 0.85 (t, J=5.9 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ d 172.73, 172.66, 169.79, 167.47, 166.93, 139.54, 134.66, 133.13, 131.50, 127.06, 121.80, 54.86, 48.85, 37.57, 35.23, 30.96, 28.31, 25.16, 21.97, 13.87; Anal. Calcd. For C<sub>21</sub>H<sub>25</sub>N<sub>3</sub>O<sub>5</sub> + 0.3 H<sub>2</sub>O: C, 62.30; H, 6.37; N, 10.38. Found : C, 62.07; H, 6.29; N, 10.23.

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-2-furylcarboxamide I-17

5

10




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.650 g, 2.22 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.600 g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 ml). After stirring for 20 min, 2-furoyl chloride (0.290 g, 2.22 mmol) was added and the mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 ml). The  $\text{CH}_2\text{Cl}_2$  solution was washed 1N HCl (30 ml),  $\text{H}_2\text{O}$  (30 ml), brine (30 ml) and dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ : EtOAc 1: 1) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-2-furylcarboxamide (0.51 g, 73%) as a white solid: mp 121-123°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.13 (s, 1H), 9.01 (t,  $J=5.7$  Hz, 1H), 7.88-7.68 (m, 4H), 7.18 (d,  $J=3.3$  Hz, 1H), 6.66 (m, 1H), 5.20-5.13 (dd,  $J=5.4$  and 12.5 Hz, 1H), 4.90 (d,  $J=5.6$  Hz, 2H), 2.97-2.84 (m, 1H), 2.65-2.49 (m, 2H), 2.10-1.98 (m, 1H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.73, 169.80, 167.52, 166.94, 158.09, 147.49, 145.27, 139.06, 134.80, 132.98, 131.52, 127.09, 121.88, 113.86, 111.91, 48.86, 37.64, 30.92, 21.97; Anal. Calcd. For  $\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_6 + 0.18 \text{H}_2\text{O}$ : C, 59.34; H, 4.03; N, 10.93. Found: C, 59.47; H, 4.16; N, 10.49.

30

35

2-Azido-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-acetamide I-18




10

A mixture of 2-chloro-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]acetamide (1.3 g, 3.57 mmol), sodium azide (0.3 g, 4.65 mmol) and sodium iodide (0.54 g, 3.57 mmol) in acetone (50 ml) was refluxed for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in EtOAc (60 ml). The EtOAc solution was washed with H<sub>2</sub>O (30 ml), brine (30 ml), and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* to give 2-azido-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]acetamide (1.2 g, 90%) as a white solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ d 8.48 (s, 1H), 7.84-7.68 (m, 3H), 7.51 (t, J=6.3 Hz, 1H), 5.04-4.97 (dd, J=4.7 and 11.8 Hz, 1H), 4.80 (d, J=6.5 Hz, 2H), 3.97 (s, 2H), 2.95-2.74 (m, 3H), 2.20-2.15 (m, 1H).

20

2-Amino-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]acetamide I-19, hydrochloride



25

30

A mixture of 2-azido-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]acetamide (1.2 g, 3.24 mmol), and 10% Pd/C (0.15 g) in 4N HCl (20 ml) and CH<sub>3</sub>OH (50 ml) was hydrogenated in Parr shake apparatus under 50 psi of hydrogen for 3

35




hours. The mixture was then filtered through celite and the filtrate was concentrated to a solid residue. The solid was slurried in ethanol (20 ml) and the suspension filtered to give 2-amino-*N*-{[2-(2,6-dioxo(3-piperidyl))1,3-dioxoisindolin-4-yl]methyl}acetamide hydrochloride (0.86 g, 84%) as a white solid: mp 270-272°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ d 11.16 (s, 1H), 9.28 (t, J=5.7 Hz, 1H), 8.33 (s, 3H), 7.83 (s, 3H), 5.20-5.13 (dd, J=5.3 and 12.5 Hz, 1H), 4.81 (d, J=5.7 Hz, 2H), 3.69 (s, 2H), 2.92-2.87 (m, 1H), 2.65-2.51 (m, 2H), 2.09-2.05 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ d 172.76, 169.80, 167.44, 166.91, 166.52, 138.24, 134.74, 133.53, 131.51, 127.17, 122.08, 48.89, 37.89, 30.94, 21.99; Anal. Calcd. For C<sub>16</sub>H<sub>17</sub>N<sub>4</sub>O<sub>5</sub>Cl: C, 50.47; H, 4.50; N, 14.71; Cl, 9.31. Found: C, 50.39; H, 4.61; N, 14.42; Cl, 9.25.

Ethyl 6-(*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)hexanoate I-20

15


20



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.65 g, 4.26 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in CH<sub>3</sub>CN (50 ml). After stirring for 20 min, 6-(chloroformyl)hexanoic acid ethyl ester (0.46 g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 ml). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 ml), H<sub>2</sub>O (30 ml), brine (30 ml) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: EtOAc 1:1) to give ethyl 6-(*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)hexanoate (0.43 g, 50%) as a white solid: mp 82-84°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ d 11.11 (s, 1H), 8.41 (t, J=5.6 Hz, 1H), 7.86-7.65 (m, 3H), 5.18-5.11 (dd, J=5.4 and 12.4 Hz, 1H), 4.72 (d, J=5.7

Hz, 2H), 4.05 (q, J=7.1 Hz, 2H), 2.97-2.83 (m, 1H), 2.64-2.48 (m, 2H), 2.30-2.15 (m, 4H), 2.08-2.04 (m, 1H), 1.56-1.47 (m, 4H), 1.32-1.23 (m, 2H), 1.17 (t, J=7.1 Hz, 3H);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  d 172.80, 172.71, 172.52, 169.77, 167.45, 166.92, 139.49, 134.67, 133.13, 131.49, 127.04, 121.78, 59.60, 48.84, 37.57, 35.02, 33.38, 30.91, 28.08, 24.83, 24.16, 21.96, 14.09; Anal. Calcd. For  $\text{C}_{23}\text{H}_{27}\text{N}_3\text{O}_7$ : C, 60.39; H, 5.95; N, 9.18. Found: C, 60.10; H, 5.82; N, 8.82.


3-[(*tert*-Butoxy)carbonylamino]-*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}propanamide I-21



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.7 g, 4.62 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 ml). After stirring for 20 min, 1-hydroxybenzotriazole (0.3 g, 2.22 mmol), *N*-BOC-b-alanine (0.42 g, 2.22 mol) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.53 g, 2.78 mmol) were added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 ml). The  $\text{CH}_2\text{Cl}_2$  solution was washed with 1N citric acid (30 ml),  $\text{H}_2\text{O}$  ( $2 \times 30$  ml), brine (30 ml) and dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ :  $\text{CH}_3\text{OH}$  100:2) to give 3-[(*tert*-butoxy)carbonylamino]-*N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}propanamide (0.57 g, 67%) as a white solid: mp 96-98°C;  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  d 11.14 (s, 1H), 8.50 (t, J=5.8 Hz, 1H), 7.82-7.67 (m, 3H), 6.81 (t, J=5.1 Hz, 1H), 5.19-5.11 (dd, J=5.4 and 12.4 Hz, 1H), 4.72 (d, J=5.7 Hz, 2H), 3.21-3.13 (dd, J=6.8 and 13.1 Hz, 2H), 2.92-2.85 (m, 1H), 2.64-2.33 (m, 4H), 2.08-2.04 (m, 1H), 1.37 (s, 9H);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  d 172.71, 170.87, 169.77, 167.46, 166.93,

155.45, 139.27, 134.67, 133.17, 131.47, 127.03, 121.80, 77.57, 48.84, 37.63, 36.69, 35.62, 30.91, 28.21, 21.96; Anal. Calcd. For  $C_{22}H_{26}N_4O_7 + 0.28 H_2O$ : C, 57.01; H, 5.78; N, 12.09. Found: C, 56.99; H, 5.89; N, 11.79.

5 3-Amino-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]propanamide I-22, hydrochloride



15 A 4N HCl solution in dioxane (1 ml) was added to a stirred solution of 3-[(*tert*-butoxy)carbonylamino]-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]propanamide (0.5 g, 1.09 mmol) in  $CH_2Cl_2$  (15 ml) and stirred for 17 hours. The resulting suspension was filtered to give 3-amino-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]propanamide hydrochloride (0.34 g, 79%) as a white solid: mp 161-163°C;  $^1H$  NMR ( $DMSO-d_6$ )  $\delta$  d 11.15 (s, 1H), 8.88 (t,  $J=5.8$  Hz, 1H), 8.06 (b, 3H), 7.87-7.79 (m, 3H), 5.19-5.12 (dd,  $J=5.3$  and 12.6 Hz, 1H), 4.76 (d,  $J=5.8$  Hz, 2H), 3.03-2.84 (m, 3H), 2.67-2.47 (m, 4H), 2.08-2.04 (m, 1H);  $^{13}C$  NMR ( $DMSO-d_6$ )  $\delta$  d 172.74, 169.88, 169.80, 167.45, 166.92, 138.92, 134.74, 133.46, 131.48, 127.09, 121.92, 48.86, 37.69, 35.11, 32.03, 30.92, 21.97; Anal. Calcd. For  $C_{17}H_{19}N_4O_5Cl + 0.13 CH_2Cl_2 + 0.57 H_2O$ : C, 49.44; H, 4.94; N, 13.46; Cl, 10.73. Found: C, 49.22; H, 4.88; N, 13.08; Cl, 10.95.


20

25

30

35

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}2-thienylcarboxamide I-  
23




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.620 g; 4.07 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))-isindoline-1,3-dione hydrochloride (0.600 g, 1.85 mmol) in CH<sub>3</sub>CN (50 ml). After stirring for 20 min, 2-thiophene-carbonyl chloride (0.3 g, 2.03 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 ml). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 ml), H<sub>2</sub>O (30 ml), brine (30 ml) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: EtOAc 6:4) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-2-thienylcarboxamide (0.35 g, 47%) as a white solid: mp 192-194°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ d 11.16 (s, 1H), 9.18 (t, J=5.8 Hz, 1H), 7.88-7.72 (m 5H), 7.20-7.17 (dd, J=3.9 and 4.7 Hz, 1H), 5.22-5.15 (dd, J=5.5 and 12.7 Hz, 1H), 4.94 (d, J=5.8 Hz, 2H), 2.98-2.85 (m, 1H), 2.66-2.50 (m, 2H), 2.11-2.06 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ d 172.72, 169.80, 167.51, 166.93, 161.52, 139.26, 139.14, 134.83, 133.14, 131.53, 131.11, 128.51, 127.96, 127.12, 121.93, 48.89, 38.09, 30.93, 21.99; Anal. Calcd. For C<sub>19</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub>S: C, 57.42; H, 3.80; N, 10.57; S, 8.07. Found: C, 57.80; H, 3.93; N, 10.22; S, 7.99.

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-2-methoxyacetamide I-  
24

5

10



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62 g, 4.07 mmol) was added to a stirred  
 15 suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione  
 hydrochloride (0.600 g, 1.85 mmol) in CH<sub>3</sub>CN (50 ml). After stirring for 20 min,  
 methoxyacetyl chloride (0.22 g, 2.03 mmol) was added. The mixture was stirred at room  
 temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved  
 in CH<sub>2</sub>Cl<sub>2</sub> (70 ml). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 ml), H<sub>2</sub>O (30 ml),  
 20 brine (30 ml) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was  
 purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 100:2.5) to give N-{[2-(2,6-dioxo(3-  
 piperidyl))-1,3-dioxoisindolin-4-yl]methyl}-2-methoxyacetamide (0.44 g, 66%) as a white  
 solid: mp 196-198°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ d 11.14 (s, 1H), 8.49 (t, J=6.1 Hz, 1H), 7.86-  
 7.79 (m, 2H), 7.68-7.65 (m, 1H), 5.19-5.12 (dd, J=5.3 and 12.5 Hz, 1H), 4.77 (d, J=6.1 Hz,  
 25 2H), 3.92 (s, 2H), 3.36 (s, 3H), 2.96-2.83 (m, 1H), 2.64-2.49 (m, 2H), 2.09-2.04 (m, 1H);  
<sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ d 172.71, 169.78, 169.58, 167.52, 166.94, 139.09, 134.70, 133.00,  
 131.51, 127.08, 121.82, 71.46, 58.70, 48.86, 37.47, 30.91, 21.95; Anal. Calcd. For  
 C<sub>17</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>: C, 56.82; H, 4.77; N, 11.69. Found: C, 57.02; H, 4.87; N, 11.36.


30

35

(N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)methyl acetate I-25

5

10



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62 g, 4.08 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.600 g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 ml). After stirring for 20 min, acetoxyacetyl chloride (0.28 g, 2.03 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 ml). The  $\text{CH}_2\text{Cl}_2$  solution was washed with 1N HCl (30 ml),  $\text{H}_2\text{O}$  (30 ml), brine (30 ml) and dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ :  $\text{CH}_3\text{OH}$  100:2.5) to give (N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)methyl acetate (0.54 g, 75%) as a white solid: mp 108-110°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  d 11.14 (s, 1H), 8.68 (t,  $J=6.0$  Hz, 1H), 7.87-7.79 (m, 2H), 7.71-7.65 (m, 1H), 5.19-5.12 (dd,  $J=5.3$  and 12.4 Hz, 1H), 4.77 (d,  $J=5.9$  Hz, 2H), 4.56 (s, 2H), 2.96-2.83 (m, 1H), 2.63-2.47 (m, 2H), 2.11 (s, 3H), 2.08-1.98 (m, 1H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  d 172.66, 169.96, 169.72, 167.46, 167.37, 166.36, 138.80, 134.68, 133.01, 132.47, 127.04, 121.87, 62.36, 48.84, 37.46, 30.38, 21.93, 20.49; Anal. Calcd. For  $\text{C}_{18}\text{H}_{17}\text{N}_3\text{O}_7 + 0.15\text{H}_2\text{O}$ : C, 55.43; H, 4.47; N, 10.77. Found: C, 55.43; H, 4.54; N, 10.44.


30

35

Ethyl 2-[(N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)amino]acetate I-26

5

10




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29 g, 1.9 mmol) was added to a stirred  
 15 suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isindoline-1,3-dione  
 hydrochloride (0.6 g, 1.85 mmol) in CH<sub>3</sub>CN (50 ml). After stirring for 20 min, ethyl  
 isocyanatacetate (0.29 g, 2.22 mmol) was added. The mixture was stirred at room  
 temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved  
 in CH<sub>2</sub>Cl<sub>2</sub> (70 ml). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 ml), H<sub>2</sub>O (30 ml),  
 20 brine (30 ml) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was  
 purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: EtOAc 1:1) to give ethyl 2-[(N-{[2-(2,6-  
 dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}carbamoyl)amino]acetate (0.30 g,  
 39%) as a white solid: mp 187-189°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ d 11.14 (s, 1H), 7.86-7.70  
 (m, 3H), 6.83 (t, J=6.1 Hz, 1H), 6.53 (t, J=6.0 Hz, 1H), 5.18-5.11 (dd, J=5.4 and 12.5 Hz,  
 25 1H), 4.65 (d, J=6.0 Hz, 2H), 4.09 (q, J=7.2 Hz, 2H), 3.79 (d, J=6.0 Hz, 2H), 2.98-2.83 (m,  
 1H), 2.64-2.48 (m, 2H), 2.08-2.04 (m, 1H), 1.18 (t, J=7.2 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ  
 d 172.72; 171.01, 169.78, 167.54, 167.00, 158.00, 140.78, 134.57, 133.25, 131.48, 126.95,  
 121.66, 60.17, 48.83, 41.58, 38.72, 30.91, 21.97, 14.06; Anal. Calcd. For C<sub>19</sub>H<sub>20</sub>N<sub>4</sub>O<sub>7</sub>: C,  
 54.81; H, 4.84; N, 13.46. Found: C, 54.73; H, 4.77; N, 13.35.

30

35

*N*-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(ethylamino)carboxamide  
**I-27**




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29 g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.60 g, 1.85 mmol) in CH<sub>3</sub>CN (50 ml). After stirring for 20 min, ethyl isocyanate (0.16 g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 ml). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCL (30 ml), H<sub>2</sub>O (30 ml), brine (30 ml) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>CN 6:4) to give *N*-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(ethylamino)carboxamide (0.2 g, 30%) as a white solid: mp 173-175 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ d 11.13 (s, 1H), 7.86-7.69 (m, 3H), 6.44 (t, J=6.1 Hz, 1H), 6.11 (t, J=5.55 Hz, 1H), 5.18-5.11 (dd, J=5.4 and 12.6 Hz, 1H), 4.63 (d, J=6.1 Hz, 2H), 3.07-2.83 (m, 3H), 2.64-2.49 (m, 2H), 2.08-2.04 (m, 1H), 0.99 (t, J=7.1 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ d 172.73, 169.80, 167.58, 167.03, 157.93, 141.15, 134.61, 133.37, 131.49, 126.96, 121.61, 48.83, 38.71, 34.17, 30.92, 21.98, 15.58; Anal. Calcd. For C<sub>17</sub>H<sub>18</sub>N<sub>4</sub>O<sub>5</sub> + 0.14 H<sub>2</sub>O: C, 56.58; H, 5.11; N, 15.53. Found: C, 56.56; H, 5.05; N, 15.28.



2-(2,6-Dioxo(3-piperidyl))-4-[(2-furylmethyl)amino]isoindoline-1,3-dione I-28

5




10

To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in acetic acid (20 ml) was added furan-2-carbaldehyde (0.20 g, 2.05 mmol). The mixture was heated to reflux for 5 hours and allowed to cool at room temperature. Sodium borohydride (80 mg, 2 mmol) was added to the reaction mixture. The reaction mixture was stirred for 24 hours. The solvent was evaporated *in vacuo* and the residue was dissolved in ethyl acetate (100 ml), washed with H<sub>2</sub>O (3 × 100 ml), saturated aqueous NaHCO<sub>3</sub> (2 × 100 ml), brine (1 × 100 ml), and dried. The solvent was evaporated and the residue was purified by chromatography (ethyl acetate/hexane, 1:1) to give 0.25 g (35%) of product as a yellow solid: mp 171-173 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.10 (s, 1H), 7.58 (t, J=7.1 Hz, 2H), 7.18 (d, J=8.6 Hz, 1H), 7.08-6.97 (m, 2H), 6.39-6.36 (m, 2H), 5.06 (dd, J=5.2 and 12.4 Hz, 1H), 4.57 (d, J=6.0 Hz, 2H), 2.96-2.82 (m, 1H), 2.63-2.46 (m, 2H), 2.06-2.02 (m, 1H; <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.76, 170.01, 168.75, 167.21, 151.97, 145.81, 142.42, 136.05, 132.09, 117.60, 111.04, 110.42, 109.73, 107.38, 48.57, 30.95, 22.10; Anal. Calcd. For C<sub>18</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub> : C, 61.19; H, 4.28; N, 11.89. Found: C, 61.02; H, 4.24; N, 11.81.

25

N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-methoxyacetamide I-29


30




35

To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added 2-methoxyacetyl chloride (0.43 g, 4.0 mmol). The stirred mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.69 g (100 %) of product as an off-white solid: mp 246-248 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.15 (s, 1H), 10.30 (s, 1H), 8.70 (d, J=8.4 Hz, 1H), 7.86 (t, J=7.8 Hz, 1H), 7.63 (d, J=7.3 Hz, 1H), 5.17 (dd, J=5.2 and 12.7 Hz, 1H), 4.11 (s, 2H), 3.49 (s, 3H), 2.98-2.84 (m, 1H), 2.66-2.47 (m, 2H), 2.12-2.07 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.67, 169.66, 169.02, 168.25, 166.61, 136.48, 135.91, 131.24, 124.35, 118.31, 116.04, 71.40, 59.10, 48.96, 30.91, 21.94; Anal. Calcd. For C<sub>16</sub>H<sub>15</sub>N<sub>3</sub>O<sub>6</sub> : C, 55.65; H, 4.38; N, 12.17. Found: C, 55.58; H, 4.40; N, 12.08.

*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]heptanamide I-30



To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added heptanoyl chloride (0.59 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.61 g (79 %) of product as an off-white solid: mp 200-202 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.16 (s, 1H), 9.65 (s, 1H), 8.48 (d, J=8.3 Hz, 1H), 7.81 (t, J=7.7 Hz, 1H), 7.58 (d, J=7.2 Hz, 1H), 5.15 (dd, J=4.8 and 12.2 Hz, 1H), 2.97-2.87 (m, 1H), 2.65-2.43 (m, 4H), 2.10-2.06 (m, 1H), 1.61-1.58 (m, 2H), 1.28 (bs, 6H), 0.85 (bs, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.67, 171.96, 169.69, 167.78, 166.61, 136.62, 136.04, 131.36, 125.97, 118.13, 116.67, 48.93, 36.56, 30.95, 28.16, 24.69, 21.94, 13.84; Anal. Calcd. For C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O<sub>5</sub> : C, 62.33; H, 6.02; N, 10.90. Found: C, 62.14; H, 6.05; N, 10.72.

{N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]carbamoyl}methyl acetate I-31

10


To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added (chlorocarbonyl)methyl acetate (0.55 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.56 g (75 %) of product as an off-white solid: mp 234-236 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.15 (s, 1H), 10.06 (s, 1H), 8.55 (d, J=8.3 Hz, 1H), 7.87 (t, J=8.0 Hz, 1H), 7.65 (d, J=7.2 Hz, 1H), 5.17 (dd, J=5.0 and 12.4 Hz, 1H), 4.78 (s, 2H), 2.99-2.85 (m, 1H), 2.65-2.52 (m, 2H), 2.22 (s, 2H), 2.11-2.07 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.70, 169.69, 169.50, 167.99, 166.57, 136.40, 135.59, 131.35, 125.34, 118.72, 116.96, 62.60, 48.98, 30.90, 21.91, 20.41; Anal.

15

20 Calcd. For C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O<sub>7</sub> : C, 54.69; H, 4.05; N, 11.26. Found: C, 54.43; H, 4.05; N, 10.97.

N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]pentanamide I-32

25




To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added pentanoyl chloride (0.48 g, 4.0 mmol). The stirred mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.61 g (85 %) of


35

product as an off-white solid: mp 178-179 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.17 (s, 1H), 9.65 (s, 1H), 8.50 (d, J=8.3 Hz, 1H), 7.82 (t, J=7.5 Hz, 1H), 7.59 (d, J=7.2 Hz, 1H), 5.18 (dd, J=5.0 and 12.4 Hz, 1H), 2.99-2.89 (m, 1H), 2.68-2.45 (m, 4H), 2.13-2.09 (m, 1H), 1.69-1.57 (m, 2H), 1.44-1.30 (m, 2H), 0.92 (t, J=7.2 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.72, 171.99, 169.73, 167.83, 166.64, 136.64, 136.07, 131.36, 125.96, 118.15, 116.68, 48.97, 36.32, 30.97, 26.88, 22.03, 21.70, 13.64; Anal. Calcd. For C<sub>18</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub> : C, 60.50; H, 5.36; N, 11.76. Found: C, 60.10; H, 5.37; N, 11.58.

N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-thienylcarboxamide I-33




To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55g, 2.0 mmol) in THF (30 ml) was added thiophene-2-carbonyl chloride (0.59 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. To the reaction mixture was added additional thiophene-2-carbonyl chloride (0.30 g, 2 mmol). The mixture was heated to reflux for an additional 8 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give an off-white solid which was recrystallized from acetic acid to give 0.50 g (65 %) of product: mp 284-286 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.19 (s, 1H), 10.34 (s, 1H), 8.47(d, J=8.3 Hz, 1H), 7.98-7.86 (m, 3H), 7.68 (d, J=7.3 Hz, 1H), 7.29 (dd, J=4.0 and 4.8 Hz, 1H), 5.19 (dd, J=5.4 and 12.6 Hz, 1H), 2.99-2.85 (m, 1H), 2.67-2.49(m, 2H), 2.12-2.08 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.72, 169.69, 167.96, 166.60, 159.67, 138.02, 136.26, 136.19, 133.26, 131.42, 129.91, 128.58, 126.42, 118.82, 117.94, 49.00, 30.93, 22.00; Anal. Calcd. For C<sub>18</sub>H<sub>13</sub>N<sub>3</sub>O<sub>5</sub>S : C, 55.77; H, 3.54; N, 10.60. Found: C, 55.67; H, 3.36; N, 10.42 + 0.2 AcOH + 0.05 H<sub>2</sub>O.

Methyl {N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]carbamoyl} formate I-34

10

To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55g, 2.0 mmol) in THF (30 ml) was added methyl (chlorocarbonyl)formate (0.49 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.55 g (76 %) of product as an off-white solid: mp 247-249 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.17 (s, 1H), 10.81 (s, 1H), 8.59 (d, J=8.3 Hz, 1H), 7.92 (t, J=7.7 Hz, 1H), 7.70 (d, J=7.3 Hz, 1H), 5.19 (dd, J=5.0 and 12.5 Hz, 1H), 3.91 (s, 3H), 2.98-2.85 (m, 1H), 2.68-2.50 (m, 2H), 2.14-2.09 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.70, 169.63, 168.03, 166.45, 159.81, 154.57, 136.65, 134.93, 131.36, 124.67, 119.34, 117.17, 53.80, 49.06, 30.92, 21.94; Anal. Calcd. For C<sub>16</sub>H<sub>13</sub>N<sub>3</sub>O<sub>7</sub> : C, 53.49; H, 3.65; N, 11.70. Found: C, 53.53; H, 3.66; N, 11.52.


N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-furancarboxamide I-35

30


To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added furan-2-carbonyl chloride (0.52 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. To the reaction mixture was added additional furan-2-carbonyl chloride (0.26 g, 2 mmol). The mixture was heated to reflux

for an additional 8 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.65 g (88 %) of product as an off-white solid: mp 299-301 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.19 (s, 1H), 10.37 (s, 1H), 8.68 (dd, J=1.4 and 8.4 Hz, 1H), 8.06 (s, 1H), 7.90 (t, J=8.1 Hz, 1H), 7.65 (d, J=7.3 Hz, 1H), 7.39 (d, J=3.5 Hz, 1H), 6.79 (dd, J=1.5 and 3.2 Hz, 1H), 5.19 (dd, J=5.1 and 12.5 Hz, 1H), 2.98-2.84 (m, 1H), 2.67-2.49(m, 2H), 2.13-2.08 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.66, 169.62, 168.44, 166.59, 155.71, 146.80, 146.34, 136.55, 136.23, 136.12, 131.30, 124.86, 118.54, 116.64, 113.06, 48.98, 30.90, 22.01; Anal. Calcd. For C<sub>18</sub>H<sub>13</sub>N<sub>3</sub>O<sub>6</sub> : C, 58.86; H, 3.57; N, 11.44. Found: C, 58.69; H, 3.54; N, 11.41.

N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]benzamide I-36




To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added benzoyl chloride (0.56 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. To the reaction mixture was added additional benzoyl chloride (0.28 g, 2.0 mmol). The mixture was heated at reflux for an additional 8 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give an of off-white solid which was recrystallized from acetic acid to give 0.49 g (65 %) of product: mp 268-270 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.18 (s, 1H), 10.43 (s, 1H), 8.64-8.59 (m, 1H), 8.00-7.89 (m, 3H), 7.72-7.60 (m, 4H), 5.19 (dd, J=5.2 and 12.5 Hz, 1H), 2.98-2.84 (m, 1H), 2.66-2.50(m, 2H), 2.12-2.07 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.62, 169.59, 168.16, 166.64, 164.91, 136.50, 136.36, 133.23, 132.67, 131.36, 129.06, 127.24, 126.00, 118.74, 117.74, 48.98, 30.89, 21.98; Anal. Calcd. For C<sub>20</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub> : C, 63.02; H, 4.08; N, 10.90. Found: C, 63.05; H, 4.06; N, 10.69 + 0.11 AcOH + 0.08 H<sub>2</sub>O.

N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]propanamide I-37

10

To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added propanoyl chloride (0.37 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.58 g (88 %) of product as an off-white solid: mp 221-223 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.15 (s, 1H), 9.65 (s, 1H), 8.50 (d, J=8.3 Hz, 1H), 7.83 (t, J=7.6 Hz, 1H), 7.60 (d, J=7.2 Hz, 1H), 5.16 (dd, J=5.2 and 12.6 Hz, 1H), 2.99-2.85 (m, 1H), 2.66-2.45 (m, 4H), 2.11-2.07(m, 1H), 1.14 (t, J=7.5 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.69, 172.65, 169.71, 167.79, 166.62, 136.65, 136.09, 131.35, 125.93, 118.12, 116.66, 48.91, 30.91, 29.73, 21.97, 9.13; Anal. Calcd. For C<sub>16</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub> : C, 58.36; H, 4.59; N, 12.76. Found: C, 58.01; H, 4.45; N, 12.61.


Methyl 3-{N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]carbamoyl}propanoate I-38

30

To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added methyl 3-(chlorocarbonyl)propanoate (0.63 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated

*in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.75 g (97 %) of product as an off-white solid: mp 224-226 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.18 (s, 1H), 9.79 (s, 1H), 8.43 (d, J=8.4 Hz, 1H), 7.83 (t, J=7.8 Hz, 1H), 7.62 (d, J=7.3 Hz, 1H), 5.16 (dd, J=5.2 and 12.6 Hz, 1H), 3.61 (s, 3H), 2.99-2.48 (m, 7H), 2.11-2.06 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.84, 172.66, 170.94, 169.75, 167.61, 166.70, 136.39, 136.15, 131.52, 126.39, 118.43, 117.04, 51.51, 48.95, 31.22, 30.96, 28.39, 22.03; Anal. Calcd. For C<sub>18</sub>H<sub>17</sub>N<sub>3</sub>O<sub>7</sub> : C, 55.82; H, 4.42; N, 10.85. Found: C, 55.68; H, 4.41; N, 10.61.

N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-phenylacetamide I-39




To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added 2-phenylacetyl chloride (0.62 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.72 g (92 %) of product as an off-white solid: mp 217-218 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.15 (s, 1H), 9.79 (s, 1H), 8.49 (d, J=8.4 Hz, 1H), 7.82 (t, J=7.9 Hz, 1H), 7.60 (t, J=7.3 Hz, 1H), 7.41-7.27 (m, 5H), 5.13 (dd, J=5.1 and 12.7 Hz, 1H), 3.85 (s, 2H), 2.98-2.83 (m, 1H), 2.64-2.44 (m, 2H), 2.08-2.04 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.71, 170.07, 169.74, 167.60, 166.61, 136.38, 136.10, 134.70, 131.41, 129.41, 128.58, 126.97, 125.96, 118.39, 116.90, 48.89, 43.47, 30.90, 21.91; Anal. Calcd. For C<sub>21</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub> : C, 64.45; H, 4.38; N, 10.74. Found: C, 64.23; H, 4.34; N, 10.53.



N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-pyridylcarboxamide I-40

5




10

To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isindoline-1,3-dione (0.55 g, 2.0 mmol) in THF (30 ml) was added pyridine-2-carbonyl chloride hydrochloride (0.71 g, 4.0 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* and the resulting solid was slurried in a biphasic mixture of diethyl ether (20 ml)/20 % NH<sub>4</sub>OH (20 ml) and filtered to give an off-white solid. The solid was re-slurried in methanol (20 ml) and filtered to give 0.30 g (40 %) of product: mp 336-338 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.83 (s, 1H), 11.20 (s, 1H), 8.92 (d, J=8.4 Hz, 1H), 8.81 (d, J=3.6 Hz, 1H), 8.22 (d, J=7.7 Hz, 1H), 8.15 (t, J=7.4 Hz, 1H), 7.92 (t, J=7.7 Hz, 1H), 7.77-7.72 (m, 1H), 7.65 (d, J=7.2 Hz, 1H), 5.22 (dd, J=5.2 and 12.5 Hz, 1H), 3.00-2.86 (m, 1H), 2.69-2.52 (m, 2H), 2.15-2.11 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.67, 169.67, 168.20, 166.61, 162.69, 148.83, 148.19, 138.45, 136.50, 136.16, 131.39, 127.72, 124.23, 122.46, 118.29, 116.32, 48.96, 30.91, 21.98; Anal. Calcd. For C<sub>19</sub>H<sub>14</sub>N<sub>4</sub>O<sub>5</sub> : C, 60.32; H, 3.73; N, 14.81. Found: C, 60.05; H, 3.57; N, 14.45.

25

N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-chloroacetamide I-41

30




35

To a stirred suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.37 g, 5.00 mmol) in THF (30 ml) was added chloroacetyl chloride (0.62 g, 5.5 mmol). The mixture was heated to reflux for 30 minutes. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 1.67 g (96 %) of product as an off-white solid: <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.18 (s, 1H), 10.31 (s, 1H), 8.54 (d, J=8.4 Hz, 1H), 7.88 (t, J=7.7 Hz, 1H), 7.68 (d, J=7.3 Hz, 1H), 5.17 (dd, J=5.2 and 12.7 Hz, 1H), 4.54 (s, 2H), 2.90-2.85 (m, 1H), 2.65-2.51 (m, 2H), 2.10-2.06 (m, 1H).

2-azido-N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]acetamide I-42

10

15




To a suspension of N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-chloroacetamide (1.53 g, 4.4 mmol) in acetone (30 ml) was added sodium azide (0.43 g, 6.6 mmol). The mixture was heated to reflux for 18 hours. The solvent was evaporated *in vacuo* to give 1.49 g (96 %) of product as an off-white solid: <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.19 (s, 1H), 10.20 (s, 1H), 8.49 (d, J=8.3 Hz, 1H), 7.88 (t, J=7.7 Hz, 1H), 7.68 (d, J=7.3 Hz, 1H), 5.17 (dd, J=5.1 and 12.7 Hz, 1H), 4.34 (s, 2H), 2.99-2.84 (m, 1H), 2.65-2.47 (m, 2H), 2.09-2.00 (m, 1H).

25

2-Amino-N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]acetamide I-43, hydrochloride


30

35




To a solution of 2-azido-*N*-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]acetamide (1.49 g, 4.2 mmol) in methanol (50 ml) was added 10% Pd-C (0.1 g). Hydrogenation at 50 psi of hydrogen in a Parr Type shaker for 2 hours yielded a slurry. The mixture was filtered leaving a gray solid that was stirred in H<sub>2</sub>O (50 ml). The pH of the aqueous mixture was adjusted to 4 by addition of 3N HCl. The aqueous mixture was filtered through celite to remove catalyst and the filtrate was stirred with 50 ml of ethyl acetate for 3 hours. The aqueous layer was separated and was evaporated *in vacuo* to give a solid which was slurried in ethyl acetate and filtered to give 0.72 g (45%) of product as an off-white solid: mp 305-307 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.17 (s, 1H), 10.35 (s, 1H), 8.44 (bs, 3H), 8.32 (d, J=8.2 Hz, 1H), 7.90 (t, J=8.0 Hz, 1H), 7.71 (d, J=7.2 Hz, 1H), 5.16 (dd, J=5.1 and 12.6 Hz, 1H), 3.97 (s, 2H), 2.99-2.84 (m, 1H), 2.65-2.46 (m, 2H), 2.10-2.06 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.81, 169.80, 166.75, 166.56, 166.19, 136.19, 134.91, 131.89, 127.62, 119.47, 118.68, 48.96, 41.13, 30.94, 22.00; Anal. Calcd. For C<sub>15</sub>H<sub>15</sub>ClN<sub>4</sub>O<sub>5</sub>: C, 48.62; H, 4.19; N, 15.12. Found: C, 48.68; H, 4.18; N, 15.05 + 0.21 H<sub>2</sub>O.

*N*-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]-2-chloroacetamide I-44




To a stirred suspension of 3-(4-amino-1-oxoisindolin-2-yl)piperidine-2,6-dione (3.89 g, 15.0 mmol) in THF (50 ml) was added chloroacetyl chloride (1.86 g, 16.5 mmol). The mixture was heated to reflux for 45 minutes. To the reaction mixture was added additional chloroacetyl chloride (0.15 g, 0.13 mmol). The reaction mixture was heated at reflux for an additional 30 minutes. The solvent was evaporated *in vacuo* and the resulting solid was slurried in diethyl ether (20 ml) and filtered to give 4.64 g (92 %) of product as an off-white solid: <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.04 (s, 1H), 10.22 (s, 1H), 7.82 (dd, J=1.6 and 7.2 Hz, 1H), 7.59-7.50 (m, 2H), 5.16 (dd, J=5.1 and 13.2 Hz, 1H), 4.46-4.30 (m, 4H), 3.00-2.85 (m, 1H), 2.65-2.58 (m, 1H), 2.44-2.28 (m, 1H), 2.06-2.01 (m, 1H).

2-azido-N-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]acetamide I-45

10 To a stirred suspension of N-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]-2-chloroacetamide (4.64 g, 13.8 mmol) in acetone (60 ml) was added sodium azide (1.35 g, 20.7 mmol). The mixture was heated to reflux for 18 hours. After 18 hours, to the reaction mixture was added NaI (2.05 g, 13.8 mmol) and additional sodium azide (0.90 g, 13.8 mmol). The mixture was heated at reflux for an additional 18 hours. The solvent was

15 evaporated *in vacuo* to give an off-white solid which was slurried in a mixture of dichloromethane (50 ml) and H<sub>2</sub>O (50 ml). This slurry was filtered to give 4.39 g (93 %) of product: <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.50-9.52 (bs, 2H), 7.87-7.84 (m, 1H), 7.59-7.50 (m, 2H), 5.17 (dd, J=5.0 and 13.1 Hz, 1H), 4.44 (d, J=17.6 Hz, 1H), 4.34 (d, J=17.6 Hz, 1H), 4.13 (s, 2H), 3.00-2.86 (m, 1H), 2.65-2.59 (m, 1H), 2.44-2.29 (m, 1H), 2.07-2.02 (m, 1H).

20

2-Amino-N-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]acetamide I-46, hydrochloride

30

To a stirred suspension of 2-azido-N-[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]acetamide (1.49 g, 4.20 mmol) in a mixture of methanol (50 ml) and 3N HCl (6 ml) was added 10% Pd-C (0.1 g). Hydrogenation at 50 psi of hydrogen in a Parr Type shaker for 24 hours gave a slurry. The mixture was filtered leaving a gray solid that was stirred in H<sub>2</sub>O


35 (100 ml). The aqueous mixture was filtered through celite to remove catalyst. The aqueous

and methanolic filtrates were combined and evaporated *in vacuo* to give a white solid. The solid was slurried in ethyl acetate (20 ml), filtered, re-slurried in methanol (20 ml), and filtered to give 2.35 g (48 %) of product: mp 293-295 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.06 (s, 1H), 10.86 (s, 1H), 8.45 (bs, 2H), 7.90 (d, J=6.3 Hz, 1H), 7.59-7.51 (m, 2H), 5.17 (dd, J=4.9 and 13.0 Hz, 1H), 4.54 (d, J=17.8 Hz, 1H), 4.39 (d, J=17.8 Hz, 1H), 3.91 (s, 2H), 3.02-2.88 (m, 1H), 2.68-2.61 (m, 1H), 2.37-2.23 (m, 1H), 2.09-2.05 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.94, 171.07, 167.74, 165.24, 133.73, 132.86, 128.95, 125.01, 119.65, 51.59, 46.74, 40.90, 31.22, 22.82; Anal. Calcd. For C<sub>15</sub>H<sub>17</sub>ClN<sub>4</sub>O<sub>4</sub>: C, 50.43; H, 4.94; N, 15.68. Found: C, 50.08; H, 4.92; N, 15.53 + 0.25 H<sub>2</sub>O.

10

3-{4-[(2-Furylmethyl)amino]-1-oxoisindolin-2-yl}piperidine-2,6-dione I-47

15




20

To a stirred suspension of 3-(4-amino-1-oxoisindolin-2-yl)piperidine-2,6-dione (0.52 g, 2.0 mmol) in methanol (50 ml) was added furan-2-carbaldehyde (0.200 g, 2.05 mmol). The mixture was heated to reflux for 4 hours. The solvent was evaporated *in vacuo* and the residue was dissolved in acetic acid (20 ml). Sodium triacetoxymethylborohydride (0.450 g, 2.05 mmol) was added to the reaction mixture. The reaction mixture was stirred for 24 hours. The solvent was evaporated *in vacuo* and the residue was dissolved in ethyl acetate (100 ml), washed with H<sub>2</sub>O (3 × 100 ml), saturated aqueous NaHCO<sub>3</sub> (2 × 100 ml), brine (1 × 100 ml), and dried over MgSO<sub>4</sub>. The solvent was evaporated *in vacuo* and the residue was partially purified by chromatography (100 % ethyl acetate) to give an off-white solid which was recrystallized from ethanol to give 0.20 g of product as a white solid: mp 248-250 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.02 (s, 1H), 7.57 (s, 1H), 7.27 (t, J=7.6 Hz, 1H), 6.96 (d, J=7.3 Hz, 1H), 6.85 (d, J=8.0 Hz, 1H), 6.36 (d, J=11.2 Hz, 2H), 6.22 (t, J=5.4 Hz, 1H), 5.12 (dd, J=4.9 and 13.1 Hz, 1H), 4.37 (d, J=5.5 Hz, 2H), 4.27 (d, J=17.2 Hz, 1H), 4.14 (d, J=17.2 Hz, 1H) 3.00-2.86 (m, 1H), 2.65-2.59 (m, 1H), 2.37-2.23 (m, 1H), 2.08-1.99 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.83, 171.16, 168.70, 152.82, 143.06, 142.00, 132.09,

35

129.02, 126.76, 112.35, 110.63, 110.33, 107.08, 51.50, 45.73, 31.20, 22.75; Anal. Calcd. For  $C_{18}H_{17}N_3O_4$  : C, 63.71; H, 5.05; N, 12.38. Found: C, 63.41; H, 5.03; N, 11.98.


3-[1-Oxo-4-(pentylamino)isoindolin-2-yl]piperidine-2,6-dione I-48



To a stirred solution of 3-(4-amino-1-oxoisoindolin-2-yl)piperidine-2,6-dione (0.52 g, 2.0 mmol) in DMF (10 ml) was added pentanal (0.26 g, 3.0 mmol), acetic acid (0.24 g, 4.0 mmol), and sodium triacetoxyborohydride (0.85 g, 4.0 mmol). The reaction mixture was stirred at room temperature for 6 hours. The solvent was evaporated *in vacuo* and the residue was dissolved in ethyl acetate (100 ml), washed with  $H_2O$  ( $3 \times 100$  ml), brine ( $1 \times 100$  ml), and dried over  $MgSO_4$ . The solvent was evaporated and the residue was partially purified by chromatography (ethyl acetate/hexane, 75:25) to give an off-white solid which was slurried in ethyl acetate and filtered to give 0.14 g (21 %) of product as a white solid: mp 244-246 °C;  $^1H$  NMR ( $DMSO-d_6$ )  $\delta$  11.00 (s, 1H), 7.28 (t,  $J=7.7$  Hz, 1H), 6.92 (d,  $J=7.3$  Hz, 1H), 6.73 (d,  $J=8.0$  Hz, 1H), 5.54 (t,  $J=5.3$  Hz, 1H), 5.12 (dd,  $J=5.1$  and 13.2 Hz, 1H), 4.24 (d,  $J=17.2$  Hz, 1H), 4.13 (d,  $J=17.2$  Hz, 1H) 3.15-3.07 (m, 2H), 3.00-2.86 (m, 1H), 2.65-2.59 (m, 1H), 2.39-2.23 (m, 1H), 2.08-1.99 (m, 1H), 1.61-1.57 (m, 2H), 1.41-1.32 (m, 4H), 0.89 (t,  $J=6.8$  Hz, 3H);  $^{13}C$  NMR ( $DMSO-d_6$ )  $\delta$  172.83, 171.19, 168.86, 143.74, 132.00, 129.15, 126.42, 111.66, 109.86, 51.46, 45.69, 42.68, 31.20, 28.80, 28.21, 22.79, 21.96, 13.89; Anal. Calcd. For  $C_{18}H_{23}N_3O_3$  : C, 65.63; H, 7.04; N, 12.76. Found: C, 65.69; H, 7.22; N, 12.55.

3-(2-Methoxy-ethylamino)-phthalic acid dimethyl ester

5




To a stirred solution of oxalyl chloride (1.75 ml, 20 mmol) in methylene chloride (20 ml) under a nitrogen atmosphere at  $-78^{\circ}\text{C}$  was added DMSO (1.42 ml, 20 mmol) in methylene chloride (10 ml) dropwise over 5 minutes. The mixture was stirred for 5 minutes followed by the dropwise addition of 2-methoxyethanol (1.58 ml, 20 mmol) in methylene chloride (10 ml) over 5 minutes. The mixture was stirred for 20 minutes followed by the dropwise addition of triethylamine (8.36 ml, 60 mmol) over 5 minutes. The resulting suspension was stirred for 30 minutes at  $-78^{\circ}\text{C}$  and then allowed to warm to room temperature. The reaction mixture was diluted with methylene chloride (20 ml). To this stirred mixture was added 3-amino-phthalic acid dimethyl ester (2.09 g, 10 mmol) and acetic acid (4.60 ml, 80 mmol). The mixture was stirred at room temperature under a nitrogen atmosphere for 5 minutes followed by the addition of sodium triacetoxyborohydride (4.24 g, 20 mmol). The mixture was stirred for 3 h. The reaction mixture was diluted with methylene chloride (50 ml) and washed with water (3x100 ml), saturated aqueous sodium bicarbonate (2x100 ml), brine (100 ml), and dried ( $\text{MgSO}_4$ ). The solvent was evaporated *in vacuo* and the residue purified by chromatography (25% ethyl acetate/hexane) to give 2.24 g (84%) of product as an oil.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  7.32 (t,  $J=8.0$  Hz, 1H), 6.88-6.79 (m, 3H), 3.85 (s, 3H), 3.83 (s, 3H), 3.62 (t,  $J=5.4$  Hz, 2H), 3.41 (s, 3H), 3.35 (q,  $J=5.2$  Hz, 2H).

30

35

3-(2-Methoxy-ethylamino)-phthalic acid


5



To a stirred solution of 3-(2-methoxy-ethylamino)-phthalic acid dimethyl ester (2.24 g, 8.38 mmol) in methanol (50 ml) was added 5N potassium hydroxide (10 ml). The mixture was stirred at room temperature overnight. The solvent was evaporated *in vacuo* and the residue dissolved in water (50 ml). The water was washed with diethyl ether (2 x 75 ml). The aqueous portion was cooled in an ice bath and the pH was adjusted to 2-3 by dropwise addition of concentrated hydrochloric acid. The aqueous solution was then extracted with ethyl acetate (3 x 75 ml). The combined ethyl acetate extracts were washed with brine (100 ml) and dried (MgSO<sub>4</sub>). The solvent was evaporated *in vacuo* and the residue, which contained a mixture of diacid and monomethyl esters, was used without further purification.

2-(2,6-Dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione I-49

25




To a stirred solution of 3-(2-methoxy-ethylamino)-phthalic acid (8.38 mmol) in pyridine (40 ml) was added 3-amino-piperidine-2,6-dione hydrochloride (1.39 g, 8.42 mmol). The reaction mixture was heated to reflux for 5 hours. The solvent was evaporated *in vacuo* and the residue dissolved in methylene chloride (125 ml). The methylene chloride mixture was treated with Norit (2 g), washed with water (2 x 100 ml), 0.1N HCl (1 x 100 ml), brine (1 x 100 ml), and dried (MgSO<sub>4</sub>). The solvent was evaporated *in vacuo* and the residue (oil) crystallized from a minimal amount of ethanol to give a yellow solid that was



purified by preparative HPLC to give 1.71 g (64%) of product as a yellow solid: mp 182-184°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.12 (s, 1H), 7.58 (t,  $J=7.7$  Hz, 1H), 7.12 (d,  $J=8.5$  Hz, 1H), 7.04 (d,  $J=6.9$  Hz, 1H), 6.59 (bs, 1H), 5.07 (dd,  $J=4.7$  and 12.1 Hz, 1H), 3.54-3.30 (m, 7H), 2.95-2.85 (m, 1 H), 2.64-2.52 (m, 2H), 2.07-2.02 (m, 1H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.79, 170.06, 168.98, 167.27, 146.38, 136.21, 132.07, 117.33, 110.67, 109.24, 70.39, 58.12, 48.58, 41.50, 30.99, 22.15; Anal. Calcd. For  $\text{C}_{16}\text{H}_{17}\text{N}_3\text{O}_5$ : C, 58.00; H, 5.17; N, 12.68. Found: C, 58.06; H, 5.12; N, 12.76.


10 2-Benzoyloxy-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-acetamide I-50



To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.10 g, 4 mmol) in THF (30 ml) was added benzyloxyacetyl chloride (1.26 ml, 8 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* and the residue was slurried in diethyl ether (30 ml), filtered, recrystallized from a minimal amount of acetic acid, slurried in ethyl acetate (15 ml), and filtered to give 1.35 g (80%) of product: mp 204-206 °C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.20 (s, 1H), 10.40 (s, 1H), 8.71 (d,  $J=8.4$  Hz, 1H), 7.86 (t,  $J=7.9$  Hz, 1H), 7.64 (d,  $J=7.2$  Hz, 1H), 7.50-7.28 (m, 5H), 5.17 (dd,  $J=5.0$  and 12.5 Hz, 1H), 4.72 (s, 2H), 4.21 (s, 2H), 2.99-2.85 (m, 1H), 2.67-2.51 (m, 2H), 2.12-2.08 (m, 1H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.78, 169.78, 169.01, 168.27, 166.67, 137.04, 136.52, 135.93, 131.29, 128.36, 127.82, 124.40, 118.38, 116.08, 72.78, 69.23, 48.97, 30.92, 21.98; Anal. Calcd. For  $\text{C}_{22}\text{H}_{19}\text{N}_3\text{O}_6$ : C, 62.70; H, 4.54; N, 9.97. Found: C, 62.77; H, 4.54; N, 9.82.


3-Pentylamino-phthalic acid dimethyl ester

5



To a stirred solution of 3-amino-phthalic acid dimethyl ester (3.14 g, 15 mmol) in methylene chloride (50 ml) under a nitrogen atmosphere were added valeraldehyde (2.0 ml, 18.75 mmol) and acetic acid (5.18 ml, 90 mmol). The mixture was stirred for 5 minutes followed by addition of sodium triacetoxyborohydride (6.36g, 30 mmol). The reaction was stirred for 30 minutes, diluted with methylene chloride (50 ml), washed with water (2 x 100 ml), saturated aqueous sodium bicarbonate (2 x 100 ml), brine (100 ml), and dried (MgSO<sub>4</sub>). The solvent was evaporated *in vacuo* to give 4.19 g of product (100%) that used without further purification. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 7.31 (t, J=7.9 Hz, 1H), 6.81-6.75 (m, 2H), 3.85 (s, 3H), 3.82 (s, 3H), 3.15 (bs, 2H), 1.68-1.60 (m, 2H), 1.45-1.32 (m, 4H), 0.89 (t, J=6.9 Hz, 3H).

20

3-Pentylamino-phthalic acid

25


3-Pentylamino-phthalic acid dimethyl ester (4.19, 15 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

30

35

2-(2,6-Dioxo-piperidin-3-yl)-4-pentylamino-isoindole-1,3-dione I-51


5



To a stirred solution of 3-pentylamino-phthalic acid (2.51 g, 10 mmol) in acetic acid (50 ml) was added 3-amino-piperidine-2,6-dione hydrochloride (1.81 g, 11 mmol). The reaction mixture was heated to reflux overnight. The solvent was evaporated *in vacuo* and the residue dissolved in ethyl acetate (100 ml). The ethyl acetate mixture was washed with water (2 x 100 ml), saturated aqueous sodium bicarbonate (2 x 100 ml), brine (1 x 100 ml), and dried (MgSO<sub>4</sub>). The solvent was evaporated *in vacuo* and the solid residue was purified by chromatography (25% ethyl acetate/hexane) to give 1.82 g (53%) of product as a yellow solid: mp 141-143 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.09 (s, 1H), 7.58 (t, J=8.3 Hz, 1H), 7.08 (d, J=8.6 Hz, 1H), 7.02 (d, J=7.0 Hz, 1H), 6.519 (t, J=5.7 Hz, 1H), 5.06 (dd, J= 5.3 and 12.4 Hz, 1H), 3.32-3.25 (m, 2H), 2.97-2.82 (m, 1 H), 2.62-2.46 (m, 2H), 2.06-2.01 (m, 1H), 1.61-1.55 (m, 2H), 1.35-1.32 (m, 4H), 0.88 (t, J=6.7 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.72, 170.01, 168.92, 167.25, 146.39, 136.21, 132.14, 117.08, 110.31, 108.99, 48.52, 41.77, 30.94, 28.46, 28.33, 22.12, 21.82, 13.85; Anal. Calcd. For C<sub>18</sub>H<sub>21</sub>N<sub>3</sub>O<sub>4</sub> : C, 62.96; H, 6.16; N, 12.24. Found: C, 62.92; H, 6.17; N, 12.15.

3-Chloro-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide I-52


30



35

To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in THF (30 ml) was added 3-chlorobenzoyl chloride (0.51 ml, 4 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated  
5 *in vacuo* leaving a solid which was slurried in diethyl ether (20 ml) and filtered to give 0.82 g (100%) of product as an off-white solid: mp 257-259 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.06 (s, 1H), 10.43 (s, 1H), 8.50 (d, J=8.4 Hz, 1H), 7.99-7.88 (m, 3H), 7.75-7.61 (m, 3H), 5.17 (dd, J=5.5 and 12.7 Hz, 1H), 2.99-2.84 (m, 1H), 2.67-2.49 (m, 2H), 2.12-2.07 (m, 1H); <sup>13</sup>C  
10 NMR (DMSO-*d*<sub>6</sub>) δ 172.36, 169.78, 169.36, 167.59, 166.39, 163.57, 135.99, 135.31, 133.59, 132.11, 131.33, 130.73, 127.07, 126.76, 125.78, 118.93, 118.55, 48.91, 30.77, 21.85; Anal. Calcd. For C<sub>20</sub>H<sub>14</sub>ClN<sub>3</sub>O<sub>5</sub> : C, 58.33; H, 3.43; N, 10.20. Found: C, 58.38; H, 3.23; N, 9.95.

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-phenoxy-  
15 acetamide I-53




25 To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in THF (30 ml) was added phenoxyacetyl chloride (0.55 ml, 4 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* leaving a solid which was slurried in diethyl ether (20 ml) and filtered to give 0.76 g (93%)  
30 of product as an off-white solid: mp 236-238 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.19 (s, 1H), 10.53 (s, 1H), 8.73 (d, J=8.4 Hz, 1H), 7.88 (t, J=7.6 Hz, 1H), 7.64 (d, J=7.3 Hz, 1H), 7.38 (t, J=7.6 Hz, 2H), 7.14-7.01 (m, 3H), 5.21 (dd, J=5.3 and 12.6 Hz, 1H), 4.81 (s, 2H), 3.00-2.86 (m, 1H), 2.67-2.51 (m, 2H), 2.13-2.09 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.77, 169.77, 168.37, 167.59, 166.67, 156.85, 136.58, 135.84, 131.28, 129.74, 124.45, 121.91,  
35

118.50, 116.28, 114.86, 67.03, 49.02, 30.96, 21.93; Anal. Calcd. For  $C_{21}H_{17}N_3O_6$ : C, 61.92; H, 4.21; N, 10.31. Found: C, 61.87; H, 4.27; N, 10.25.

3-(2-Benzoyloxy-ethylamino)-phthalic acid dimethyl ester

5




10

Benzoyloxyacetaldehyde (5.27 ml, 37.5 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester. The residue (oil) was purified by chromatography (6:3:1 methylene chloride/hexane/ethyl acetate) to give 7.98 g (78%) of yellow oil:  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  7.38-7.26 (m, 5H), 6.89-6.77 (m, 3H), 4.57 (s, 2H), 3.85 (s, 3H), 3.82 (s, 3H), 3.69 (t, J=5.5 Hz, 2H), 3.38 (q, J=5.4 Hz, 2H).

15

3-(2-Benzoyloxy-ethylamino)-phthalic acid

20




25 3-(2-Benzoyloxy-ethylamino)-phthalic acid dimethyl ester (2.50 g, 7.28 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

30

35

4-(2-Benzyloxy-ethylamino)-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione I-54

5




- 10 3-(2-Benzyloxy-ethylamino)-phthalic acid (1.78 g, 5.65 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-Dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione. The solid yellow residue was recrystallized from a minimal amount of ethanol to give 1.32 g (57%) of product as a yellow solid: mp 158-160 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.11 (s, 1H), 7.57 (t, J=7.3 Hz, 1H), 7.37-7.24 (m, 5H), 7.14 (d, J=8.6 Hz, 1H), 7.04 (d, J=7.0 Hz, 1H), 6.67 (t, J=5.7 Hz, 1H), 5.07 (dd, J= 5.4 and 12.5 Hz, 1H), 4.54 (s, 2H), 3.66-3.62 (m, 2H), 3.55-3.49 (m, 2H), 2.97-2.82 (m, 1 H), 2.63-2.45 (m, 2H), 2.06-2.02 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.81, 170.08, 168.96, 167.28, 146.40, 138.23, 136.16, 132.06, 128.22, 127.42, 117.47, 110.69, 109.30, 71.90, 68.09, 48.54, 41.68, 30.97, 22.13; Anal. Calcd. For C<sub>22</sub>H<sub>21</sub>N<sub>3</sub>O<sub>5</sub>: C, 64.86; H, 5.20; N, 10.31.
- 20 Found: C, 64.95; H, 5.03; N, 10.27.

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-fluorobenzamide I-55

25


30



- To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in THF (30 ml) was added 3-Fluorobenzoyl chloride (0.49 ml, 4 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature,
- 35

methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* leaving a solid which was slurried in diethyl ether (20 ml) and filtered to give 0.69 g (96%) of product as an off-white solid: mp 260-262 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.00 (s, 1H), 10.39 (s, 1H), 8.53 (d, J=8.3 Hz, 1H), 7.94-7.50 (m, 6H), 5.15 (dd, J=5.6 and 12.7 Hz, 1H), 2.97-2.83 (m, 1H), 2.66-2.49 (m, 2H), 2.14-2.06 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.36, 169.36, 167.66, 166.39, 163.62, 136.03, 131.31, 131.12, 130.99, 126.60, 123.19, 123.15, 119.44, 119.11, 118.89, 118.42, 114.31, 113.94, 48.91, 30.77, 21.83; Anal. Calcd. For C<sub>20</sub>H<sub>14</sub>N<sub>3</sub>O<sub>5</sub>: C, 60.76; H, 3.57; N, 10.63. Found: C, 60.88; H, 3.38; N, 10.51.

10 N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methyl-benzamide I-56




20

To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in THF (30 ml) was added *m*-toluoyl chloride (0.53 ml, 4 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* leaving a solid that was slurried in diethyl ether (20 ml), filtered, reslurried in ethyl acetate (20 ml), and filtered to give 0.69 g (88%) of product as an off-white solid: mp 234-236 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.97 (s, 1H), 10.31 (s, 1H), 8.64(d, J=8.4 Hz, 1H), 7.92-7.47 (m, 6H), 5.16 (dd, J=5.5 and 12.7 Hz, 1H), 2.97-2.84 (m, 1H), 2.68-2.43 (m, 5H), 2.14-2.09 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.13, 169.15, 168.03, 166.30, 164.80, 138.18, 136.54, 135.95, 133.14, 132.87, 131.09, 128.57, 127.45, 126.64, 123.95, 118.23, 117.33, 48.88, 30.67, 21.76, 20.62; Anal. Calcd. For C<sub>21</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>: C, 64.45; H, 4.38; N, 10.74. Found: C, 64.23; H, 4.18; N, 10.56.

35

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methoxy-benzamide I-57




To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in THF (30 ml) was added *m*-anisoyl chloride (0.56 ml, 4 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* leaving a solid that was slurried in diethyl ether (20 ml), filtered, reslurried in ethyl acetate (20 ml), filtered, and recrystallized from minimal acetic acid to give 0.51 g (63%) of product as an off-white solid: mp 240-242 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.15 (s, 1H), 10.38 (s, 1H), 8.60 (d, J=8.3 Hz, 1H), 7.91 (t, J=7.5 Hz, 1H), 7.67 (d, J=7.2 Hz, 1H), 7.55-7.50 (m, 3H), 7.29-7.22 (m, 1H), 5.19 (dd, J=5.3 and 12.5 Hz, 1H), 3.86 (s, 3H), 2.99-2.84 (m, 1H), 2.66-2.50 (m, 2H), 2.12-2.07 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.02, 169.71, 168.12, 166.64, 164.70, 159.56, 136.52, 136.31, 134.73, 131.36, 130.26, 126.18, 119.19, 118.78, 118.26, 117.92, 112.74, 55.41, 48.97, 30.93, 21.99; Anal. Calcd. For C<sub>21</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>: C, 59.73; H, 4.46; N, 9.29. Found: C, 59.37; H, 4.34; N, 9.10 + 0.75 AcOH.



N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-trifluoromethyl-benzamide I-58

5

10



15


20

To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in THF (30 ml) was added 3-trifluoromethylbenzoyl chloride (0.60 ml, 4 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* leaving a solid that was slurried in diethyl ether (20 ml) and filtered to give 0.41 g (46%) of product as an off-white solid: mp 257-259 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.86 (s, 1H), 10.45 (s, 1H), 8.50 (d, J=8.5 Hz, 1H), 8.24-8.25 (m, 2H), 8.02-7.82 (m, 3H), 7.69 (d, J=7.3 Hz, 1H), 5.14 (dd, J=5.7 and 12.8 Hz, 1H), 2.95-2.82 (m, 1H), 2.67-2.48 (m, 2H), 2.14-2.07 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 171.75, 168.80, 167.31, 166.06, 163.43, 135.80, 135.60, 134.28, 131.16, 130.71, 129.75, 128.35, 128.29, 126.63, 123.59, 123.52, 118.65, 118.55, 48.84, 30.51, 21.62; Anal. Calcd. For C<sub>21</sub>H<sub>14</sub>F<sub>3</sub>N<sub>3</sub>O<sub>5</sub>: C, 56.64; H, 3.17; N, 9.44. Found: C, 56.48; H, 3.15; N, 9.41.

25

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-nitro-benzamide I-59

30



35


To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.10 g, 4 mmol) in THF (30 ml) was added 3-nitrobenzoyl chloride (1.48 g, 8 mmol). The mixture

was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* leaving a solid that was slurried in diethyl ether (30 ml) and filtered to give 1.60 g (95%) of product as an off-white solid: mp 245-247 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.85 (s, 1H), 10.53 (s, 1H), 8.76 (s, 1H), 8.50-8.37 (m, 3H), 7.95-7.87 (m, 2H), 7.70 (d, J=7.4 Hz, 1H), 5.14 (dd, J=5.7 and 12.7 Hz, 1H), 3.02-2.83 (m, 1H), 2.67-2.47 (m, 2H), 2.15-2.07 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 171.73, 168.79, 167.19, 166.04, 162.85, 147.84, 135.60, 134.79, 132.92, 131.21, 130.19, 126.87, 126.20, 121.69, 118.83, 48.86, 30.51, 21.63; Anal. Calcd. For C<sub>20</sub>H<sub>14</sub>N<sub>4</sub>O<sub>7</sub>: C, 56.88; H, 3.34; N, 13.27. Found: C, 56.87; H, 3.33; N, 13.05.

10

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-butyramide I-60

15




20

To a suspension of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.55 g, 2 mmol) in THF (30 ml) was added butanoyl chloride (0.42 ml, 4 mmol). The mixture was heated to reflux for 18 hours. The reaction was cooled to room temperature, methanol (2 ml) was added, and the mixture stirred for 1 hour. The solvent was evaporated *in vacuo* leaving a solid that was slurried in diethyl ether (20 ml) and filtered to give 0.55 g (80%) of product as an off-white solid: mp 171-173 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.14 (s, 1H), 9.68 (s, 1H), 8.48 (d, J=8.3 Hz, 1H), 7.83 (t, J=7.5 Hz, 1H), 7.61 (d, J=7.2 Hz, 1H), 5.15 (dd, J=5.2 and 12.6 Hz, 1H), 2.96-2.84 (m, 1H), 2.65-2.42 (m, 4H), 2.11-2.06 (m, 1H), 1.73-1.58 (m, 2H), 0.95 (t, J=7.4 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.70, 171.84, 169.72, 167.70, 166.63, 136.54, 136.07, 131.41, 126.17, 118.23, 116.89, 48.90, 30.91, 21.96, 18.25, 13.46; Anal. Calcd. For C<sub>17</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>: C, 59.47; H, 4.99; N, 12.24. Found: C, 59.45; H, 4.82; N, 12.15.

35

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methylamino-acetamide hydrochloride I-61


5



10

To a suspension of N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]-2-chloroacetamide (0.95 g, 2.72 mmol) in THF (30 ml) was added sodium iodide (0.41 g, 2.72 mmol) and 2M methyl amine in THF (4.08 ml, 8.15 mmol). The mixture was stirred at  
15 room temperature for 5 hours. The solvent was evaporated *in vacuo* leaving a white solid. The solid was slurried in ethyl acetate (200 ml) for 2 h. The suspension was then washed with water (3 x 100 ml), brine (100 ml), and dried (MgSO<sub>4</sub>). The solvent was evaporated *in vacuo* leaving an off-white solid. The solid was dissolved in acetonitrile (20 ml) and to this solution was added 2M HCl in ether (2 ml). The mixture was stirred for 1 h and the solvent  
20 evaporated *in vacuo*. The residue was slurried in ethyl acetate for 3 h and filtered to give an off-white solid. The solid was dissolved in water (40 ml) and washed with ethyl acetate (2 x 50 ml). The pH of the aqueous portion was adjusted to 11-12 by dropwise addition of saturated aqueous sodium carbonate. The aqueous mixture was washed with ethyl acetate (3 x 100 ml). The combined ethyl acetate extracts were washed with brine (100 ml), and  
25 dried (MgSO<sub>4</sub>). The solvent was evaporated *in vacuo* leaving a white solid. The solid was dissolved in acetonitrile (15 ml) and 2M HCl in ether (2 ml) was added to the solution. The mixture was stirred for 1 h and the solvent was evaporated *in vacuo* leaving a white solid. The solid was slurried in diethyl ether (20 ml) and filtered to give 0.18 g (17 %) of product as a white solid: mp 228-230 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.13 (s, 1H), 10.50 (s, 1H), 9.35  
30 (bs, 2H), 8.28-8.21 (m, 1H), 7.93-7.63 (m, 2H), 5.15 (dd, J=5.1 and 12.6 Hz, 1H), 4.09 (s, 2H), 3.03-2.85 (m, 1H), 2.63-2.47 (m, 5H), 2.10-2.05 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 171.99, 169.03, 166.35, 166.07, 164.75, 135.63, 134.46, 131.56, 127.45, 119.14, 118.85, 49.56, 48.82, 32.49, 30.59, 21.69; Anal. Calcd. For C<sub>16</sub>H<sub>17</sub>ClN<sub>4</sub>O<sub>5</sub>: C, 48.99; H, 4.70; N, 14.28. Found: C, 48.82; H, 4.72; N, 14.02 + 0.64 H<sub>2</sub>O.

35

2-(2,6-Dioxo-piperidin-3-yl)-4-heptylamino-isoindole-1,3-dione I-62

A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.1 g, 4.0 mmol) and heptanal (3.4 mL, 24 mmol) and acetic acid (2 mL) in DMF (20 mL) was heated until all solid was dissolved. To the mixture was added sodium borohydride (605 mg, 16 mmol) and kept at room temperature for 18h. To the mixture was added sodium

15 borohydride (150 mg, 3.9 mmol) and kept at room temperature for 1 d. The mixture was extracted with ethyl acetate (200 mL) and water (100 mL). The organic layer was washed with water (100 mL). The solvent was removed *in vacuo* to give an oil. The oil was purified by column chromatography (Silca Gel, 33% EtOAc:CH<sub>2</sub>Cl<sub>2</sub>) to give 2-(2,6-Dioxo-piperidin-3-yl)-4-heptylamino-isoindole-1,3-dione as a yellow solid (610 mg, 41% yield):

20 mp, 107-109 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.82-0.87 (m, 3H, CH<sub>3</sub>), 1.24-1.29 (m, 8H, 4 CH<sub>2</sub>), 2.00-2.04 (m, 1H, CHH), 2.43-2.62 (m, 2H, CH<sub>2</sub>), 2.82-2.96 (m, 1H, CHH), 3.23-3.31 (m, 2H, CH<sub>2</sub>), 5.06 (dd, *J* = 5.3, 12.4 Hz, 1H, NCH), 6.51 (t, *J* = 5.9 Hz, 1H, NH), 7.01 (d, *J* = 7.0 Hz, 1H, Ar), 7.07 (d, *J* = 8.6 Hz, 1H, Ar), 7.57 (dd, *J* = 7.4, 8.4 Hz, 1H, Ar), 11.10 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 13.91, 22.03, 22.17, 26.28, 28.42, 28.69, 30.98,

25 31.22, 34.84, 48.55, 109.03, 110.36, 117.13, 132.18, 136.24, 146.43, 167.29, 168.96, 170.05, 172.77; Anal Calcd for C<sub>20</sub>H<sub>25</sub>N<sub>3</sub>O<sub>4</sub>: C, 64.67; H, 6.78; N, 11.31. Found: C, 64.62; H, 6.76; N, 11.13.


30

35

4-Chloro-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide I-63

5

10



15

20


A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.2 g, 4.5 mmol) and 4-chlorobenzoyl chloride (1.1 mL, 8.8 mmol) in THF (40 mL) was heated to reflux for 15 h. To the mixture was added methanol (5 mL) to give a suspension. The suspension was filtered and washed with ether (2 x 10 mL) then methanol (5 mL) to give 4-chloro-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide as a white solid (1.5 g, 81 % yield): mp, 261-263 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.05-2.09 (m, 1H, CHH), 2.49-2.65 (m, 2H, CH<sub>2</sub>), 2.83-2.98 (m, 1H, CHH), 5.18 (dd, *J* = 5.5, 12.8 Hz, 1H, NCH), 7.68-7.72 (m, 3H, Ar), 7.89-8.01 (m, 3H, Ar), 8.52 (d, *J* = 8.2, Hz, 1H, Ar), 10.47 (s, 1H, NH), 11.16 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 21.99, 30.92, 48.98, 118.48, 119.06, 126.76, 129.10, 129.29, 131.46, 132.13, 136.28, 137.47, 164.08, 166.60, 167.86, 169.67, 172.72; Anal Calcd for C<sub>20</sub>H<sub>14</sub>N<sub>3</sub>O<sub>5</sub>Cl + 0.2 H<sub>2</sub>O: C, 57.83; H, 3.49; N, 10.12; Cl, 8.53; H<sub>2</sub>O, 0.87. Found: C, 57.88; H, 3.33; N, 9.93; Cl, 8.53; H<sub>2</sub>O, 0.73.

25

Cyclopropanecarboxylic acid [2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide I-64


30

35



A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.60 g, 2.2 mmol) and cyclopropanecarbonyl chloride (0.4 mL, 4.4 mmol) in THF (20 mL) was heated to reflux for 15 h. To the mixture was added methanol (5 mL). The solvent was removed *in vacuo* to give a solid. The solid was slurried in ether (30 mL) for 1 h. The suspension  
 5 was filtered and washed with ether (30 mL) to give cyclopropanecarboxylic acid [2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide as a solid (630 mg, 84 % yield): mp, 237-239 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.87-0.90 (m, 4H, 2CH<sub>2</sub>), 1.93-2.09 (m, 2H, CH, CHH), 2.49-2.65 (m, 2H, CH<sub>2</sub>), 2.64-2.96 (m, 1H, CHH), 5.15 (dd, *J* = 5.2, 12.6 Hz, 1H, NCH), 7.61 (d, *J* = 7.2 Hz, 1H, Ar), 7.82 (t, *J* = 7.7 Hz, 1H, Ar), 8.41 (d, *J* =  
 10 8.3 Hz, 1H, Ar), 9.99 (s, 1H, NH), 11.16 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 8.10, 14.93, 22.00, 30.93, 48.92, 117.07, 118.30, 126.69, 131.49, 135.97, 136.43, 166.67, 167.57, 169.77, 172.55, 172.74; Anal Calcd for C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub>: C, 59.82; H, 4.43; N, 12.31. Found: C, 59.50; H, 4.39; N, 12.04.


15 N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-fluoro-benzamide I-65



25 A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.1 g, 4.0 mmol) and 4-fluorobenzoyl chloride (0.95 mL, 8.0 mmol) in THF (40 mL) was heated to reflux for 15 h. To the mixture was added methanol (5 mL) to give a suspension. The suspension was filtered and washed with ether (2 x 10 mL) then methanol (5 mL) to give N-  
 30 [2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-fluoro-benzamide as a yellow solid (1.2 g, 77 % yield): mp, 283-285 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.06-2.10 (m, 1H, CHH), 2.48-2.65 (m, 2H, CH<sub>2</sub>), 2.83-2.97 (m, 1H, CHH), 5.18 (dd, *J* = 5.4, 12.6 Hz, 1H, NCH), 7.42-7.49 (m, 2H, Ar), 7.69 (d, *J* = 7.2, Hz, 1H, Ar), 7.91 (t, *J* = 8.2 Hz, 1H, Ar), 8.03-8.08 (m, 2H, Ar), 8.54 (d, *J* = 8.3 Hz, 1H, Ar), 10.41 (s, 1H, NH), 11.15 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 22.10, 30.98, 49.29, 115.96 (d, *J*<sub>C-F</sub> = 22 Hz); 118.03,  
 35 118.67, 128.09 (d, *J*<sub>C-F</sub> = 235 Hz), 130.12, 131.49, 136.18, 136.75, 162.70, 164.08, 166.56,

168.22, 169.29, 172.27; Anal Calcd for  $C_{20}H_{14}N_3O_5F + 0.2 H_2O$ : C, 60.21; H, 3.64; N, 10.53; F, 4.76;  $H_2O$ , 0.90. Found: C, 60.17; H, 3.55; N, 10.47; F, 4.90;  $H_2O$ , 0.95.

5 N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-trifluoromethyl-  
benzamide I-66




- 15 A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (0.6 g, 2.2 mmol) and 4-(trifluoromethyl)benzoyl chloride (1 g, 4.8 mmol) in THF (20 mL) was heated to reflux for 15 h. The solvent was removed *in vacuo* to give a solid. The solid was slurried in methanol (20 mL) for 2h. The suspension was filtered and washed with ether (15 mL) then methanol (15 mL) to give N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-
- 20 isoindol-4-yl]-4-trifluoromethyl-benzamide as a white solid (750 mg, 77 % yield): mp, 213-215 °C;  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  2.05-2.10 (m, 1H, CHH), 2.49-2.65 (m, 2H,  $CH_2$ ), 2.83-2.98 (m, 1H, CHH), 5.18 (dd,  $J = 5.2, 12.5$  Hz, 1H, NCH), 7.72 (d,  $J = 7.2$  Hz, 1H, Ar), 7.90-8.02 (m, 3H, Ar), 8.16-8.19 (m, 2H, Ar), 8.49 (d,  $J = 8.4$  Hz, 1H, Ar), 10.58 (s, 1H, NH), 11.17 (s, 1H, NH);  $^{13}C$  NMR (DMSO- $d_6$ )  $\delta$  21.99, 30.92, 48.99, 118.88, 119.36,
- 25 119.32, 123.76 (q,  $J_{C-F} = 271$  Hz); 125.99 (q,  $J_{C-F} = 3.6$  Hz), 127.10 128.37, 131.54, 132.16 (q,  $J_{C-F} = 32$  Hz), 136.00, 136.26, 137.21, 164.05, 166.59, 167.65, 169.68, 172.73; Anal Calcd for  $C_{21}H_{14}N_3O_5F_3$ : C, 56.64; H, 3.17; N, 9.44; F, 12.80. Found: C, 56.25; H, 3.05; N, 9.32; F, 12.69.

30


35

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-methyl-benzamide I-67



A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.1 g, 4.0 mmol) and 4-methylbenzoyl chloride (1.1 g, 8.0 mmol) in THF (40 mL) was heated to reflux for 36 h. To the mixture was added methanol (5 mL) to give a suspension. The suspension was filtered and washed with methanol (15 mL) to give N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-methyl-benzamide as a yellow solid (1.3 g, 83 % yield): mp, 322-324 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.06-2.10 (m, 1H, CHH), 2.41 (s, 3H, CH<sub>3</sub>), 2.50-2.65 (m, 2H, CH<sub>2</sub>), 2.83-2.97 (m, 1H, CHH), 5.19 (dd, *J* = 5.3, 12.6 Hz, 1H, NCH), 7.42 (d, *J* = 8.1 Hz, 2H, Ar), 7.67 (d, *J* = 7.2 Hz, 1H, Ar), 7.86-7.93 (m, 3H, Ar), 8.62 (d, *J* = 8.4 Hz, 1H, Ar), 10.37 (s, 1H, NH), 11.18 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 21.07, 21.99, 30.93, 48.99, 117.57, 118.58, 125.94, 127.28, 129.58, 130.44, 131.33, 136.34, 136.77, 142.96, 164.80, 166.65, 168.29, 169.69, 172.72; Anal Calcd for C<sub>21</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>: C, 64.45; H, 4.38; N, 10.74. Found: C, 64.65; H, 4.17; N, 10.70.


N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-nitro-benzamide I-68





A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (2.2 g, 8.0 mmol) and 4-nitrobenzoyl chloride (3.0 g, 16.0 mmol) in THF (80 mL) was heated to reflux for 15 h. To the mixture was added methanol (20 mL) to give a suspension. The suspension was filtered and washed with methanol (20 mL) to give N-[2-(2,6-Dioxo-  
 5 piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-nitro-benzamide as a white solid (2.5 g, 73 % yield): mp, 298-300 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.06-2.10 (m, 1H, CHH), 2.49-2.65 (m, 2H, CH<sub>2</sub>), 2.83-2.98 (m, 1H, CHH), 5.18 (dd, *J* = 5.2, 12.6 Hz, 1H, NCH), 7.73 (d, *J* = 7.2 Hz, 1H, Ar), 7.93 (t, *J* = 8.0 Hz, 1H, Ar), 8.19-8.22 (m, 2H, Ar), 8.42-8.47 (m, 3H, Ar), 10.65 (s, 1H, NH), 11.18 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 21.99, 30.92,  
 10 48.99, 119.30, 119.56, 124.06, 127.49, 129.01, 131.58, 135.77, 136.23, 138.96, 149.69, 163.65, 166.57, 167.48, 169.68, 172.72; Anal Calcd for C<sub>20</sub>H<sub>14</sub>N<sub>4</sub>O<sub>7</sub>: C, 56.88; H, 3.34; N, 13.27. Found: C, 57.15; H, 3.02; N, 13.22.

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-ethoxy-  
 15 acetamide I-69




To a solution of ethoxyacetic acid (0.8 mL, 8.5 mmol) and oxalyl chloride (0.7 mL, 8.0 mmol) in ether (5 mL) was added DMF (0.03 mL) at room temperature. After 3h, 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.1 g, 4.0 mmol) and THF (40 mL) was added to the mixture. Then the mixture was heated to reflux for 15 h. To the mixture was added methanol (10 mL) to give a suspension. The suspension was filtered and washed with methanol (10 mL) to give N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-  
 25 isoindol-4-yl]-2-ethoxy-acetamide as a white solid (1.3 g, 87 % yield): mp, 253-255 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.27 (t, *J* = 7.0 Hz, 3H, CH<sub>3</sub>), 2.06-2.10 (m, 1H, CHH), 2.46-2.64 (m, 2H, CH<sub>2</sub>), 2.84-2.98 (m, 1H, CHH), 3.66 (q, *J* = 7.0 Hz, 2H, CH<sub>2</sub>), 4.14 (s, 2H, CH<sub>2</sub>), 5.17 (dd, *J* = 5.2, 12.7 Hz, 1H, NCH), 7.62 (d, *J* = 7.2 Hz, 1H, Ar), 7.87 (t, *J* = 8.2 Hz, 1H, Ar), 8.75 (d, *J* = 8.4 Hz, 1H, Ar), 10.39 (s, 1H, NH), 11.16 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-  
 30 d<sub>6</sub>) δ 14.88, 21.93, 30.92, 48.98, 66.89, 69.49, 116.00, 118.28, 124.25, 131.31, 135.99,

136.53, 166.69, 168.31, 169.49, 169.73, 174.71; Anal Calcd for  $C_{17}H_{17}N_3O_6$ : C, 56.82; H, 4.77; N, 11.69. Found: C, 56.82; H, 4.71; N, 11.60.

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methoxy-

5 benzamide I-70



15 To a solution of (methylthio)acetic acid (0.77 mL, 8.9 mmol) and oxalyl chloride (0.7 mL, 8.0 mmol) in ether (5 mL) was added DMF (0.02 mL) at room temperature. After 3h, 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.1 g, 4.0 mmol) and THF (40 mL) was added to the mixture. Then the mixture was heated to reflux for 15 h. To the mixture was added methanol (10 mL) to give a suspension. The suspension was filtered and  
20 washed with methanol (10 mL) to give N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methylsulfanylacetamide as a white solid (1.0 g, 69 % yield): mp, 228-230 °C;  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  2.05-2.10 (m, 1H, CHH), 2.18 (s, 3H, CH<sub>3</sub>), 2.46-2.65 (m, 2H, CH<sub>2</sub>), 2.82-2.95 (m, 1H, CHH), 3.53 (s, 2H, CH<sub>2</sub>), 5.17 (dd,  $J$  = 5.2, 12.6 Hz, 1H, NCH), 7.63 (d,  $J$  = 7.2 Hz, 1H, Ar), 7.86 (t,  $J$  = 7.5 Hz, 1H, Ar), 8.61 (d,  $J$  = 8.4 Hz, 1H, Ar), 10.39 (s, 1H, NH), 11.16 (s, 1H, NH);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  15.62, 21.96,  
25 30.93, 37.99, 48.94, 116.69, 118.46, 125.28, 131.44, 136.31, 166.67, 167.88, 168.63, 169.78, 172.75; Anal Calcd for  $C_{16}H_{15}N_3O_5S$ : C, 53.18; H, 4.18; N, 11.63. Found: C, 53.26; H, 4.17; N, 11.52.


30

35

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methoxy-benzamide I-71

5

10



A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (660 mg, 2.4 mmol) and 2-methoxybenzoyl chloride (0.7 mL, 4.7 mmol) in THF (20 mL) was heated to reflux for 15 h. To the mixture was added methanol (5 mL) to give a suspension. The suspension was filtered and washed with methanol (20 mL) to give N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methoxy-benzamide as a white solid (760 mg, 78 % yield): mp, 286-287 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) (at 340K) δ 2.09-2.14 (m, 1H, CHH), 2.55-2.66 (m, 2H, CH<sub>2</sub>), 2.85-2.98 (m, 1H, CHH), 4.14 (s, 3H, OCH<sub>3</sub>), 5.19 (dd, *J* = 5.5, 12.9 Hz, 1H, NCH), 7.17 (t, *J* = 7.2 Hz, 1H, Ar), 7.30 (d, *J* = 8.3 Hz, 1H, Ar), 7.61-7.68 (m, 2H, Ar), 7.89 (t, *J* = 7.7 Hz, 1H, Ar), 8.12 (dd, *J* = 1.8, 7.9 Hz, 1H Ar), 9.03 (d, *J* = 8.5 Hz, 1H, Ar), 11.17 (s, 1H, NH), 11.64 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) (at 340K) δ 22.09, 31.02, 49.29, 56.19, 112.65, 116.31, 117.93, 120.66, 121.14, 125.52, 131.59, 131.75, 134.34, 136.22, 137.00, 157.64, 163.82, 166.69, 168.15, 169.43, 172.32; Anal Calcd for C<sub>21</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>: C, 61.92; H, 4.21; N, 10.31. Found: C, 62.05; H, 4.10; N, 10.38.


30

35

N-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-fluoro-benzamide I-72

5

10




A mixture of 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.1 g, 4.0 mmol) and 2-fluorobenzoyl chloride (1.0 mL, 8.4 mmol) in THF (40 mL) was heated to reflux for 15 h. To the mixture was added methanol (10 mL) to give a suspension. The suspension was filtered and washed with methanol (20 mL) to give N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-fluoro-benzamide as a white solid (1.5 g, 93 % yield): mp, 300-302 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.05-2.12 (m, 1H, CHH), 2.45-2.65 (m, 2H, CH<sub>2</sub>), 2.83-2.97 (m, 1H, CHH), 5.18 (dd, *J* = 5.5, 12.9 Hz, 1H, NCH), 7.40-7.49 (m, 2H, Ar), 7.67-7.76 (m, 2H, Ar), 7.88-7.98 (m, 1H, Ar), 8.01-8.05 (m, 1H, Ar), 8.76 (d, *J* = 8.4 Hz, 1H Ar), 10.56 (d, *J*<sub>N,F</sub> = 10 Hz, 1H, NH), 11.17 (s, 1H, NH); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 21.96, 30.92, 48.97, 116.48, 116.97 (d, *J*<sub>C,F</sub> = 14 Hz), 118.82, 120.87 (d, *J*<sub>C,F</sub> = 12 Hz), 125.32 (d, *J*<sub>C,F</sub> = 1.5 Hz), 125.74, 131.32, 131.39, 134.88 (d, *J*<sub>C,F</sub> = 9 Hz), 136.22, 136.48, 159.75 (d, *J*<sub>C,F</sub> = 252 Hz), 161.76 (d, *J*<sub>C,F</sub> = 7 Hz), 166.58, 168.04, 169.70, 172.71; Anal Calcd for C<sub>20</sub>H<sub>14</sub>N<sub>3</sub>O<sub>5</sub>F: C, 60.76; H, 3.57; N, 10.63; F, 4.81. Found: C, 60.70; H, 3.64; N, 10.64; F, 4.91.

7-Amino-N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}heptanamide Hydrochloride I-73 :


30

35



Step 1: 1,8-Diazabicyclo[5,4,0]undec-7-ene (0.7g, 4.62 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, 1-hydroxybenzotriazole (0.3g, 2.22 mmol), N-BOC-7-aminoheptanonic acid (0.54g, 2.22 mmol) and 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.53g, 2.78 mmol) were added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N citric acid (30 mL), H<sub>2</sub>O (2x30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: EtOAc 1:1) to give 7-[(tert-butoxy)carbonylamino]-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]heptanamide (0.74g, 77%) as a white solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 11.4 (s, 1H), 8.44 (t, J=5.7 Hz, 1H), 7.83-7.78 (m, 2H), 7.68-7.65 (m, 1H), 6.77 (t, J=5.1 Hz, 1H), 5.19-5.11 (dd, J=5.4 and 12.4 Hz, 1H), 4.71 (d, J=5.8 Hz, 2H), 2.93-2.84 (m, 3H), 2.63-2.49 (m, 2H), 2.21-2.05 (m, 1H), 1.55-1.49 (m, 2H), 1.36 (s, 9H), 1.36-1.20 (m, 6H).

Step 2: A 4N HCl solution in dioxane (1.5 mL) was added to a stirred solution of 7-[(tert-butoxy)carbonylamino]-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]heptanamide (0.72g, 1.40 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (25 mL) and stirred for 17 hours. The resulting suspension was filtered to give 7-amino-N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]heptanamide hydrochloride (0.26g, 41%) as a white solid: mp 187-189°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.12 (s, 1H), 8.52 (t, J=5.7 Hz, 1H), 7.93 (b, 3H), 7.88-7.67 (m, 3H), 5.18-5.11 (dd, J=5.3 and 12.4 Hz, 1H), 4.72 (d, J=5.7 Hz, 2H), 2.91-2.50 (m, 5H), 2.21 (t, J=7.2 Hz, 2H), 2.08-2.04 (m, 1H), 1.57-1.52 (m, 4H), 1.31-1.29 (m, 4H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 172.70, 172.55, 169.77, 167.44, 166.90, 139.54, 134.68, 133.08, 131.47, 127.01, 121.77, 48.82, 38.64, 37.53, 35.00, 30.90, 28.05, 26.73, 25.50, 24.89, 21.95; Anal. Calcd. For C<sub>21</sub>H<sub>27</sub>N<sub>4</sub>O<sub>5</sub>Cl + 0.64 H<sub>2</sub>O: C, 54.95; H, 6.13; N, 12.21; Cl, 7.72. Found: C, 54.56; H, 6.10; N, 11.96; Cl, 8.04.


N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}butanamide I-74

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62g, 4.08 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, butyryl chloride (0.24g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>:EtOAc 1:1) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}butanamide (0.41g, 62%) as a white solid: mp 121-123°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.14 (s, 1H), 8.44 (t, J=5.55 Hz, 1H), 7.87-7.66 (m, 3H), 5.19-5.12 (dd, J=5.1 and 12.4 Hz, 1H), 4.72 (d, J=5.6 Hz, 2H), 2.96-2.85 (m, 1H), 2.63-2.51 (m, 2H), 2.17 (t, J=7.2 Hz, 2H), 2.08-2.04 (m, 1H), 1.63-1.51 (m, 2H), 0.87 (t, J=7.3 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.71, 172.47, 169.77, 167.46, 166.92, 139.53, 134.68, 133.11, 131.49, 127.04, 121.77, 48.84, 37.55, 37.16, 30.91, 21.96, 18.60, 13.63; Anal. Calcd. For C<sub>18</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>: C, 60.50; H, 5.36; N, 11.76. Found: C, 60.46; H, 5.36; N, 11.59.

N-([2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl)benzamide I-75

5

10



15


20

25

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62g, 4.08 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 mL). After stirring for 20 min, benzoyl chloride (0.31g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 mL). The  $\text{CH}_2\text{Cl}_2$  solution was washed with  $\text{H}_2\text{O}$  (30 mL), brine (30 mL) and dried ( $\text{MgSO}_4$ ). The solvent was removed and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ : EtOAc 6:4) to give N-([2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl)benzamide (0.55g, 76%) as a white solid: mp 227-229°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.16 (s, 1H), 9.16 (t,  $J=5.7$  Hz, 1H), 7.95-7.72 (m, 5H), 7.60-7.46 (m, 3H), 5.22-5.12 (dd,  $J=5.4$  and 12.8 Hz, 1H), 4.96 (d,  $J=5.7$  Hz, 2H), 2.98-2.85 (m, 1H), 2.65-2.50 (m, 2H), 2.11-2.06 (m, 1H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.72, 169.80, 167.54, 166.96, 166.60, 139.34, 134.77, 133.92, 133.02, 131.52, 131.42, 128.34, 127.28, 127.12, 121.83, 48.88, 38.32, 30.93, 21.98; Anal. Calcd. For  $\text{C}_{21}\text{H}_{17}\text{N}_3\text{O}_5$ : C, 64.45; H, 4.38; N, 10.74. Found: C, 64.47; H, 4.50; N, 10.34.

30

35

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}phenylacetamide I-76


10

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.65g, 4.26mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.60g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, phenylacetyl chloride (0.35g, 2.22mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: EtOAc 6:4) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-2-phenylacetamide (0.41g, 55%) as a white solid: mp 128-130°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.14 (s, 1H), 8.67 (t, J=5.5 Hz, 1H), 7.80-7.61 (m, 3H), 7.29 (s, 5H), 5.19-5.12 (dd, J=5.1 and 12.4 Hz, 1H), 4.71 (d, J=5.5 Hz, 2H), 3.53 (s, 2H), 2.96-2.83 (m, 1H), 2.63-2.50 (m, 2H), 2.08-2.03 (m, 1H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.77, 170.65, 169.82, 167.45, 166.93, 139.19, 136.15, 134.68, 133.18, 131.53, 129.05, 128.25, 127.13, 126.43, 121.90, 48.85, 42.24, 37.85, 30.93, 21.98; Anal. Calcd. For C<sub>22</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>: C, 65.18; H, 4.72; N, 10.36. Found: C, 65.16; H, 4.75; N, 10.11.

30

35



N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}2pyridylcarboxamide I-77

10


1,8-Diazabicyclo[5,4,0]undec-7-ene (0.98g, 6.48mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.60g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, picolinoyl chloride hydrochloride (0.41g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCL (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 97.5:2.5) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-2-pyridylcarboxamide (0.40g, 55%) as a white solid: mp 155-157°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.15 (s, 1H), 9.50 (t, J=6.2 Hz, 1H), 8.70 (d, J=4.6 Hz, 1H), 8.08-7.98 (m, 2H), 7.82-7.62 (m, 4H), 5.21-5.14 (dd, J=5.4 and 12.6 Hz, 1H), 4.97 (d, J=6.3 Hz, 2H), 2.99-2.84 (m, 1H), 2.65-2.50 (m, 2H), 2.10-2.06 (m, 1H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.77, 169.85, 167.59, 166.99, 164.38, 149.59, 148.56, 139.02, 137.87, 134.79, 132.95, 131.57, 127.16, 126.76, 122.05, 121.87, 48.87, 38.37, 30.94, 21.97; Anal. Calcd. For C<sub>20</sub>H<sub>16</sub>N<sub>4</sub>O<sub>5</sub>+0.08H<sub>2</sub>O: C, 61.08; H, 4.13; N, 14.25. Found: C, 61.48; H, 4.22; N, 13.87.

30

35

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}undecamide I-78

5




10

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62g, 4.08 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.60 g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 mL). After stirring for 20 min, undecanoyl chloride (0.45g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 mL). The  $\text{CH}_2\text{Cl}_2$  solution was washed with 1N HCl (30 mL),  $\text{H}_2\text{O}$  (30 mL), brine (30 mL) and dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ : EtOAc 6:4) to give N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}undecanamide (0.53g, 63%) as a white solid: mp 138-139°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.12 (s, 1H), 8.42 (t,  $J=5.9$  Hz, 1H), 7.85-7.78 (m, 2H), 7.71-7.65 (m, 1H), 5.18-5.11 (dd,  $J=5.4$  and 12.5 Hz, 1H), 4.72 (d,  $J=5.9$  Hz, 2H), 2.96-2.83 (m, 1H), 2.64-2.47 (m, 2H), 2.18 (t,  $J=7.3$  Hz, 2H), 2.08-2.04 (m, 1H), 1.55-1.50 (m, 2H), 1.24 (s, 14H), 0.85 (t,  $J=6.1$  Hz, 3H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.74, 172.63, 169.80, 167.47, 166.93, 139.55, 134.65, 133.10, 131.50, 127.04, 121.79, 48.83, 37.55, 35.21, 31.28, 30.92, 28.94, 28.74, 28.67, 25.20, 22.08, 21.97, 13.94; Anal. Calcd. For  $\text{C}_{25}\text{H}_{33}\text{N}_3\text{O}_5$ : C, 65.91; H, 7.30; N, 9.23. Found: C, 66.08; H, 7.13; N, 9.23.

30

35

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}2methylpropanamide I-79

10

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62g, 4.08 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.60g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, isobutyryl chloride (0.24g, 2.22 mmol) was added. The mixture was stirred at room temperature for 15 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the solid was purified from ether (10 mL) and hexane (10 mL) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-2-methylpropanamide (0.48g, 73%) as a white solid: mp 218-20 220°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.13 (s, 1H), 8.39 (t, J=5.8 Hz, 1H), 7.87-7.78 (m, 2H), 7.66-7.63 (m, 1H), 5.19-5.12 (dd, J=6.9 and 12.5 Hz, 1H), 4.71 (d, J=5.8 Hz, 2H), 2.97-2.83 (m, 1H), 2.63-2.43 (m, 3H), 2.08-2.04 (m, 1H), 1.07 (d, J=6.9 Hz, 6H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 176.52, 172.75, 169.82, 167.49, 166.95, 139.54, 134.77, 132.85, 131.50, 127.02, 121.77, 48.83, 37.48, 33.98, 30.92, 21.96, 19.53; Anal. Calcd. For C<sub>18</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>: C, 60.50; H, 25 5.36; N, 11.76. Found: C, 60.48; H, 5.33; N, 11.64.

30


35

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}cyclopentylcarboxamide

**I-80**

5

10



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62g, 4.08 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.60g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, cyclopentanecarbonyl chloride (0.29g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the solid was stirred with ether (20 mL) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}cyclopentylcarboxamide (0.59g, 83%) as a white solid: mp 175-177°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.13 (s, 1H), 8.41 (t, J=5.7 Hz, 1H), 7.87-7.78 (m, 2H), 7.66-7.63 (m, 1H), 5.19-5.12 (dd, J=5.3 and 12.5 Hz, 1H), 4.72 (d, J=5.8 Hz, 2H), 2.98-2.83 (m, 1H), 2.73-2.51 (m, 3H), 2.08-2.04 (m, 1H), 1.81-1.51 (m, 8H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 175.71, 172.75, 169.82, 167.49, 166.95, 139.68, 134.76, 132.91, 131.50, 127.01, 121.77, 48.84, 44.20, 37.60, 30.93, 29.96, 25.60, 21.97; Anal. Calcd. For C<sub>20</sub>H<sub>21</sub>N<sub>3</sub>O<sub>5</sub>: C, 62.65; H, 5.52; N, 10.96. Found: C, 62.52; H, 5.55; N, 10.81.


30

35

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}cyclohexylcarboxamide  
I-81

5

10




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.62g, 4.08 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.60g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 mL). After stirring for 20 min, cyclohexanecarbonyl chloride (0.33g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 mL). The  $\text{CH}_2\text{Cl}_2$  solution was washed with 1N HCL (30 mL),  $\text{H}_2\text{O}$  (30 mL), brine (30 mL) and dried ( $\text{MgSO}_4$ ). The solvent was removed and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ : EtOAc 6:4) to give N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}cyclohexylcarboxamide (0.53g, 72%) as a white solid: mp 142-144°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.13 (s, 1H), 8.36 (t,  $J=5.8$  Hz, 1H), 7.86-7.77 (m, 2H), 7.64-7.61 (m, 1H), 5.18-5.11 (dd,  $J=5.3$  and 12.5 Hz, 1H), 4.70 (d,  $J=5.8$  Hz, 2H), 2.97-2.83 (m, 1H), 2.63-2.47 (m, 2H), 2.26-2.17 (m, 1H), 2.08-2.03 (m, 1H), 1.79-1.61 (m, 5H), 1.43-1.12 (m, 5H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  175.58, 172.75, 169.82, 167.49, 166.96, 139.68, 134.75, 132.76, 131.49, 126.99, 121.73, 48.83, 43.90, 37.43, 30.92, 29.20, 25.43, 25.24, 21.96; Anal. Calcd. For  $\text{C}_{21}\text{H}_{23}\text{N}_3\text{O}_5$ : C, 63.47; H, 5.83; N, 10.57. Found: C, 63.12; H, 5.68; N, 10.41.

30


35

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-(phenylamino)-carboxamide I-82




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, phenyl isocyanate (0.33g, 2.77 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub> : EtOAc 7:3) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-(phenylamino)carboxamide (0.23g, 31%) as a white solid: mp 212-214°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.15 (s, 1H), 8.78 (s, 1H), 7.88-7.76 (m, 3H), 7.37 (d, J=7.7 Hz, 2H), 7.21 (t, J=7.7 Hz, 2H), 6.89 (t, J=7.3 Hz, 1H), 6.76 (t, J=5.9 Hz, 1H), 5.20-5.13 (dd, J=5.3 and 12.5 Hz, 1H), 4.72 (d, J=5.9 Hz, 2H), 2.97-2.84 (m, 1H), 2.65-2.49 (m, 2H), 2.09-2.05 (m, 1H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.72, 169.79, 167.58, 166.99, 155.22, 140.25, 134.69, 133.63, 131.59, 128.60, 127.18, 121.83, 121.18, 117.70, 48.86, 38.71, 30.92, 21.97; Anal. Calcd. For C<sub>21</sub>H<sub>18</sub>N<sub>4</sub>O<sub>5</sub>: C, 62.07; H, 4.46; N, 13.79. Found: C, 62.14; H, 4.49; N, 13.49.

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(butylamino)carboxamide  
I-83



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, n-butyl isocyanate (0.27g, 2.77 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: EtOAc 1:1) to give N-{[2-(2,6-dioxo(3-piperidyl))1,3-dioxoisindolin-4-yl]methyl}(butylamino)carboxamide (0.44g, 61%) as a white solid: mp 172-174°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.13 (s, 1H), 7.86-7.68 (m, 3H), 6.42 (t, J=5.9 Hz, 1H), 6.12 (t, J=5.4 Hz, 1H), 5.18-5.11 (dd, J=5.2 and 12.4 Hz, 1H), 4.63 (d, J=5.9 Hz, 1H), 3.03-2.83 (m, 3H), 2.64-2.51 (m, 2H), 2.08-2.04 (m, 1H), 1.37-1.22 (m, 4H), 0.86 (t, J=7.0 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.70, 169.77, 167.54, 167.00, 157.98, 141.14, 134.56, 133.33, 131.47, 126.94, 121.58, 48.80, 38.98, 38.70, 32.01, 30.90, 21.95, 19.46, 13.64; Anal. Calcd. For C<sub>19</sub>H<sub>22</sub>N<sub>4</sub>O<sub>5</sub>: C, 59.06; H, 5.74; N, 14.50. Found: C, 59.24; H, 5.53; N, 14.37.

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl}(propylamino)-carboxamide I-84




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 mL). After stirring for 20 min, propyl isocyanate (0.24 g, 2.77 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 mL). The  $\text{CH}_2\text{Cl}_2$  solution was washed with 1N HCl (30 mL),  $\text{H}_2\text{O}$  (30 mL), brine (30 mL) and dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ :  $\text{CH}_3\text{OH}$  100: 3) to give N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl}(propylamino)carboxamide (0.13g, 20%) as a white solid: mp 160-162°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.14 (s, 1H), 7.86-7.69 (m, 3H), 6.44 (t,  $J=5.9$  Hz, 1H), 6.16 (t,  $J=1$  Hz), 5.18-5.11 (dd,  $J=5.3$  and 12.4 Hz, 1H), 4.63 (d,  $J=5.9$  Hz, 2H), 2.99-2.83 (m, 3H), 2.64-2.50 (m, 2H), 2.08-2.04 (m, 1H), 1.42-1.32 (m, 2H), 0.83 (t,  $J=7.3$  Hz, 3H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.78, 169.84, 167.59, 167.05, 158.03, 141.16, 134.62, 133.34, 131.51, 126.96, 121.63, 48.82, 41.18, 30.94, 23.15, 21.99, 11.33; Anal. Calcd. For  $\text{C}_{18}\text{H}_{20}\text{N}_4\text{O}_5$ : C, 58.06; H, 5.41; N, 15.05. Found: C, 57.94; H, 5.31; N, 14.90.



N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(cyclohexylamino)-carboxamide I-85

5

10



15

20


25

30

35


1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in  $\text{CH}_3\text{CN}$  (50 mL). After stirring for 20 min, cyclohexyl isocyanate (0.35g, 2.77 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 mL). The  $\text{CH}_2\text{Cl}_2$  solution was washed with 1N HCl (30 mL),  $\text{H}_2\text{O}$  (30 mL), brine (30 mL) and dried ( $\text{MgSO}_4$ ). The solvent was removed *in vacuo* and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ : EtOAc 1:1) to give N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(cyclohexylamino)carboxamide (0.37g, 49%) as a white solid: mp 208-210°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.13 (s, 1H), 7.86-7.68 (m, 3H), 6.34 (t,  $J=5.8\text{Hz}$ , 1H), 6.04 (d,  $J=7.9\text{Hz}$ , 1H), 5.18-5.11 (dd,  $J=5.3$  and  $12.4\text{Hz}$ , 1H), 4.62 (d,  $J=5.8\text{Hz}$ , 2H), 3.37 (m, 1H), 2.96-2.83 (m, 1H), 2.63-2.50 (m, 2H), 2.08-2.04 (m, 1H), 1.76-1.02 (m, 10H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.78, 169.84, 167.59, 167.04, 157.28, 141.13, 134.64, 133.42, 131.51, 126.98, 121.64, 48.82, 47.91, 38.66, 33.23, 30.94, 25.27, 24.47, 21.99; Anal. Calcd. For  $\text{C}_{21}\text{H}_{24}\text{N}_4\text{O}_5$ : C, 61.16; H, 5.87; N, 13.58. Found: C, 61.21; H, 5.79; N, 13.63.

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}{(methylethylamino)}]-  
carboxamide I-86



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, isopropyl isocyanate (0.24g, 2.77 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 97.5:2.5) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}{(methylethylamino)]carboxamide (0.25g, 36%) as a white solid: mp 180-182 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.19 (s, 1H), 7.87-7.68 (m, 3H), 6.33 (t, J=5.9 Hz, 1H), 6.02 (d, J=7.5 Hz, 1H), 5.18-5.11 (dd, J=5.2 and 12.4 Hz, 1H), 4.62 (d, J=5.9 Hz, 2H), 3.73-3.35 (m, 1H), 2.98-2.83 (m, 1H), 2.63-2.50 (m, 2H), 2.08-2.04 (m, 1H), 1.04 (d, J=6.5 Hz, 6H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.78, 169.85, 167.59, 167.05, 157.36, 141.16, 134.65, 133.39, 131.52, 126.98, 121.64, 48.82, 41.03, 38.64, 30.94, 23.18, 21.99; Anal. Calcd. For C<sub>18</sub>H<sub>20</sub>N<sub>4</sub>O<sub>5</sub>: C, 58.06; H, 5.41; N, 15.05. Found: C, 58.20; H, 5.44; N, 14.95.

N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(octylamino)carboxamide  
I-87




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, octyl isocyanate (0.44g, 2.77 mmol) was added. The mixture was stirred at room temperature for 17 hours. The slurry mixture was filtered and the solid was recrystallized from methanol to give N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(octylamino)carboxamide (0.46g, 56%) as a white solid: mp 160-162°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.14 (s, 1H), 7.85-7.68 (m, 3H), 6.43 (t, J=6.0 Hz, 1H), 6.13 (t, J=5.6 Hz, 1H), 5.18-5.11 (dd, J=5.3 and 12.5 Hz, 1H), 4.62 (d, J=6.0 Hz, 2H), 3.02-2.83 (m, 3H), 2.64-2.50 (m, 2H), 2.08-2.04 (m, 1H), 1.36-1.24 (m 12H), 0.85 (t, J=6.2 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.75, 169.82, 167.57, 167.03, 157.99, 141.19, 134.57, 133.33, 131.50, 126.95, 121.61, 48.82, 39.32, 38.83, 31.21, 30.93, 29.92, 28.73, 28.69, 26.37, 22.07, 21.98, 13.93; Anal. Calcd. For C<sub>23</sub>H<sub>30</sub>N<sub>4</sub>O<sub>5</sub>: C, 62.43; H, 6.83; N, 12.66. Found: C, 62.27; H, 6.94; N, 12.54.

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(benzylamino)-  
carboxamide I-88

5

10




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, benzyl isocyanate (0.32g, 2.41 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (70 mL). The CH<sub>2</sub>Cl<sub>2</sub> solution was washed with 1N HCl (30 mL), H<sub>2</sub>O (30 mL), brine (30 mL) and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue was purified by chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 96:4) to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(benzylamino)carboxamide (0.42g, 54%) as a white solid: mp 192-194°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.13 (s, 1H), 7.86-7.69 (m, 3H), 7.34-7.19 (m, 5H), 6.67 (t, J=5.8 Hz, 1H), 6.60 (t, J=5.9 Hz, 1H), 5.18-5.11 (dd, J= 5.3 and 12.5 Hz, 1H), 4.67 (d, J=5.9 Hz, 2H), 4.23 (d, J=5.8 Hz, 2H), 2.97-2.83 (m, 1H), 2.63-2.50 (m, 2H), 2.07-2.03 (m, 1H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 175.63, 172.75, 167.56, 167.03, 158.05, 141.01, 140.70, 134.61, 133.31, 131.52, 128.19, 126.98, 126.55, 121.66, 48.83, 42.99, 30.93, 21.98; Anal. Calcd. For C<sub>22</sub>H<sub>20</sub>N<sub>4</sub>O<sub>5</sub>: C, 62.85; H, 4.79; N, 13.33. Found: C, 62.78; H, 4.53; N, 13.18.

30

35

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(cyclopropylamino)-carboxamide I-89

5




10 1,8-Diazabicyclo[5,4,0]undec-7-ene (0.58g, 3.81mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, 4-nitrophenyl-N-cyclopropylcarbamate (0.41g, 1.85 mmol) was added. The mixture was stirred at room temperature for 17 hours. The mixture was filtered and the solid was

15 recrystallized from methanol to give N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(cyclopropylamino)carboxamide (0.53g, 77%) as a white solid: mp 245-247°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.14 (s, 1H), 7.87-7.69 (m, 3H), 6.58 (t, J=5.7 Hz, 1H), 6.45 (d, J=2.2 Hz, 1H) 5.19-5.11 (dd, J=5.4 and 12.5 Hz, 1H), 4.65 (d, J=6.1 Hz, 2H), 2.96-2.83 (m, 1H), 2.64-2.40 (m, 3H), 2.08-2.04 (m, 1H), 0.62-0.55 (m, 2H), 0.40-0.34 (m, 2H); <sup>13</sup>C NMR

20 (DMSO-d<sub>6</sub>) δ 172.75, 169.82, 167.63, 167.04, 158.69, 141.11, 134.61, 133.26, 131.49, 126.94, 121.60, 48.83, 30.93, 22.37, 21.97, 6.59; Anal. Calcd. For C<sub>18</sub>H<sub>18</sub>N<sub>4</sub>O<sub>5</sub>: C, 58.37; H, 4.90; N, 15.13. Found: C, 58.26; H, 4.82; N, 14.85.

25 2-(2,6-Dioxo(3-piperidyl))-4-({[(ethylamino)thioxomethyl]amino}methyl)isindoline-1,3-dione I-114


30



35 1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29g, 1.90 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6g, 1.85 mmol) in CH<sub>3</sub>CN (50 mL). After stirring for 20 min, ethyl

idothiocyanate (0.2g, 2.22 mmol) was added. The mixture was stirred at room temperature for 17 hours. The solvent was removed *in vacuo* and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (70 mL). The  $\text{CH}_2\text{Cl}_2$  solution was washed with 2N HCl (30 mL),  $\text{H}_2\text{O}$  (30 mL), brine (30 mL) and dried ( $\text{MgSO}_4$ ). The solvent was removed and the residue was purified by chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ :EtOAc 6:4) to give 2-(2,6-dioxo(3-piperidyl))-4-  
 5 {{{(ethylamino)thioxomethyl}amino}methyl}isoindoline-1,3-dione (0.33g, 48%) as a white solid: mp 154-156°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  11.14 (s, 1H), 7.86-7.66 (m, 5H), 5.19-5.09 (m, 3H), 3.38 (m, 2H), 2.98-2.83 (m, 1H), 2.64-2.50 (m, 2H), 2.08-2.04 (m, 1H), 1.09 (t,  $J=7.1$  Hz, 3H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  172.78, 169.85, 167.56, 167.02, 139.64, 134.53,  
 10 133.22, 131.54, 127.06, 121.76, 48.85, 42.74, 30.84, 22.80, 21.99, 14.32; Anal. Calcd. For  $\text{C}_{17}\text{H}_{18}\text{N}_4\text{O}_4\text{S}$ : C, 54.53; H, 4.85; N, 14.96; S, 8.56. Found: C, 54.89; H, 4.82; N, 14.72; S, 8.51.

{5-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]carbamoyl}-  
 15 pentyl}-carbamic acid benzyl ester I-117




A solution of 6-benzyloxycarbonylamino-hexanoic acid (2.65 g, 10mmol) in thionyl chloride (15 ml) was heated to reflux 1h. The reaction mixture was allowed to cool to room temperature and the solvent was evaporated *in vacuo* to give (5-chlorocarbonyl-pentyl)-  
 25 carbamic acid benzyl ester as tan oil. The oil was used without further purification. (5-Chlorocarbonyl-pentyl)-carbamic acid benzyl ester was dissolved in THF (50 ml). To this solution was added 4-amino-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione (1.37 g, 5 mmol). The resulting suspension was to reflux for 4 hours. The solvent was evaporated *in vacuo* and the resulting solid was partially purified by flash chromatography (60/40 ethyl  
 30 acetate/hexane) to give a light yellow solid. The solid was recrystallized from minimal ethyl acetate to give 1.24 g (48%) of product as an off-white solid: mp 122-125°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ) 11.18 (s, 1H), 9.70 (s, 1H), 8.49 (d,  $J=8.4$  Hz, 1H), 7.83 (t,  $J=7.7$  Hz, 1H), 7.61 (d,  $J=7.2$  Hz, 1H), 7.34-7.24 (m, 6H), 5.16 (dd,  $J=5.1$  and 12.6 Hz, 1H), 5.00 (s, 2H), 3.02-2.84 (m, 3H), 2.65-2.43 (m, 4H), 2.12-2.05 (m, 1H), 1.65-1.15 (m, 6H;  $^{13}\text{C}$  NMR ( $\text{DMSO}-$   
 35  $d_6$ ) 172.75, 171.95, 169.77, 167.72, 166.67, 156.08, 137.29, 136.57, 136.09, 131.43,

128.32, 127.70, 126.21, 118.25, 116.90, 65.08, 48.92, 36.47, 30.94, 30.67, 29.14, 25.77, 24.47, 21.99; Anal. Calcd. For  $C_{27}H_{28}N_4O_7$ : C, 62.30; H, 5.42; N, 10.76. Found: C, 62.40; H, 5.31; N, 10.48.

5 2-Methoxy-N-[2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-acetamide I-118

10




To a suspension of 4-amino-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione (0.10 g, 0.35 mmol) in THF (15 ml) was added methoxyacetyl chloride (0.08 g, 0.70 mmol).

- 15 The mixture was heated to reflux for 6 hours. The reaction mixture was allowed to cool and to the solution was added methanol (2 ml). The reaction mixture was stirred for an additional 15 minutes followed by solvent evaporation *in vacuo*. The resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.11 g (87 %) of product as an off-white solid: mp 244-246°C;  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ) 11.07 (s, 1H), 10.26 (s, 1H), 8.68 (d,  $J=8.3$  Hz, 1H), 7.82 (t,  $J=7.7$  Hz, 1H), 7.55 (d,  $J=7.2$  Hz, 1H), 4.10 (s, 2H), 3.47 (s, 3H), 2.77-2.51 (m, 3H), 2.09-2.04 (m, 1H), 1.90 (s, 3H);  $^{13}\text{C}$  NMR ( $\text{DMSO}-d_6$ ) 172.12, 171.99, 169.15, 168.99, 167.38, 136.22, 135.68, 131.06, 124.25, 117.90, 115.91, 71.43, 59.06, 58.88, 29.01, 28.53, 20.92; Anal. Calcd. For  $C_{17}H_{17}N_3O_6$ : C, 56.48; H, 4.81; N, 11.62. Found: C, 56.10; H, 4.55; N, 11.39 + 0.12  $\text{H}_2\text{O}$ .

25

Pentanoic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide I-119


30



35

To a suspension of 4-amino-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione (0.10 g, 0.35 mmol) in THF (15 ml) was added valeryl chloride (0.08 g, 0.70 mmol). The mixture was heated to reflux for 18 h. The reaction mixture was allowed to cool and to the solution was added methanol (2 ml). The reaction mixture was stirred for an additional 15 minutes followed by solvent evaporation *in vacuo*. The resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.11 g (81 %) of product as an off-white solid: mp 190-192°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) 11.04 (s, 1H), 9.65 (s, 1H), 8.50 (d, J=8.4 Hz, 1H), 7.79 (t, J=7.8 Hz, 1H), 7.52 (d, J=7.2 Hz, 1H), 2.74-2.44 (m, 5H), 2.10-2.03 (m, 1H), 1.90 (s, 3H), 1.66-1.54 (m, 2H), 1.42-1.27 (m, 2H), 0.91 (t, J=7.3 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) 172.13, 172.03, 168.75, 167.36, 136.42, 135.91, 131.20, 125.68, 117.71, 116.29, 58.81, 36.25, 29.07, 28.54, 26.83, 21.65, 20.98, 13.64; Anal. Calcd. For C<sub>19</sub>H<sub>21</sub>N<sub>3</sub>O<sub>5</sub>: C, 61.39; H, 5.70; N, 11.30. Found: C, 61.03; H, 5.51; N, 11.13 + 0.02 H<sub>2</sub>O.


Heptanoic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide I-120



To a suspension of 4-amino-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione (0.10 g, 0.35 mmol) in THF (15 ml) was added heptanoyl chloride (0.10 g, 0.70 mmol). The mixture was heated to reflux for 18 h. The reaction mixture was allowed to cool and to the solution was added methanol (2 ml). The reaction mixture was stirred for an additional 15 minutes followed by solvent evaporation *in vacuo*. The resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.10 g (74 %) of product as an off-white solid: mp 172-174°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) 11.05 (s, 1H), 9.65 (s, 1H), 8.50 (d, J=8.3 Hz, 1H), 7.79 (t, J=7.6 Hz, 1H), 7.52 (d, J=7.2 Hz, 1H), 2.77-2.43 (m, 5H), 2.08-2.03 (m, 1H), 1.90 (s, 3H), 1.61 (t, J=6.4 Hz, 2H), 1.29 (bs, 6H), 0.87 (t, J=6.3 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) 172.12, 172.02, 168.74, 167.36, 136.41, 135.90, 131.20, 125.70, 117.71, 116.31, 58.81, 36.51, 30.93, 29.07, 28.54, 28.12, 24.69, 21.94, 20.98, 13.86; Anal. Calcd. For C<sub>21</sub>H<sub>25</sub>N<sub>3</sub>O<sub>5</sub>: C, 62.83; H, 6.33; N, 10.47. Found: C, 62.45; H, 6.18; N, 10.24 + 0.50 H<sub>2</sub>O.




3-Chloro-N-[2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide I-121



To a suspension of 4-amino-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione (0.10 g, 0.35 mmol) in THF (15 ml) was added 3-chlorobenzoyl chloride (0.12 g, 0.70 mmol). The mixture was heated to reflux for 18 h. The reaction mixture was allowed to cool and to the solution was added methanol (2 ml). The reaction mixture was stirred for an additional 15 minutes followed by solvent evaporation *in vacuo*. The resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.11 g (73 %) of product as an off-white solid: mp 283-285 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) 11.06 (s, 1H), 10.43 (s, 1H), 8.48 (d, J=8.3 Hz, 1H), 7.98-7.61 (m, 6H), 2.80-2.51 (m, 3H), 2.11-2.01 (m, 1H), 1.92 (s, 3H), 1.61 (t, J=6.4 Hz, 2H), 1.29 (bs, 6H), 0.87 (t, J=6.3 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) 171.72, 171.64, 168.53, 167.11, 163.55, 135.81, 135.75, 135.35, 133.59, 132.07, 131.03, 130.73, 127.03, 126.37, 125.58, 118.43, 118.06, 58.80, 28.90, 28.37, 20.84; Anal. Calcd. For C<sub>21</sub>H<sub>16</sub>ClN<sub>3</sub>O<sub>5</sub>: C, 59.23; H, 3.79; N, 9.87. Found: C, 59.00; H, 3.80; N, 9.70.


N-[2-(3-Methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-propionamide I-122



To a suspension of 4-amino-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione (0.10 g, 0.35 mmol) in THF (15 ml) was added propionyl chloride (0.07 g, 0.70 mmol).

The mixture was heated to reflux for 18 h. The reaction mixture was allowed to cool and to the solution was added methanol (2 ml). The reaction mixture was stirred for an additional 15 minutes followed by solvent evaporation *in vacuo*. The resulting solid was slurried in diethyl ether (20 ml) and filtered to give 0.07 g (63 %) of product as an off-white solid: mp 222-224°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) 11.03 (s, 1H), 9.63 (s, 1H), 8.51 (d, J=8.3 Hz, 1H), 7.79 (t, J=7.6 Hz, 1H), 7.51 (d, J=7.3 Hz, 1H), 2.76-2.45 (m, 5H), 2.10-2.03 (m, 1H), 1.90 (s, 3H), 1.12 (t, J=7.5 Hz, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) 172.70, 172.14, 172.03, 168.80, 167.37, 136.48, 135.94, 131.20, 125.59, 117.67, 116.21, 58.81, 29.71, 29.07, 28.53, 20.97, 14.74, 9.18; Anal. Calcd. For C<sub>17</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>: C, 59.47; H, 4.99; N, 12.24 Found: C, 59.24; H, 5.06; N, 11.97.


Thiophene-2-carboxylic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide I-123



To a suspension of 4-amino-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione (0.10 g, 0.35 mmol) in THF (15 ml) was added 2-thiophenecarbonyl chloride (0.10 g, 0.70 mmol). The mixture was heated to reflux for 18 h. The reaction mixture was allowed to cool and to the solution was added methanol (2 ml). The reaction mixture was stirred for an additional 15 minutes followed by solvent evaporation *in vacuo*. The resulting solid was slurried in ethyl acetate (10 ml) and filtered to give 0.13 g of light yellow solid. The solid was recrystallized from acetonitrile to give 0.10 g of product (72%) as a off-white solid: mp 288-290°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) 11.07 (s, 1H), 10.32 (s, 1H), 8.49 (d, J=8.4 Hz, 1H), 7.98 (d, J=4.8 Hz, 1H), 7.89-7.83 (m, 2H), 7.59 (d, J=7.0 Hz, 1H), 7.31 (t, J=4.1 Hz, 1H), 2.80-2.51 (m, 3H), 2.12-2.01 (m, 1H), 1.92 (s, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) 172.10, 171.99, 169.00, 167.33, 159.60, 137.99, 136.05, 136.03, 133.26, 131.13, 129.82, 128.60, 125.91, 118.29, 117.43, 58.93, 29.04, 28.55, 20.96; Anal. Calcd. For C<sub>19</sub>H<sub>15</sub>N<sub>3</sub>O<sub>5</sub>S: C, 57.17; H, 3.84; N, 10.53 Found: C, 56.91; H, 3.48; N, 10.55 + 0.1 H<sub>2</sub>O.

3-[(5-Methyl-furan-2-ylmethyl)-amino]-phthalic acid dimethyl ester


5



3-Amino-phthalic acid dimethyl ester (1.05 g, 5 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester. The product of the reaction was used without further purification.

3-[(5-Methyl-furan-2-ylmethyl)-amino]-phthalic acid


15



3-[(5-Methyl-furan-2-ylmethyl)-amino]-phthalic acid dimethyl ester (5 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

2-(2,6-Dioxo-piperidin-3-yl)-4-[(5-methyl-furan-2-ylmethyl)-amino]-isoindole-1,3-dione25 **I-124**


30



3-[(5-Methyl-furan-2-ylmethyl)-amino]-phthalic acid (5 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione. The residue was purified by preparative HPLC

(Symmetry C<sub>18</sub>, isocratic, 35/65 acetonitrile/water) to give 0.31 g (20%) of product as a yellow solid: mp 305-307°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.10 (s, 1H), 7.58 (t, J=7.7 Hz, 1H), 7.18 (d, J=8.6 Hz, 1H), 7.06 (d, J=7.1 Hz, 1H), 6.93 (t, J=6.0 Hz, 1H), 6.22 (d, J=2.9 Hz, 1H), 5.98 (d, J=1.9 Hz, 1H), 5.07 (dd, J= 5.3 and 12.5 Hz, 1H), 4.49 (d, J=6.0 Hz, 2H), 2.97-2.82 (m, 1 H), 2.62-2.44 (m, 2H), 2.22 (s, 3H), 2.06-1.99 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.77, 170.02, 168.77, 167.21, 151.04, 150.02, 145.80, 136.04, 132.08, 117.57, 110.96, 109.63, 108.27, 106.34, 48.55, 38.94, 30.95, 22.10, 13.24; Anal. Calcd. For C<sub>19</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>: C, 62.12; H, 4.66; N, 11.44. Found: C, 61.75; H, 4.71; N, 11.15.


10 3-[(5-Hydroxymethyl-furan-2-ylmethyl)-amino]-phthalic acid dimethyl ester



5-Hydroxymethyl-furan-2-carbaldehyde (0.95 ml, 7.5 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester.

20 The residue (oil) was purified by chromatography (SiO<sub>2</sub>, 40/60 ethyl acetate/hexanes) to give 0.97 g (76%) of yellow oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.35-7.27 (m, 1H), 7.05 (bs, 1H), 6.87-6.83 (m, 2H), 6.21 (d, J=3.1 Hz, 1H), 6.17 (d, J=3.2 Hz, 1H), 4.57 (s, 2H), 4.38 (bs, 2H) 3.86 (s, 3H), 3.83 (s, 3H), 1.98 (bs, 1H).

25 3-[(5-Hydroxymethyl-furan-2-ylmethyl)-amino]-phthalic acid




3-[(5-Hydroxymethyl-furan-2-ylmethyl)-amino]-phthalic acid dimethyl ester (0.97 g, 3.04 mmol) was treated in the same manner as described above for the synthesis of 3-(2-

35


methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

5 2-(2,6-Dioxo-piperidin-3-yl)-4-[(5-hydroxymethyl-furan-2-ylmethyl)-amino]-isoindole-1,3-  
dione I-125



10 3-[(5-Hydroxymethyl-furan-2-ylmethyl)-amino]-phthalic acid (3.04 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-Dioxo-piperidin-3-yl)-4-(2-  
 15 methoxy-ethylamino)-isoindole-1,3-dione. The solid yellow residue was purified by chromatography (SiO<sub>2</sub>, 60% ethyl acetate/hexanes) to give 0.40 g of solid which was dissolved in methylene chloride (70 ml) and washed with saturated NaHCO<sub>3</sub> (2 x 100 ml), brine (1 x 100 ml), dried (magnesium sulfate), and filtered. The solvent was evaporated leaving 0.32 g (27%) of product as a yellow solid: mp 172-174°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  
 20 δ 11.11 (s, 1H), 7.58 (t, J=7.9 Hz, 1H), 7.18 (d, J=8.6 Hz, 1H), 7.06 (d, J=7.0 Hz, 1H), 6.99 (t, J=6.0 Hz, 1H), 6.27 (d, J=2.9 Hz, 1H), 6.20 (d, J=2.9 Hz, 1H), 5.18 (t, J=5.7 Hz, 1H), 5.07 (dd, J= 5.2 and 12.5 Hz, 1H), 4.54 (d, J=5.9 Hz, 2H), 4.33 (d, J=5.7 Hz, 2H), 2.96-2.82 (m, 1 H), 2.63-2.46 (m, 2H), 2.06-1.99 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.80, 170.07, 168.80, 167.24, 155.00, 151.10, 145.79, 136.08, 132.12, 117.57, 111.04, 109.71, 108.06,  
 25 107.60, 55.63, 48.60, 39.05, 30.98, 22.13; Anal. Calcd. For C<sub>19</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>: C, 59.53; H, 4.47; N, 10.96. Found: C, 59.30; H, 4.54; N, 10.70.


30 3-[(Thiophen-2-ylmethyl)-amino]-phthalic acid dimethyl ester



35


2-Thiophenecarboxaldehyde (1.12 g, 10 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester. The residue (oil) was purified by preparative HPLC (Symmetry C<sub>18</sub>, isocratic, 45% acetonitrile/water) to give 0.88 g (58%) of yellow oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.38-7.15 (m, 3H), 7.00-6.93 (m, 2H), 6.87-6.82 (m, 2H), 4.58 (d, J=5.3 Hz, 2H), 3.86 (s, 3H), 3.83 (s, 3H).

3-[(Thiophen-2-ylmethyl)-amino]-phthalic acid



3-[(Thiophen-2-ylmethyl)-amino]-phthalic acid dimethyl ester (0.88 g, 2.88 mmol) was treated in the same manner as described above for the synthesis 3-(2-methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

2-(2,6-Dioxo-piperidin-3-yl)-4-[(thiophen-2-ylmethyl)-amino]-isoindole-1,3-dione I-126




3-[(Thiophen-2-ylmethyl)-amino]-phthalic acid (2.88 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione. The solid yellow residue was purified by preparative HPLC (Symmetry C<sub>18</sub>, isocratic, 35% acetonitrile/water) to give 0.31 g (29%) of product as a yellow solid: mp 223-225°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.12 (s, 1H), 7.56 (t, J=7.8 Hz, 1H), 7.39 (d, J=4.8 Hz, 1H), 7.23-6.96 (m, 5H), 5.07 (dd, J= 5.3 and 12.4 Hz, 1H), 4.75 (d, J=6.0 Hz, 2H), 2.97-2.83 (m, 1 H), 2.63-2.46 (m, 2H), 2.09-1.99 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.77, 170.03, 168.67, 167.20, 145.60, 142.51, 136.04, 132.15, 126.88,

125.40, 125.00, 117.67, 111.05, 109.80, 48.56, 40.88, 30.96, 22.10; Anal. Calcd. For  $C_{18}H_{15}N_3O_4S$ : C, 58.53; H, 4.09; N, 11.38. Found: C, 58.20; H, 3.96; N, 10.99.

3-(3-Chloro-benzylamino)-phthalic acid dimethyl ester

5




10 3-Chlorobenzaldehyde (1.14 ml, 10 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester. The residue (oil) was purified by chromatography ( $SiO_2$ , 15% ethyl acetate/hexanes) to give 1.61 g (96%) of yellow oil:  $^1H$  NMR ( $CDCl_3$ )  $\delta$  7.35-7.18 (m, 6), 6.83 (d,  $J=7.4$  Hz, 1H), 6.65 (d,  $J=8.5$  Hz, 1H), 4.39 (d,  $J=5.8$  Hz, 2H), 3.86 (s, 3H), 3.84 (s, 3H).

15

3-(3-Chloro-benzylamino)-phthalic acid


20



25 3-(3-Chloro-benzylamino)-phthalic acid dimethyl ester (1.61 g, 4.8 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

4-(3-Chloro-benzylamino)-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione I-127

30




35

3-(3-Chloro-benzylamino)-phthalic acid (4.8 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione. The solid yellow residue was slurried in diethyl ether (30 ml) for 18 h and filtered to give 1.42 g (89 %) of product as a yellow solid: mp 207-209°C;  
5 <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.11 (s, 1H), 7.55-7.28 (m, 6H), 7.03 (d, J=7.0 Hz, 1H), 6.95 (d, J=8.6 Hz, 1H), 5.08 (dd, J= 5.4 and 12.4 Hz, 1H), 4.58 (d, J=6.3 Hz, 2H), 2.98-2.83 (m, 1 H), 2.64-2.46 (m, 2H), 2.08-2.04 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.56, 169.84, 168.59, 167.11, 145.76, 141.68, 136.01, 133.10, 132.16, 130.25, 126.83, 126.68, 125.48, 117.40, 110.81, 109.74, 48.54, 41.54, 44.80, 30.87, 22.05; Anal. Calcd. For C<sub>20</sub>H<sub>16</sub>ClN<sub>3</sub>O<sub>4</sub>: C, 60.38; H, 4.05; N, 10.56. Found: C, 60.22; H, 4.05; N, 10.38.

3-[(Pyridin-3-ylmethyl)-amino]-phthalic acid dimethyl ester

15




20 3-Pyridinecarboxaldehyde (0.94 ml, 10 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester. After removal of pyridine the residue was dissolved in methylene chloride (100 ml) and washed with water (2 x 100 ml), saturated NaHCO<sub>3</sub> (2 x 100ml), brine (1 x 100 ml), dried (magnesium sulfate), and filtered. The solvent was evaporated *in vacuo* to give an oil. The oil was dissolved in  
25 diethyl ether (100 ml) and extracted with 0.1 N HCl (2 x 100 ml). The combined aqueous HCl extracts were washed with diethyl ether (2 x 100 ml) and the pH adjusted to 10 by dropwise addition of saturated Na<sub>2</sub>CO<sub>3</sub>. The product was extracted with diethyl ether (3 x 100 ml). The combined ether extracts were washed with brine (1 x 100 ml), dried (magnesium sulfate), and filtered. The solvent was evaporated *in vacuo* to give 0.91 g (61  
30 %) of product as a light yellow oil which was used without further purification: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.61 (d, J=2.0 Hz, 1H), 8.53 (dd, J=1.1 Hz and 4.8 Hz, 1H), 7.65 (d, J=7.9 Hz, 1H), 7.31-7.21 (m, 3H), 6.85 (d, J=7.3 Hz, 1H), 6.69 (d, J=8.5 Hz, 1H), 4.45 (d, J=5.7 Hz, 2H), 3.87 (s, 3H), 3.85 (s, 3H).

35



3-[(Pyridin-3-ylmethyl)-amino]-phthalic acid


5



3-[(Pyridin-3-ylmethyl)-amino]-phthalic acid dimethyl ester (0.91 g, 3.03 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxy-ethylamino)-  
 10 phthalic acid except the pH of crude reaction mixture was adjusted to 2-3 by dropwise addition of concentrated HCl. The solvent was then evaporated *in vacuo* leaving a dry salt mixture that was used without further purification.

15 2-(2,6-Dioxo-piperidin-3-yl)-4-[(pyridin-3-ylmethyl)-amino]-isoindole-1,3-dione I-128

20




3-[(Pyridin-3-ylmethyl)-amino]-phthalic acid (3.03 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxy-  
 25 ethylamino)-isoindole-1,3-dione. The solid yellow residue was slurried in 50% methanol/ethyl acetate (20 ml) for 18 h to give 0.50 g (46%) of product as a yellow solid: mp 231-233 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.14 (s, 1H), 8.63 (s, 1H), 8.47 (s, 1H), 7.78 (d, J=7.7 Hz, 1H), 7.53 (t, J=7.8 Hz, 1H), 7.39-7.30 (m, 2H), 7.03 (t, J=6.6 Hz, 2H), 5.09 (dd, J= 5.2 and 12.4 Hz, 1H), 4.61 (d, J=6.0 Hz, 2H), 2.97-2.84 (m, 1 H), 2.64-2.48 (m, 2H),  
 30 2.08-2.04 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.78, 170.05, 168.65, 167.22, 148.66, 148.27, 145.75, 136.14, 134.79, 134.54, 132.25, 123.58, 117.50, 110.92, 109.82, 48.57, 42.99, 30.96, 22.12; Anal. Calcd. For C<sub>19</sub>H<sub>16</sub>N<sub>4</sub>O<sub>4</sub>: C, 62.63; H, 4.43; N, 15.38. Found: C, 62.30; H, 4.27; N, 15.30.

35

3-[(5-Carboxy-furan-2-yl)methyl]-amino]-phthalic acid dimethyl ester


5



5-Formyl-furan-2-carboxylic acid (1 g, 7.14 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester except the combined aqueous  $\text{NaHCO}_3$  extracts were washed with methylene chloride (1 x 70 ml) and the pH adjusted to 2-3 by dropwise addition of concentrated HCl. The mixture was then extracted with ethyl acetate (3 x 75 ml). The combined ethyl acetate extracts were washed with brine (1 x 100 ml), dried (magnesium sulfate), and filtered. The solvent was evaporated *in vacuo* to give 0.90 g product as an off-white solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.42-7.36 (m, 1H), 7.16 (d,  $J=3.5$  Hz, 1H), 7.03-6.97 (m, 2H), 6.90 (dd,  $J=1.0$  Hz and 7.3 Hz, 1H), 6.48 (d,  $J=3.5$  Hz, 1H), 4.60 (d,  $J=5.2$  Hz, 2H), 3.82 (s, 3H), 3.81 (s, 3H).

3-[(5-Carboxy-furan-2-yl)methyl]-amino]-phthalic acid

20




3-[(5-Carboxy-furan-2-yl)methyl]-amino]-phthalic acid dimethyl ester (0.90 g, 2.70 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

30

35

5-[[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylamino]-methyl]-furan-2-carboxylic acid I-129

5




3-[(5-Carboxy-furan-2-ylmethyl)-amino]-phthalic acid (1.78 g, 5.65 mmol) was treated in  
 10 the same manner as described above for the synthesis of 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione. The solid yellow residue was purified by preparative HPLC (Symmetry, C<sub>18</sub>, isocratic, 30% acetonitrile/water) to give 0.39 g (36%) of product as a yellow solid: mp 202-204°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 13.01 (bs, 1H), 11.11 (s, 1H), 7.58 (t, J=7.8 Hz, 1H), 7.17-7.07 (m, 4H), 6.51 (d, J=3.2 Hz, 1H), 5.08 (dd, J=5.2  
 15 Hz and 12.4 Hz, 1H), 4.65 (d, J=5.9 Hz, 2H), 2.97-2.83 (m, 1H), 2.63-2.47 (m, 2H), 2.07-2.03 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.78, 170.02, 168.65, 167.20, 159.13, 156.61, 145.58, 144.19, 136.13, 132.17, 118.59, 117.52, 111.24, 109.97, 109.55, 48.59, 30.96, 22.11; Anal. Calcd. For C<sub>19</sub>H<sub>15</sub>N<sub>3</sub>O<sub>7</sub>: C, 56.76; H, 3.89; N, 10.45. Found: C, 56.53; H, 4.16; N, 10.24 + 1.18% H<sub>2</sub>O.

20

3-[(4,5-Dimethyl-furan-2-ylmethyl)-amino]-phthalic acid dimethyl ester

25




4,5-Dimethyl-furan-2-carbaldehyde (0.98 ml, 8 mmol) was treated in the same manner as  
 described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester. The  
 residue (oil) was purified by chromatography (SiO<sub>2</sub>, 10% ethyl acetate/hexanes) to give 0.95  
 30 g (75%) of product as a yellow oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.32 (t, J=8.2 Hz, 1H), 6.97 (bs, 1H), 6.88-6.83 (m, 2H), 5.99 (s, 1H), 4.28 (d, J=5.5 Hz, 2H), 3.86 (s, 3H), 3.83 (s, 3H), 2.17 (s, 3H), 1.87 (s, 3H).

35

3-[(4,5-Dimethyl-furan-2-ylmethyl)-amino]-phthalic acid


5



3-[(4,5-Dimethyl-furan-2-ylmethyl)-amino]-phthalic acid dimethyl ester (0.95 g, 3 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxy-ethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

4-[(4,5-Dimethyl-furan-2-ylmethyl)-amino]-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione I-130

15



20


3-[(4,5-Dimethyl-furan-2-ylmethyl)-amino]-phthalic acid (3 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione. The solid yellow residue was purified by preparative HPLC (Symmetry C<sub>18</sub>, isocratic, 40% acetonitrile/water) to give 0.25 g (22%) of product as a yellow solid: mp 127-129°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.10 (s, 1H), 7.58 (dd, J=7.4 Hz and 8.4 Hz, 1H), 7.17 (d, J=8.6 Hz, 1H), 7.06 (d, J=7.1 Hz, 1H), 6.91 (t, J=6.1 Hz, 1H), 5.11 (s, 1H), 5.06 (dd, J= 5.4 and 12.5 Hz, 1H), 4.45 (d, J=6.0 Hz, 2H), 2.97-2.82 (m, 1 H), 2.63-2.44 (m, 2H), 2.13 (s, 3H), 2.08-1.99 (m, 1H), 1.85 (s, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.75, 170.02, 168.77, 167.21, 148.75, 146.18, 145.82, 136.05, 132.07, 117.57, 114.16, 110.94, 110.55, 109.61, 48.56, 38.94, 30.95, 22.10, 11.08, 9.50; Anal. Calcd. For C<sub>20</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>: C, 62.99; H, 5.02; N, 11.02. Found: C, 62.75; H, 4.97; N, 10.72.

30

35

3-[(Benzofuran-2-ylmethyl)-amino]-phthalic acid dimethyl ester


5



Benzofuran-2-carbaldehyde (1.17 ml, 8 mmol) was treated in the same manner as described above for the synthesis of 3-pentylamino-phthalic acid dimethyl ester. The residue (oil) was  
 10 purified by chromatography (SiO<sub>2</sub>, 50% methylene chloride/hexane) to give 1.12 g (83%) of product as a yellow oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.50-7.43 (m, 2H), 7.33-7.16 (m, 3H), 6.88-6.84 (m, 2H), 6.58 (s, 1H), 4.54 (d, J=6.1 Hz, 2H), 3.86 (s, 3H), 3.85 (s, 3H).

3-[(Benzofuran-2-ylmethyl)-amino]-phthalic acid


15



20 3-[(Benzofuran-2-ylmethyl)-amino]-phthalic acid dimethyl ester (1.12 g, 3.3 mmol) was treated in the same manner as described above for the synthesis of 3-(2-methoxyethylamino)-phthalic acid. The product of the reaction, which contained a mixture of diacid and monomethyl esters, was used without further purification.

4-[(Benzofuran-2-ylmethyl)-amino]-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione I-131


30



3-[(Benzofuran-2-ylmethyl)-amino]-phthalic acid (3.3 mmol) was treated in the same manner as described above for the synthesis of 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxyethylamino)-isoindole-1,3-dione. The solid yellow residue was slurried in methylene  
 35 chloride (20 ml) for 18 h and filtered to give 0.59 g (37%) of product as a yellow solid: mp


199-201 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.14 (s, 1H), 7.61-7.52 (m, 3H), 7.29-7.18 (m, 4H), 7.08 (d, J=7.0 Hz, 1H), 6.80 (s, 1H), 5.10 (dd, J= 5.4 and 12.5 Hz, 1H), 4.77 (d, J=6.1 Hz, 2H), 2.98-2.83 (m, 1 H), 2.64-2.46 (m, 2H), 2.07-2.03 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.76, 170.02, 168.71, 167.20, 155.33, 154.33, 154.22, 145.71, 136.09, 132.15, 127.93, 123.96, 122.81, 120.88, 117.64, 111.19, 110.93, 109.96, 103.85, 48.59, 30.96, 22.12; Anal. Calcd. For C<sub>22</sub>H<sub>17</sub>N<sub>3</sub>O<sub>5</sub>: C, 65.50; H, 4.25; N, 10.42. Found: C, 65.35; H, 4.12; N, 10.34.

4-(3-Chloro-benzylamino)-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione I-132




To a stirred solution of 3-(3-Chloro-benzylamino)-phthalic acid (4 mmol) in pyridine (50 ml) was added 3-amino-3-methyl-piperidine-2,6-dione hydrochloride (0.71 g, 4 mmol). The reaction mixture was heated to reflux for 18 h. The solvent was evaporated *in vacuo* and the residue dissolved in methylene chloride (150 ml). The methylene chloride mixture was washed with water (2 x 100 ml), 0.1N HCl (2 x 100 ml), brine (1 x 100 ml), dried (magnesium sulfate), and filtered. The solvent was evaporated *in vacuo* to give a yellow semi-solid. The residue was purified by preparative HPLC (Symmetry C<sub>18</sub>, isocratic, 50% acetonitrile/water) to give 0.67 g (41%) of product as a yellow solid: mp 199-201 °C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.99 (s, 1H), 7.52-7.28 (m, 6H), 6.97-6.89 (m, 2H), 4.54 (d, J=6.1 Hz, 1H), 2.78-2.50 (m, 3H), 2.08-1.98 (m, 1H), 1.89 (s, 3H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.42, 172.19, 169.62, 167.92, 145.59, 141.82, 136.02, 133.14, 132.06, 130.35, 126.91, 126.78, 125.60, 117.23, 110.42, 109.49, 58.37, 44.74, 29.27, 28.62, 21.02; Anal. Calcd. For C<sub>21</sub>H<sub>18</sub>ClN<sub>3</sub>O<sub>4</sub>: C, 61.24; H, 4.41; N, 10.20. Found: C, 61.30; H, 4.28; N, 9.97.

3-[4-(3-Chloro-benzylamino)-1-oxo-1,3-dihydro-isoindol-2-yl]-piperidine-2,6-dione I-133



3-(4-Amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (1.04 g, 4 mmol) was dissolved in acetic acid (7 ml) with heat. The mixture was cooled slightly and methylene chloride (50 ml) was added slowly followed by 3-chlorobenzaldehyde (0.91 ml, 8 mmol) and sodium triacetoxyborohydride (2.54 g, 12 mmol). The reaction mixture was stirred at room temperature under a nitrogen atmosphere for 2 h, diluted with methylene chloride (100 ml), washed with water (2 x 100 ml), saturated aqueous sodium bicarbonate (2 x 100 ml), brine (100 ml), dried (magnesium sulfate), and filtered. The solvent was evaporated *in vacuo* to give a white solid. The solid was recrystallized from methylene chloride/methanol to give 0.48 g (31%) of product as a white solid: mp 253-255°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.05 (s, 1H), 7.45 (s, 1H), 7.37-7.19 (m, 4H), 6.95 (d, J=7.4 Hz, 1H), 6.63 (d, J=8.0 Hz, 1H), 6.45 (t, J=5.9 Hz, 1H), 5.14 (dd, J= 5.1 and 13.1 Hz, 1H), 4.41 (d, J=5.8 Hz, 2H), 4.34 (d, J=17.3 Hz, 1H), 4.20 (d, J=17.2 Hz, 1H) 3.01-2.87 (m, 1H), 2.66-2.60 (m, 1H), 2.42-2.25 (m, 1 H), 2.08-2.04 (m, 1H); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 172.90, 171.24, 168.74, 143.03, 142.59, 133.08, 132.16, 130.19, 129.12, 126.80, 126.70, 125.73, 112.27, 110.55, 51.54, 45.73, 45.47, 31.25, 22.79; Anal. Calcd. For C<sub>20</sub>H<sub>18</sub>ClN<sub>3</sub>O<sub>3</sub>: C, 62.58, H, 4.73; N, 10.95. Found: C, 62.32; H, 4.61; N, 10.80.

N-{{2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(cyclopentylamino)carboxamide I-134




1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29 g, 1.9 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in acetonitrile (50 mL). After stirring for 20 min, 4-nitrophenyl-N-cyclopentylcarbamate (0.44 g, 1.85 mmol) was added. The mixture was stirred at room temperature for 17 hours. The mixture was filtered to afford N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(cyclopentylamino)carboxamide (0.2 g, 27%) as a white solid: mp 151-153°C; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.13 (s, 1H), 7.86-7.68 (m, 3H), 6.31 (t, 1H), 6.17 (d, J=6.9 Hz, 1H), 5.18-5.11 (dd, J=4.6 and 12.3 Hz, 1H), 4.62 (d, J=5.4 Hz, 2H), 3.83 (q, J=6.4 Hz, 2H), 2.96-2.85 (m, 1H), 2.64-2.50 (m, 2H), 2.07-2.04 (m, 1H),

1.70-1.29 (m, 8H);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  172.77, 169.83, 167.59, 167.04, 157.63, 141.12, 134.64, 133.43, 131.52, 128.99, 121.64, 51.09, 48.83, 38.70, 32.91, 30.94, 23.16, 21.99; Anal. Calcd. For  $\text{C}_{20}\text{H}_{22}\text{N}_4\text{O}_5$ : C, 60.29; H, 5.57; N, 14.06. Found: C, 60.09; H, 5.66; N, 14.15.

5

N- {[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl} (3-pyridylamino)carboxamide Hydrochloride I-135

10



15

20

25


30

1,8-Diazabicyclo[5,4,0]undec-7-ene (0.29 g, 1.9 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.85 mmol) in acetonitrile (50 mL). After stirring for 20 min, (2,5-dioxopyrrolidinyl)-N-(3-pyridyl)carboxamide (0.44 g, 1.85 mmol) was added. The mixture was stirred at room temperature for 17 hours. The mixture was filtered and the solid was recrystallized from methanol (25 mL) to give N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl} (3-pyridylamino)carboxamide (0.23 g, 30%). 2N HCl/ether was added to a stirred solution of N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl} (3-pyridylamino)carboxamide (0.23 g) in methanol (5 mL) and ethyl acetate (10 mL). The mixture was stirred for 2 hours and filtered to give N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl} (3-pyridylamino)carboxamide hydrochloride (0.2 g) as a white solid: mp 263-265°C;  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  11.17 (s, 1H), 10.43 (s, 1H), 9.08 (d, J=1.7 Hz, 1H), 8.44 (d, J=5.4 Hz, 1H), 8.33-8.29 (dd, J=1.2 and 8.5 Hz, 1H), 7.93-7.78 (m, 4H), 7.55 (t, J=6.0 Hz, 1H), 5.21-5.14 (dd, J=5.4 and 12.5 Hz, 1H), 4.80 (d, J=5.9 Hz, 2H), 2.97-2.84 (m, 1H), 2.65-2.50 (m, 2H), 2.10-2.06 (m, 1H);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  172.78, 169.83, 167.54, 166.96, 154.78, 139.97, 139.51, 134.80, 134.37, 133.35, 132.34, 131.60, 130.43, 127.17, 127.10, 121.99, 48.88, 38.68, 30.94, 21.99; Anal. Calcd. For  $\text{C}_{20}\text{H}_{18}\text{N}_5\text{O}_5\text{Cl}$ : C, 54.12; H, 4.09; N, 15.78; Cl, 7.99. Found: C, 54.11; H, 4.10; N, 15.44; Cl, 7.81.

35




N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}piperidylcarboxamide  
**I-136**




Diisopropylethylamine (0.88 g, 6.79 mmol) was added to a stirred suspension of 4-  
 10 (aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (1.0 g, 3.09  
 mmol) in acetonitrile (50 mL). After stirring for 20 min, the mixture was slowly added to a  
 stirred solution of triphosgene (0.34 g, 1.14 mmol) in acetonitrile (15 mL) over 30 min.  
 After a further 10 min of stirring, a solution of piperidine (0.26 g, 3.09 mmol) and  
 diisopropylethylamine (0.48 g, 3.71 mmol) in acetonitrile (10 mL) was added in one  
 15 portion. The mixture was stirred at room temperature for 17 hours. The mixture was  
 concentrated and the residue was dissolved in methylene chloride (80 mL). The methylene  
 chloride solution was washed with 10% KHSO<sub>4</sub> (30 mL), H<sub>2</sub>O (2x30 mL), brine (30 mL)  
 and dried (MgSO<sub>4</sub>). Solvent was removed and the residue was purified by chromatography  
 (CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 97.5:2.5) to give N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-  
 20 yl]methyl}piperidylcarboxamide (0.66 g, 53%) as a white solid: mp 156-158°C; <sup>1</sup>H NMR  
 (DMSO-d<sub>6</sub>) δ 11.14 (s, 1H), 7.86-7.75 (m, 2H), 7.69 (d, J=7.4 Hz, 1H), 7.12 (t, J=5.6 Hz,  
 1H), 5.18-5.11 (dd, J=5.3 and 12.5 Hz, 1H), 4.69 (d, J=5.5 Hz, 2H), 3.42-3.32 (m, 4H),  
 2.96-2.83 (m, 1H), 2.63-2.50 (m, 2H), 2.08-2.03 (m, 1H), 1.54-1.44 (m, 6H); <sup>13</sup>C NMR  
 (DMSO-d<sub>6</sub>) δ 172.74, 169.83, 167.60, 157.26, 141.46, 134.58, 132.82, 131.38, 126.74,  
 25 121.43, 48.82, 44.37, 30.93, 25.37, 24.12, 21.98; Anal. Calcd. For C<sub>20</sub>H<sub>22</sub>N<sub>4</sub>O<sub>5</sub>+0.22 H<sub>2</sub>O:  
 C, 59.70; H, 5.62; N, 13.92. Found: C, 60.14; H, 5.59; N, 13.47.

Tert-Butyl 4-(N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-  
 yl]methyl}(carbamoyl)piperazinecarboxylate **I-137**



Diisopropylethylamine (1.05 g, 8.16 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (1.2 g, 3.71 mmol). After stirring for 20 min, the mixture was added slowly to a stirred solution of triphosgene (0.41 g, 1.37 mmol) in acetonitrile (20 mL) over 30 min. After a further 10 min of stirring, a solution of t-BOC-1-piperazine carboxylate (0.69 g, 3.71 mmol) and diisopropylethylamine (0.58 g, 4.45 mmol) in acetonitrile (10 mL) was added in one portion. The mixture was stirred at room temperature for 17 hours. The mixture was concentrated and the residue was dissolved in methylene chloride (100 mL). The methylene chloride solution was washed with water (2x40 mL), brine (40 mL) and dried (MgSO<sub>4</sub>). Solvent was removed and the residue was purified by chromatography (CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 97.5:2.5) to give tert-butyl 4-(N-{[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(carbamoyl)piperazinecarboxylate (0.98 g, 52%) as a white solid: mp 158-160°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.14 (s, 1H), 7.86-7.69 (m, 3H), 7.27 (t, J=5.5 Hz, 1H), 5.18-5.11 (dd, J=5.4 and 12.5 Hz, 1H), 4.71 (d, J=5.4 Hz, 2H), 3.33 (s, 8H), 2.98-2.83 (m, 1H), 2.63-2.49 (m, 2H), 2.08-2.04 (m, 1H), 1.41 (s, 9H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.74, 170.04, 169.83, 167.58, 167.03, 157.30, 153.86, 140.99, 136.52, 134.64, 132.94, 131.39, 126.80, 121.53, 111.33, 79.03, 74.43, 48.82, 43.22, 30.92, 28.03, 21.98; Anal. Calcd. For C<sub>24</sub>H<sub>29</sub>N<sub>5</sub>O<sub>7</sub>+0.13H<sub>2</sub>O: C, 57.44; H, 5.88; N, 13.95. Found: C, 57.83; H, 6.09; N, 13.45.


N-{[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl}(diethylamino)carboxamide I-138



Diisopropylethylamine (0.88 g, 6.80 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (1.0 g, 3.09 mmol) in acetonitrile (50 mL). After stirring for 20 min, the mixture was slowly added to a stirred solution of triphosgene (0.34 g, 1.14 mmol) in acetonitrile (20 mL) over 20 min. After a further 10 min of stirring, a solution of diethylamine (0.23 g, 3.09 mmol) and diisopropylethylamine (0.48 g, 3.71 mmol) in acetonitrile (10 mL) was added in one portion. The mixture was stirred at room temperature for 17 hours. The mixture was concentrated and the residue was dissolved in methylene chloride (80 mL). The methylene

chloride solution was washed with 1N HCl (40 mL), H<sub>2</sub>O (2x40 mL), brine (40 mL) and dried (MgSO<sub>4</sub>). Solvent was removed and the residue was purified by chromatography (CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 97.5:2.5) to afford N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxisoindolin-4-yl)methyl}(diethylamino)carboxamide (0.8 g, 67%) as a white solid: mp 142-144°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.14 (s, 1H), 7.86-7.66 (m, 3H), 6.91 (t, J=5.7 Hz, 1H), 5.19-5.12 (dd, J=5.4 and 12.6 Hz, 1H), 4.70 (d, J=5.6 Hz, 2H), 3.26 (q, J=6.9 Hz, 4H), 2.98-2.83 (m, 1H), 2.64-2.49 (m, 2H), 2.08-2.04 (m, 1H), 1.06 (t, J=7.0 Hz, 6H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.73, 169.81, 167.63, 167.04, 156.74, 134.54, 132.80, 131.38, 126.72, 121.38, 48.82, 40.24, 30.92, 21.97, 13.90; Anal. Calcd. For C<sub>19</sub>H<sub>22</sub>N<sub>4</sub>O<sub>5</sub>: C, 59.06; H, 5.74; N, 14.50. Found: C, 58.71; H, 5.76; N, 14.10.

Cyclopropyl-N-{{2-(3-methyl-2,6-dioxo(3-piperidyl))-1,3-dioxisoindolin-4-yl)methyl}carboxamide I-139



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.68 g, 4.44 mmol) was added to a stirred suspension of 4-(aminomethyl)-2-(3-methyl-2,6-dioxo(3-piperidyl))isoindoline-1,3-dione hydrochloride (0.6 g, 1.78 mmol) in acetonitrile (50 mL). After stirring for 20 min, cyclopropanecarbonyl chloride (0.22 g, 2.14 mmol) was added. The mixture was stirred at room temperature for 17 hours. The mixture was filtered and the solid was washed with acetonitrile (10 mL) to give cyclopropyl-N-{{2-(3-methyl-2,6-dioxo(3-piperidyl))-1,3-dioxisoindolin-4-yl)methyl}carboxamide (0.49 g, 74%) as a white solid: mp 243-245°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.04 (s, 1H), 8.68 (t, J=5.8 Hz, 1H), 7.84-7.63 (m, 3H), 5.76 (s, 1H), 4.70 (d, J=5.8 Hz, 2H), 2.71-2.54 (m, 2H), 2.10-2.02 (m, 1H), 1.90 (s, 3H), 1.68-1.63 (, 1H), 0.71-0.68 (m, 4H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 173.07, 172.20, 172.15, 168.29, 167.68, 139.01, 134.61, 133.19, 131.40, 126.78, 121.50, 58.71, 54.88, 37.72, 29.09, 28.57, 21.00, 13.51, 6.42; Anal. Calcd. For C<sub>19</sub>H<sub>19</sub>N<sub>3</sub>O<sub>5</sub>: C, 61.78; H, 5.18; N, 11.38. Found: C, 61.53; H, 5.20; N, 11.39.

Methyl 3-bromo-2-methylbenzoate

A mixture of 3-bromo-2-methylbenzoic acid (16 g, 74.4 mmol), sodium bicarbonate (12.5 g, 148.8 mmol) and iodomethane (21.2 g, 148.8 mmol) in DMF (160 mL) was heated at 60°C for 2 hours. The mixture was cooled to room temperature and poured into ice water (400

mL). The mixture was extracted with ethyl acetate (4x100 mL). The EtOAc solution was washed with water (3x100 mL), brine (100 mL) and dried (MgSO<sub>4</sub>). Solvent was removed to give methyl 3-bromo-2-methylbenzoate (17.6 g, 100%) as an oil: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.70 (t, J=7.6Hz, 2H), 7.08 (t, J=7.9 Hz, 1H), 3.90 (s, 3H), 2.67 (s, 3H).

Methyl 3-bromo-2-bromomethylbenzoate

A mixture of methyl 3-bromo-2-methylbenzoate (17.0 g, 74.22 mmol) and N-bromosuccinimide (15.85 g, 89.06 mmol) in acetonitrile (200 mL) was heated under gently refluxing for 17 hours while a 200W bulb situated 2 cm away was shining on the reaction flask. The mixture was cooled to room temperature and solvent was removed. The residue was dissolved in ethyl acetate (200 mL) and washed with water (3x80 mL), brine (80 mL) and dried (MgSO<sub>4</sub>). Solvent was removed to give methyl 3-bromo-2-bromomethylbenzoate (24.5 g, 97% by HPLC): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.90-7.86 (dd, J=1.0 and 7.9 Hz, 1H), 7.78-7.74 (dd, J=1.2 and 8.2 Hz, 1H), 7.22 (t, J=7.9 Hz, 1H), 5.12 (s, 2H), 3.95 (s, 3H).

Tert-Butyl 2-(4-bromo-1-oxoisindolin-2-yl)-4-carbamoylbutanoate

Triethylamine (4.66 g, 46.09 mmol) was added to a stirred suspension of methyl 3-bromo-2-bromomethylbenzoate (6.45 g, 20.95 mmol) and L-glutamine t-butyl ester hydrochloride (5.0 g, 20.95 mmol) in THF (100 mL). The mixture was heated to reflux for 18 hours. The mixture was cooled to room temperature and solvent was removed. The residue was dissolved in methylene chloride (100 mL) and washed with water (2x80 mL), brine (80 mL) and dried (MgSO<sub>4</sub>). Solvent was removed and the residue was purified by chromatography (CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 97.5:2.5) to afford tert-butyl 2-(4-bromo-1-oxoisindolin-2-yl)-4-carbamoylbutanoate (3.97 g, 48%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.74 (d, J=7.1 Hz, 1H), 7.65 (d, J=7.9 Hz, 1H), 7.35 (t, J=7.8 Hz, 1H), 6.15 (s, 1H), 5.77 (s, 1H), 5.02-4.96 (m, 1H), 4.58 (d, J=17.4 Hz, 1H), 4.34 (d, J=17.4 Hz, 1H), 2.45-2.10 (m, 4H), 1.46 (s, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 173.83, 169.42, 168.74, 142.28, 134.70, 133.65, 129.80, 122.70, 117.75, 82.66, 54.29, 47.80, 32.27, 27.95, 25.54.

Tert-Butyl 2-(4-cyano-1-oxoisindolin-2-yl)-4-carbamoylbutanoate


A mixture of tert-butyl 2-(4-bromo-1-oxoisindolin-2-yl)-4-carbamoylbutanoate (1.2 g, 3.02 mmol), zinc cyanide (0.21 g, 1.81 mmol), tris(dibenzylideneacetone)dipalladium (0.06g, 0.06 mmol), 1,1'-bis(diphenylphosphino)ferrocene (0.067 g, 0.12 mmol) in deoxygenated DMF (15 mL) was heated to 120°C under N<sub>2</sub> for 6 hours. The mixture was cooled to room temperature and poured into EtOAc (100 mL) and sat. NaHCO<sub>3</sub> (40 mL). The EtOAc

solution was washed with water (2x40 mL), brine (40 mL), and dried (MgSO<sub>4</sub>). Solvent was removed and the residue was purified by chromatography (CH<sub>2</sub>Cl<sub>2</sub>: CH<sub>3</sub>OH 97.5:2.5) to afford tert-butyl 2-(4-cyano-1-oxoisindolin-2-yl)-4-carbamoylbutanoate (0.77 g, 74%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.07-8.03 (dd, J=0.7 and 7.4 Hz, 1H), 7.86-7.83 (dd, J=1.0 and 7.9 Hz, 1H), 7.61 (t, J=7.6 Hz, 1H), 5.91 (s, 1H), 5.68 (s, 1H), 5.03-4.98 (m, 1H), 4.85 (d, J=17.9 Hz, 1H), 4.62 (d, J=17.9 Hz, 1H), 2.48-2.41(m, 4H), 1.46 (s, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 173.56, 169.21, 167.70, 145.23, 134.94, 133.11, 128.93, 128.21, 115.65, 107.86, 82.98, 63.68, 54.47, 46.55, 32.34, 27.95, 25.46.

10 2-(4-Cyano-1-oxoisindolin-2-yl)-4-carbamoylbutanoic acid

A mixture of tert-butyl 2-(4-cyano-1-oxoisindolin-2-yl)-4-carbamoylbutanoate (1.0 g, 2.91 mmol) and trifluoroacetic acid (5 mL) was stirred for 2 hours. The mixture was concentrated and the residue was crystallized from ether (15 mL) to give 2-(4-cyano-1-oxoisindolin-2-yl)-4-carbamoylbutanoic acid (0.74 g, 89%) as a white solid: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.12 (b, 1H), 8.12 (d, J=7.6 Hz, 1H), 8.03 (d, J=7.6 Hz, 1H), 7.73 (t, J=7.6 Hz, 1H), 7.23 (s, 1H), 6.76 (s, 1H), 4.80-4.62 (m, 3H), 2.32-2.09 (m, 4H).

20 2-(2,6-Dioxo(3-piperidyl))-1-oxoisindoline-4-carbonitrile




A mixture of 2-(4-cyano-1-oxoisindolin-2-yl)-4-carbamoylbutanoic acid (0.6 g, 2.09 mmol) and 1,1-carbonyldiimidazole (0.44 g, 2.72 mmol) in acetonitrile (20 mL) was heated to reflux for 2 hours. The mixture was cooled to room temperature and filtered to give 2-(2,6-dioxo(3-piperidyl))-1-oxoisindoline-4-carbonitrile (0.44 g, 83%) as a white solid: mp 312-314°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.04 (s, 1H), 8.16 (d, J=7.0 Hz, 1H), 8.06 (d, J=7.1 Hz, 1H), 7.74 (t, J=7.7 Hz, 1H), 5.20-5.13 (dd, J=5.2 and 13.2 Hz, 1H), 4.69 (d, J=18.1 Hz, 1H), 4.57 (d, J=18.1 Hz, 1H), 2.97-2.85 (m, 1H), 2.64-2.45 (m, 2H), 2.05-1.97 (m, 1H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.81, 170.69, 166.49, 145.36, 135.37, 132.85, 129.36, 127.91, 116.01, 107.09, 51.77, 46.72, 31.14, 22.22; Anal. Calcd. For C<sub>14</sub>H<sub>11</sub>N<sub>3</sub>O<sub>3</sub>: C, 62.45; H, 4.12; N, 15.61. Found: C, 62.26; H, 4.10; N, 15.51.

35 3-[4-(Aminomethyl)-1-oxoisindolin-2-yl]piperidine-2,6-dione Hydrochloride


A mixture of 2-(2,6-dioxo(3-piperidyl))-1-oxoisindoline-4-carbonitrile (1.0 g, 3.71 mmol) and 10% Pd/C (0.2 g) in 4N HCl (20 mL) and methanol (600 mL) was hydrogenated at 50 psi for 17 hours. The mixture was filtered through celite and the filtrate was concentrated. The residue was stirred with ether (30 mL) to give 3-[4-(aminomethyl)-1-oxoisindolin-2-yl]piperidine-2,6-dione hydrochloride (1.1 g, 99%) as a white solid:  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  11.06 (s, 1H), 8.64 (s, 3H), 7.79 (d,  $J=7.5$  Hz, 1H), 7.74 (d,  $J=7.4$  Hz, 1H), 7.59 (t,  $J=7.5$  Hz, 1H), 5.21-5.14 (dd,  $J=5.0$  and 13.2 Hz, 1H), 4.69 (d,  $J=17.4$  Hz, 1H), 4.51 (d,  $J=17.4$  Hz, 1H), 4.09 (d,  $J=4.9$  Hz, 2H), 3.00-2.87 (m, 1H), 2.66-2.35 (m, 2H), 2.03-1.98 (m, 1H);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  172.87, 170.92, 167.77, 141.58, 132.50, 131.88, 129.39, 128.48, 123.25, 51.52, 46.19, 38.39, 31.17, 22.69.

N-{[2-(2,6-Dioxo(3-piperidyl))-1-oxoisindolin-4-yl]methyl}cyclopropylcarboxamide  
**I-140**



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.7 g, 4.62 mmol) was added to a stirred suspension of 3-[4-(aminomethyl)-1-oxoisindolin-2-yl]piperidine-2,6-dione hydrochloride (0.65 g, 2.10 mmol) in acetonitrile (50 mL). After stirring for 30 min, cyclopropanecarbonyl chloride (0.24 g, 2.31 mmol) was added. The mixture was stirred at room temperature for 17 hours. The mixture was filtered and the solid was recrystallized from methanol (100 mL) to afford N-{[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]methyl}cyclopropylcarboxamide (0.36 g, 50%) as a white solid: mp 262-264°C;  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  11.02 (s, 1H), 8.63 (t,  $J=5.7$  Hz, 1H), 7.65-7.47 (m, 3H), 5.18-5.11 (dd,  $J=5.0$  and 13.2 Hz, 1H), 4.46 (d,  $J=17.3$  Hz, 1H), 4.38-4.31 (m, 3H), 3.00-2.85 (m, 1H), 2.65-2.30 (m, 2H), 2.04-1.99 (m, 1H), 1.64-1.54 (m, 1H), 0.69-0.65 (m, 4H);  $^{13}\text{C}$  NMR (DMSO- $d_6$ )  $\delta$  172.82, 172.69, 170.96, 168.03, 140.08, 134.76, 131.66, 130.70, 128.30, 121.64, 51.54, 46.15, 31.16, 22.59, 13.47, 6.27; Anal. Calcd. For  $\text{C}_{18}\text{H}_{19}\text{N}_3\text{O}_4$ : C, 63.33; H, 5.61; N, 12.31. Found: C, 62.97; H, 5.55; N, 12.33.

N-[[2-(2,6-Dioxo(3-piperidyl))-1-oxoisindolin-4-yl]methyl](ethylamino)carboxamide  
**I-141**



1,8-Diazabicyclo[5,4,0]undec-7-ene (0.44 g, 2.91 mmol) was added to a stirred suspension of 3-[4-(aminomethyl)-1-oxoisindolin-2-yl]piperidine-2,6-dione hydrochloride (0.6 g, 1.94 mmol) in acetonitrile (50 mL). After stirring for 30 min, ethyl isocyanate (0.21 g, 2.91 mmol) was added. The mixture was stirred at room temperature for 17 hours. Solvent was removed and the residue was stirred with methylene chloride (70 mL) to give N-[[2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl]methyl](ethylamino)carboxamide (0.28 g, 42%) as a white solid: mp 341-343°C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 11.02 (s, 1H), 7.62-7.47 (m, 3H), 6.40 (t, J=5.9 Hz, 1H), 5.95 (t, J=5.6 Hz, 1H), 5.18-5.10 (dd, J=5.3 and 13.3 Hz, 1H), 4.52-4.30 (dd, J=17.3 and 37.7 Hz, 2H), 4.28 (d, J=5.9 Hz, 2H), 3.04 (q, J=7.2 Hz, 2H), 3.01-2.86 (m, 1H), 2.65-2.34 (m, 2H), 2.04-1.99 (m, 1H), 0.98 (t, J=7.2 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 172.82, 170.97, 168.10, 157.86, 139.78, 136.24, 131.55, 130.25, 128.16, 121.32, 51.51, 46.13, 34.14, 31.16, 22.59, 15.67; Anal. Calcd for C<sub>17</sub>H<sub>20</sub>N<sub>4</sub>O<sub>4</sub>: C, 59.29; H, 5.85; N, 16.27. Found: C, 58.98; H, 5.85; N, 16.89.


The present invention is not to be limited in scope by the specific embodiments disclosed in the examples that are intended as illustrations of a few aspects of the invention and any embodiments that are functionally equivalent are within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art and are intended to fall within the appended claims.

What is claimed is:

1. A compound of the formula:

5

10



or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof, wherein:

15

one of X and Y is C=O and the other is CH<sub>2</sub> or C=O;

R<sup>1</sup> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, C(O)R<sup>3</sup>, C(S)R<sup>3</sup>,  
 20 C(O)OR<sup>4</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, C(O)NHR<sup>3</sup>, C(S)NHR<sup>3</sup>, C(O)NR<sup>3</sup>R<sup>3'</sup>, C(S)NR<sup>3</sup>R<sup>3'</sup> or (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>;

R<sup>2</sup> is H, F, benzyl, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, or (C<sub>2</sub>-C<sub>8</sub>)alkynyl;

25 R<sup>3</sup> and R<sup>3'</sup> are independently (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>0</sub>-C<sub>8</sub>)alkyl-N(R<sup>6</sup>)<sub>2</sub>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-C(O)OR<sup>5</sup>, (C<sub>1</sub>-C<sub>8</sub>)alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>;

30 R<sup>4</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>, benzyl, aryl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>1</sub>-C<sub>6</sub>)heterocycloalkyl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

R<sup>5</sup> is (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>2</sub>-C<sub>8</sub>)alkenyl, (C<sub>2</sub>-C<sub>8</sub>)alkynyl, benzyl, aryl, or (C<sub>2</sub>-C<sub>5</sub>)heteroaryl;

35



each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl-C(O)O- $R^5$  or the  $R^6$  groups can join to form a heterocycloalkyl group;

- 5 n is 0 or 1; and the \* represents a chiral-carbon center; with the proviso that when n is 0 then  $R^1$  is not H.

2. The compound of claim 1, wherein the compound is the R-enantiomer or substantially R.

10

3. The compound of claim 1, wherein the compound is the S-enantiomer or substantially S.

4. The compound of claim 1, wherein the compound is a racemic mixture.

15


5. The compound of claim 1, wherein the enantiomeric excess is about 90% ee or more.

6. A compound of claim 1, wherein  $R^2$  is H or  $(C_1-C_4)$ alkyl.

20

7. A compound of claim 1, wherein  $R^1$  is H,  $(C_1-C_4)$ alkyl,  $CH_2OCH_3$ ,  $CH_2CH_2OCH_3$ , or

25



wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl-N( $R^6$ )<sub>2</sub>,  $(C_1-C_8)$ alkyl-OR<sup>5</sup>,  $(C_1-C_8)$ alkyl-C(O)OR<sup>5</sup>,  $(C_1-C_8)$ alkyl-O(CO)R<sup>5</sup>, or C(O)OR<sup>5</sup>, or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl ring.

8. A compound of claim 1, wherein  $R^1$  is C(O)R<sup>3</sup>.


35

9. A compound of claim 1, wherein  $R^1$  is  $C(O)OR^4$ .

5

10. The compound of claim 1 having the formula:

10



- 15 or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof, wherein:

$R^1$  is H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(S)R^3$ ,  
 20  $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(O)NHR^3$ ,  $C(S)NHR^3$ ,  $C(O)NR^3R^3$ ,  $C(S)NR^3R^3$  or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;

$R^2$  is H or  $(C_1-C_8)$ alkyl;

25  $R^3$  and  $R^3$  are independently  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ ;

30  $R^4$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  $(C_1-C_4)$ alkyl- $OR^5$ , benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl, or  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl;

$R^5$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, or  $(C_2-C_5)$ heteroaryl;


35

each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl-C(O)O- $R^5$  or the  $R^6$  groups can join to form a heterocycloalkyl group; and

5 the \* represents a chiral-carbon center.

11. A compound of claim 10, wherein  $R^1$  is H,  $(C_1-C_4)$ alkyl,  $CH_2OCH_3$ ,  $CH_2CH_2OCH_3$ , or

10



15

wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl-OR $^5$ ,  $(C_1-C_8)$ alkyl-C(O)OR $^5$ ,  $(C_1-C_8)$ alkyl-O(CO)R $^5$ , or C(O)OR $^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl ring.


20

12. A compound of claim 10, wherein  $R^1$  is C(O)R $^3$ .

13. A compound of claim 10, wherein  $R^1$  is C(O)OR $^4$ .

14. The compound of claim 1 having the formula:

25



30

or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof, wherein:

35

$R^1$  is H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(S)R^3$ ,  $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(O)NHR^3$ ,  $C(S)NHR^3$ ,  $C(O)NR^3R^3$ ,  $C(S)NR^3R^3$  or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;

5

$R^2$  is H or  $(C_1-C_8)$ alkyl;

$R^3$  and  $R^3$  are independently  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,

10  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ ;

$R^4$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  $(C_1-C_4)$ alkyl- $OR^5$ , benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl, or  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl;

15


$R^5$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, or  $(C_2-C_5)$ heteroaryl;

each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl- $C(O)O-R^5$  or the  $R^6$  groups can join to

20 form a heterocycloalkyl group; and

the \* represents a chiral-carbon center.

15. A compound of claim 14, wherein  $R^1$  is H,  $(C_1-C_4)$ alkyl,  $CH_2OCH_3$ ,  
25  $CH_2CH_2OCH_3$ , or




30 wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl  
35 ring.

16. A compound of claim 14, wherein  $R^1$  is  $C(O)R^3$ .

17. A compound of claim 14, wherein  $R^1$  is  $C(O)OR^4$ .

5 18. A compound of claim 1 having the formula:



or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer,  
15 racemate, or mixture of stereoisomers thereof, wherein:

$R^1$  is H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  
 $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(S)R^3$ ,  
 $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(O)NHR^3$ ,  
20  $C(S)NHR^3$ ,  $C(O)NR^3R^3$ ,  $C(S)NR^3R^3$  or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;

$R^2$  is H or  $(C_1-C_8)$ alkyl;

$R^3$  and  $R^3$  are independently  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  
25  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  
 $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  
 $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ ;

$R^4$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  $(C_1-C_4)$ alkyl- $OR^5$ , benzyl, aryl,  
30  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl, or  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl;


$R^5$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, or  $(C_2-C_5)$ heteroaryl;

each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  
35 benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl- $C(O)O-R^5$  or the  $R^6$  groups can join to  
form a heterocycloalkyl group; and

the \* represents a chiral-carbon center.

19. A compound of claim 18, wherein  $R^1$  is H,  $(C_1-C_4)$ alkyl,  $CH_2OCH_3$ ,  $CH_2CH_2OCH_3$  or

5



10 wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl  
15 ring.


20. A compound of claim 18, wherein  $R^1$  is  $C(O)R^3$ .

21. A compound of claim 18, wherein  $R^1$  is  $C(O)OR^4$ .

20

22. A compound of claim 1 having the formula:

25



30 or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof, wherein:

$R^1$  is H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(S)R^3$ ,  
35  $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(O)NHR^3$ ,  $C(S)NHR^3$ ,  $C(O)NR^3R^3$ ,  $C(S)NR^3R^3$  or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;

- $R^3$  and  $R^{3'}$  are independently  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  
 5  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ ;

$R^4$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  $(C_1-C_4)$ alkyl- $OR^5$ , benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl, or  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl;

- 10  $R^5$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, or  $(C_2-C_5)$ heteroaryl;


each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl- $C(O)O-R^5$  or the  $R^6$  groups can join to form a heterocycloalkyl group; and

15

the \* represents a chiral-carbon center.

23. A compound of claim 22, wherein  $R^1$  is  $(C_1-C_8)$ alkyl, benzyl,  $CH_2OCH_3$ ,  $CH_2CH_2OCH_3$ , or

20



- 25 wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl  
 30 ring.


24. A compound of claim 22, wherein  $R^1$  is  $C(O)R^3$ .

25. A compound of claim 24, wherein  $R^3$  is  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  
 35  $(C_1-C_8)$ alkyl, aryl, or  $(C_0-C_4)$ alkyl- $OR^5$ .

26. A compound of claim 25, wherein heteroaryl is pyridyl, furyl, or thienyl.

27. A compound of claim 22, wherein  $R^1$  is  $C(O)OR^4$ .

5 28. A compound of claim 1 having the formula:



or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer,  
15 racemate, or mixture of stereoisomers thereof, wherein:

$R^1$  is H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  
 $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(S)R^3$ ,  
 $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(O)NHR^3$ ,  
20  $C(S)NHR^3$ ,  $C(O)NR^3R^3$ ,  $C(S)NR^3R^3$  or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;

$R^3$  and  $R^3$  are independently  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  
 $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  
 $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  
25  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ ;

$R^4$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  $(C_1-C_4)$ alkyl- $OR^5$ , benzyl, aryl,  
 $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl, or  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl;

30  $R^5$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, or  $(C_2-C_5)$ heteroaryl;

each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  
benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl- $C(O)O-R^5$  or the  $R^6$  groups can join to  
form a heterocycloalkyl group; and


35

the \* represents a chiral-carbon center.



29. A compound of claim 28, wherein  $R^1$  is  $(C_1-C_8)$ alkyl, benzyl,  $CH_2OCH_3$ ,  $CH_2CH_2OCH_3$ , or

5



wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,   
 10  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl ring.

15

30. A compound of claim 28, wherein  $R^1$  is  $C(O)R^3$ .

31. A compound of claim 30, wherein  $R^3$  is  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_1-C_8)$ alkyl, aryl, or  $(C_0-C_4)$ alkyl- $OR^5$ .


20

32. A compound of claim 31, wherein heteroaryl is pyridyl, furyl, or thienyl.

33. A compound of claim 28, wherein  $R^1$  is  $C(O)OR^4$ .

34. A compound of claim 1 having the formula:

25



30

or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer, racemate, or mixture of stereoisomers thereof, wherein:

35

$R^1$  is H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $C(O)R^3$ ,  $C(S)R^3$ ,  $C(O)OR^4$ ,  $(C_1-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $C(O)NHR^3$ ,  $C(S)NHR^3$ ,  $C(O)NR^3R^3$ ,  $C(S)NR^3R^3$  or  $(C_1-C_8)$ alkyl- $O(CO)R^5$ ;

5

$R^3$  and  $R^3$  are independently  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ ;

10

$R^4$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl,  $(C_1-C_4)$ alkyl- $OR^5$ , benzyl, aryl,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl, or  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl;

$R^5$  is  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, or  $(C_2-C_5)$ heteroaryl;


15

each occurrence of  $R^6$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl,  $(C_2-C_5)$ heteroaryl, or  $(C_0-C_8)$ alkyl- $C(O)O-R^5$  or the  $R^6$  groups can join to form a heterocycloalkyl group; and

20 the \* represents a chiral-carbon center.

35. A compound of claim 34, wherein  $R^1$  is  $(C_1-C_8)$ alkyl, benzyl,  $CH_2OCH_3$ ,  $CH_2CH_2OCH_3$ , or

25



wherein Q is O or S, and each occurrence of  $R^7$  is independently H,  $(C_1-C_8)$ alkyl,  $(C_3-C_7)$ cycloalkyl,  $(C_2-C_8)$ alkenyl,  $(C_2-C_8)$ alkynyl, benzyl, aryl, halogen,  $(C_0-C_4)$ alkyl- $(C_1-C_6)$ heterocycloalkyl,  $(C_0-C_4)$ alkyl- $(C_2-C_5)$ heteroaryl,  $(C_0-C_8)$ alkyl- $N(R^6)_2$ ,  $(C_1-C_8)$ alkyl- $OR^5$ ,  $(C_1-C_8)$ alkyl- $C(O)OR^5$ ,  $(C_1-C_8)$ alkyl- $O(CO)R^5$ , or  $C(O)OR^5$ , or adjacent occurrences of  $R^7$  can be taken together to form a bicyclic alkyl or aryl ring.

35

36. A compound of claim 34, wherein  $R^1$  is  $C(O)R^3$ .

37. A compound of claim 36, wherein R<sup>3</sup> is (C<sub>0</sub>-C<sub>4</sub>)alkyl-(C<sub>2</sub>-C<sub>5</sub>)heteroaryl, (C<sub>1</sub>-C<sub>8</sub>)alkyl, aryl, or (C<sub>0</sub>-C<sub>4</sub>)alkyl-OR<sup>5</sup>.
38. A compound of claim 37, wherein heteroaryl is pyridyl, furyl, or thienyl.
39. A compound of claim 34, wherein R<sup>1</sup> is C(O)OR<sup>4</sup>.
40. A compound of claim 1 having the formula:
- I-1 (2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-ylmethyl)  
-carbamic acid *tert*-butyl ester;
- I-2 4-(aminomethyl)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione;
- I-3 *N*-(2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-ylmeth  
yl)-acetamide;
- I-4 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoiindolin-4-yl)methyl}cyclopropyl-  
carboxamide;
- I-5 (2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-ylmethyl)  
-carbamic acid ethyl ester;
- I-6 2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-ylmethyl)-  
carbamic acid benzyl ester;
- I-7 2-chloro-*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoiindolin-4-  
yl)methyl}acetamide;
- I-8 2-(dimethylamino)-*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoiindolin-4-  
yl)methyl}-acetamide;
- I-9 1-*tert*-butyl-3-(2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoind  
ol-4-ylmethyl)-urea;
- I-10 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoiindolin-4-yl)methyl}-3,3-  
dimethylbutanamide;

- I-11 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)-3-pyridylcarboxamide;
- 5 I-12 3-{1-oxo-4-(benzylamino)isoindolin-2-yl}piperidine-2,6-dione;
- I-13 2-(2,6-dioxo(3-piperidyl))-4-(benzylamino)isoindoline-1,3-dione;
- I-14 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}propanamide;
- 10 I-15 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}-3-pyridylcarboxamide;
- I-16 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}heptanamide;
- 15 I-17 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}-2-furylcarboxamide;
- I-18 2-azido-*N*-(2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-yl-methyl)-acetamide;
- 20 I-19 2-amino-*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}acetamide;
- 25 I-20 ethyl 6-(*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}carbamoyl)hexanoate;
- I-21 3-((*tert*-butoxy)carbonylamino)-*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}propanamide;
- 30 I-22 3-amino-*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}propanamide;
- I-23 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}-2-thienylcarboxamide;
- 35

- I-24 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}-2-methoxyacetamide;
- 5 I-25 (*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}carbamoyl)methyl acetate;
- I-26 ethyl 2-((*N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}carbamoyl) amino)acetate;
- 10 I-27 *N*-{(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}(ethylamino)carboxamide;
- I-28 2-(2,6-Dioxo(3-piperidyl))-4-[(2-furylmethyl)amino]isoindoline-1,3-dione
- 15 I-29 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)-2-methoxyacetamide;
- I-30 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)heptanamide;
- 20 I-31 {*N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)carbamoyl} methyl acetate;
- I-32 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)pentanamide;
- 25 I-33 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)-2-thienylcarboxamide;
- I-34 methyl {*N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)carbamoyl} formate;
- 30 I-35 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)-2-furylcarboxamide;
- I-36 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)benzamide;
- 35 I-37 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)propanamide;

- 5  
10  
15  
20  
25  
30  
35
- I-38 methyl 3-{*N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)carbamoyl}propanoate;
- I-39 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)-2-phenylacetamide;
- I-40 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)-2-pyridylcarboxamide;
- I-41 *N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)-2-chloroacetamide;
- I-42 2-azido-*N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)acetamide;
- I-43 2-amino-*N*-(2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)acetamide;
- I-44 *N*-(2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl)-2-chloroacetamide;
- I-45 2-azido-*N*-(2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl)acetamide;
- I-46 2-amino-*N*-(2-(2,6-dioxo(3-piperidyl))-1-oxoisindolin-4-yl)acetamide;
- I-47 3-{4-((2-furylmethyl)amino)-1-oxoisindolin-2-yl}piperidine-2,6-dione; or
- I-48 3-(1-oxo-4-(pentylamino)isindolin-2-yl)piperidine-2,6-dione;
- I-49 2-(2,6-dioxo-piperidin-3-yl)-4-(2-methoxy-ethylamino)-isoindole-1,3-dione;
- I-50 2-benzyloxy-*N*-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-yl]-acetamide;
- I-51 2-(2,6-dioxo-piperidin-3-yl)-4-pentylamino-isoindole-1,3-dione;
- I-52 3-chloro-*N*-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-yl]-benzamide;
- I-53 *N*-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1*H*-isoindol-4-yl]-2-phenoxy-acetamide;

- I-54 4-(2-benzyloxy-ethylamino)-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione;
- 5 I-55 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-fluoro-benzamide;
- I-56 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methyl-benzamide;
- 10 I-57 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methoxy-benzamide;
- I-58 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-trifluoromethyl-benzamide;
- 15 I-59 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-nitro-benzamide;
- I-60 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-butyramide;
- 20 I-61 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methylamino-acetamide;
- I-62 2-(2,6-dioxo-piperidin-3-yl)-4-heptylamino-isoindole-1,3-dione;
- 25 I-63 4-chloro-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide;
- I-64 cyclopropanecarboxylic acid [2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide;
- 30 I-65 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-fluoro-benzamide;
- 35

- I-66 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-trifluoromethyl-benzamide;
- 5 I-67 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-methyl-benzamide;
- I-68 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-nitro-benzamide;
- 10 I-69 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-ethoxy-acetamide;
- I-70 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methylsulfanyl-acetamide;
- 15 I-71 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methoxy-benzamide;
- I-72 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-fluoro-benzamide;
- 20 I-73 7-amino-N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl} heptanamide;
- 25 I-74 N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl} butanamide;
- I-75 N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl} benzamide;
- I-76 N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl} phenylacetamide;
- 30 I-77 N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl} -2-pyridylcarboxamide;
- 35 I-78 N- {[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]methyl} undecamide;



- I-79 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}-2-methylpropanamide;
- 5 I-80 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}cyclopentylcarboxamide;
- I-81 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}cyclohexylcarboxamide;
- 10 I-82 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(phenylamino)carboxamide;
- I-83 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(butylamino)carboxamide;
- 15 I-84 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(propylamino)carboxamide;
- I-85 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(cyclohexylamino)carboxamide ;
- 20 I-86 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}[(methylethylamino)]carboxamide;
- 25 I-87 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(octylamino)carboxamide;
- I-88 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(benzylamino)carboxamide;
- 30 I-89 N-{{2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl}methyl}(cyclopropylamino)carboxamide;
- 35 I-90 2-chloro-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide;

- I-91 [2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-carbamic acid benzyl ester;
- 5 I-92 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-acetamide;
- I-93 Pentanoic acid [2-(2,6-dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-amide;
- 10 I-94 N-[2-(2,6-dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-propionamide;
- I-95 N-[2-(2,6-dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-nicotinamide;
- 15 I-96 2-(2,6-dioxo-piperidin-3-yl)-4-[[furan-2-ylmethyl)-amino]-methyl]-isoindole-1,3-dione;
- I-97 N-[2-(2,6-dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide;
- 20 I-98 2-dimethylamino-N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-acetamide;
- 25 I-99 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-methyl-benzamide;
- I-100 Heptanoic acid[2-(2,6-dioxo-piperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-dihydro-1H-isoindol-4-yl]-amide;
- 30 I-101 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3,3-dimethyl-butylamide;
- I-102 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-isobutylamide;
- 35

- I-103 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-phenyl-propionamide;
- 5 I-104 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4-methoxy-benzamide
- I-105 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-trifluoromethyl-benzamide;
- 10 I-106 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-malonamic acid methyl ester;
- I-107 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-3-methoxy-propionamide;
- 15 I-108 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-2-hydroxy-acetamide
- I-109 4-[(furan-2-ylmethyl)-amino]-2-(1-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione;
- 20 I-110 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-isonicotinamide;
- I-111 N-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]-acetamide;
- 25 I-112 {5-[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylcarbamoyl]-pentyl}-carbamic acid benzyl ester;
- 30 I-113 2-(2,6-Dioxo(3-piperidyl))-4-([(cyclohexylamino)thioxomethyl]amino)methyl)isoindole -1,3-dione;
- I-114 2-(2,6-Dioxo(3-piperidyl))-4-([(ethylamino)thioxomethyl]amino)methyl)isoindole -1,3-dione;
- 35

- I-115 2-(2,6-Dioxo(3-piperidyl))-4-({[(propylamino)thioxomethyl]amino} methyl)isoindole -1,3-dione;
- 5 I-116 N-[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisoindolin-4-yl]-2-chloro-benzylamine;
- I-117 {5-[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylcarbamoyl]-pentyl}-carbamic acid benzyl ester;
- 10 I-118 2-Methoxy-N-[2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-acetamide;
- I-119 Pentanoic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide;
- 15 I-120 Heptanoic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide;
- I-121 3-Chloro-N-[2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-benzamide;
- 20 I-122 N-[2-(3-Methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-propionamide;
- 25 I-123 Thiophene-2-carboxylic acid [2-(3-methyl-2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-amide;
- I-124 2-(2,6-Dioxo-piperidin-3-yl)-4-[(5-methyl-furan-2-ylmethyl)-amino]-isoindole-1,3-dione;
- 30 I-125 2-(2,6-Dioxo-piperidin-3-yl)-4-[(5-hydroxymethyl-furan-2-ylmethyl)-amino]-isoindole-1,3-dione;
- I-126 2-(2,6-Dioxo-piperidin-3-yl)-4-[(thiophen-2-ylmethyl)-amino]-isoindole-1,3-dione;
- 35

- I-127 4-(3-Chloro-benzylamino)-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione;
- I-128 2-(2,6-Dioxo-piperidin-3-yl)-4-[(pyridin-3-ylmethyl)-amino]-isoindole-1,3-dione;
- 5 I-129 5-[[2-(2,6-Dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylamino]-methyl]-furan-2-carboxylic acid;
- I-130 4-[(4,5-Dimethyl-furan-2-ylmethyl)-amino]-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione ;
- 10 I-131 4-[(Benzofuran-2-ylmethyl)-amino]-2-(2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione ;
- I-132 4-(3-Chloro-benzylamino)-2-(3-methyl-2,6-dioxo-piperidin-3-yl)-isoindole-1,3-dione;
- 15 I-133 3-[4-(3-Chloro-benzylamino)-1-oxo-1,3-dihydro-isoindol-2-yl]-piperidine-2,6-dione;
- 20 I-134 N-[[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl](cyclopentylamino)carboxamide;
- I-135 N-[[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl](3-pyridylamino)carboxamide Hydrochloride;
- 25 I-136 N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl]piperidylcarboxamide;
- I-137 Tert-Butyl 4-(N-[[2-(2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl](carbamoyl)piperazinecarboxylate;
- 30 I-138 N-[[2-(2,6-Dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl]methyl](diethylamino)carboxamide;
- 35

- I-139 Cyclopropyl-N-{{2-(3-methyl-2,6-dioxo(3-piperidyl))-1,3-dioxoisindolin-4-yl)methyl}carboxamide;
- 5 I-140 N-{{2-(2,6-Dioxo(3-piperidyl))-1-oxoisindolin-4-yl)methyl}cyclopropylcarboxamide;
- I-141 N-{{2-(2,6-Dioxo(3-piperidyl))-1-oxoisindolin-4-yl)methyl}(ethylamino)carboxamide; or
- 10 I-142 Piperazine-1-carboxylic acid  
[2-(2,6-dioxo-piperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-ylmethyl]  
-amide

or a pharmaceutically acceptable salt, hydrate, solvate, clathrate, enantiomer, diastereomer,  
15 racemate, or mixture of stereoisomers thereof.

41. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable vehicle or carrier.
- 20 42. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 10 and a pharmaceutically acceptable vehicle or carrier.
43. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 14 and a pharmaceutically acceptable vehicle or carrier.
- 25 44. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 18 and a pharmaceutically acceptable vehicle or carrier.
45. A pharmaceutical composition comprising a therapeutically effective amount  
30 of a compound of claim 22 and a pharmaceutically acceptable vehicle or carrier.
46. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 28 and a pharmaceutically acceptable vehicle or carrier.
- 35 47. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 34 and a pharmaceutically acceptable vehicle or carrier.

48. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 40 and a pharmaceutically acceptable vehicle or carrier.

49. A method of modulating the production of TNF- $\alpha$  in a mammal comprising  
5 administering to said mammal an effective amount of a compound of claim 1.

50. A method of modulating the production of IL-1 $\beta$  in a mammal comprising administering to said mammal an effective amount of a compound of claim 1.

10 51. A method of modulating the production of IL-10 in a mammal comprising administering to said mammal an effective amount of a compound of claim 1.

52. A method of modulating the production of T-cells in a mammal comprising administering to said mammal an effective amount of a compound of claim 1.

15

53. A method of treating cancer in a mammal, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1, 10, 14, 18, 22, 28, 34, or 40.

20 54. The method of claim 53, wherein the cancer is a solid tumor or a blood born tumor.

55. The method of claim 53, wherein the cancer is cancer of the skin, blood, lymph node, breast, cervix, uterus, gastrointestinal tract, lung, ovary, prostate, mouth, brain,  
25 head, neck, throat, colon, rectum, testes, kidney, pancreas, bone, spleen, liver, bladder, larynx, or nasal passages.

56. The method of claim 53, wherein the cancer is melanoma, multiple myeloma, or a leukemia.

30

57. A method of treating cancer in a mammal, comprising administering to a mammal in need thereof a therapeutically effective amount of a compound of claim 1 and another chemotherapeutic agent.

35

58. The method of claim 57, wherein the other cancer chemotherapeutic agent is paclitaxel, cisplatin, tamoxifen, docetaxel, epirubicin, doxorubicin, irinotecan, leuprolide, bicalutamide, goserelin implant, gemcitabine, or sargramostim.

5 59. The method of claim 57, wherein the other cancer chemotherapeutic agent is an anti-cancer vaccine.

60. A method of treating an inflammatory disorder in a mammal, comprising administering to a mammal in need thereof a therapeutically effective amount of a  
10 compound of claim 1.

61. The method of claim 60, wherein the inflammatory disorder is arthritis, rheumatoid spondylitis, psoriasis, inflammatory bowel disease, post ischemic perfusion injury, or chronic inflammatory pulmonary disease.

15

62. The method of claim 61, wherein the arthritis is rheumatoid arthritis or osteoarthritis.

63. A method of treating heart disease in a mammal comprising administering to  
20 a mammal in need thereof a therapeutically effective amount of a compound of claim 1.

64. A method of modulating the production of TNF- $\alpha$  in a mammalian cell or tissue comprising contacting an effective amount of a compound of claim 1.

25 65. A method of modulating the production of IL-1 $\beta$  in a mammalian cell or tissue comprising contacting an effective amount of a compound of claim 1.

66. A method of modulating the production of IL-10 in a mammalian cell or tissue comprising contacting an effective amount of a compound of claim 1.

30

67. A method of modulating the production of T-cells in a mammalian cell or tissue comprising contacting an effective amount of a compound of claim 1.

35



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/50401

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/04 C07D401/14 C07D407/14 C07D409/14 A61K31/445  
A61K31/44 A61K31/495 A61P29/00 A61P35/00 A61P37/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 46258 A (CHEN ROGER SHEN CHU ;MAN HON WAH (US); CELGENE CORP (US); MULLER G) 16 September 1999 (1999-09-16) Claims 1,9 the whole document	1-67
X	US 5 874 448 A (CHEN ROGER SHEN-CHU ET AL) 23 February 1999 (1999-02-23) Claims 1,9 the whole document	1-67
X	WO 98 54170 A (CHEN ROGER SHEN CHU ;CELGENE CORP (US); MULLER GEORGE W (US); STIR) 3 December 1998 (1998-12-03) Claim 6, ex. 12 the whole document	1-67

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

4 June 2002

Date of mailing of the international search report

13/06/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fritz, M

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/50401

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 03502 A (CHEN ROGER SHEN CHU ;CELGENE CORP (US); MULLER GEORGE W (US); STIR) 29 January 1998 (1998-01-29) Claim 1, ex. 15-17 the whole document ----	1-67
X	US 5 635 517 A (CHEN ROGER S-C ET AL) 3 June 1997 (1997-06-03) Ex. 1 the whole document ----	1-67
X	MULLER G W ET AL: "Amino-substituted thalidomide analogs: potent inhibitors of TNF-alpha production" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 9, no. 11, 7 June 1999 (1999-06-07), pages 1625-1630, XP004169632 ISSN: 0960-894X Compound 8a the whole document ----	1-67
X	MIYACHI H ET AL: "Tumor necrosis factor-alpha production enhancing activity of substituted 3'-methylthalidomide: Influence of substituents at the phthaloyl moiety on the activity and stereoselectivity" CHEMICAL AND PHARMACEUTICAL BULLETIN, PHARMACEUTICAL SOCIETY OF JAPAN. TOKYO, JP, vol. 46, no. 7, July 1998 (1998-07), pages 1165-1168, XP002107774 ISSN: 0009-2363 Compound 9c the whole document -----	1-67

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International Application No  
**PCT/US 01/50401**

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9946258	A	16-09-1999	US 5955476 A	21-09-1999
			AU 1413899 A	27-09-1999
			CA 2317834 A1	16-09-1999
			CN 1282330 T	31-01-2001
			EP 1062214 A1	27-12-2000
			FI 20001192 A	14-07-2000
			HU 0100615 A2	28-10-2001
			JP 2002506068 T	26-02-2002
			NO 20002529 A	30-06-2000
			NZ 504650 A	30-11-2001
			PL 340547 A1	12-02-2001
			SK 7382000 A3	11-12-2000
			TR 200002627 T2	21-11-2000
			WO 9946258 A1	16-09-1999
US 5874448	A	23-02-1999	FI 20001192 A	14-07-2000
			NO 20002529 A	30-06-2000
			SK 7382000 A3	11-12-2000
			US 5955476 A	21-09-1999
WO 9854170	A	03-12-1998	AU 715779 B2	10-02-2000
			AU 741982 B2	13-12-2001
			AU 7701298 A	30-12-1998
			CA 2261762 A1	29-01-1998
			CN 1239959 A	29-12-1999
			CN 1258293 T	28-06-2000
			CZ 9900202 A3	16-06-1999
			EP 0925294 A1	30-06-1999
			EP 0984955 A1	15-03-2000
			FI 990101 A	19-03-1999
			FI 992490 A	27-01-2000
			HU 0003217 A2	28-06-2001
			HU 9903929 A2	28-05-2000
			JP 2002501536 T	15-01-2002
			JP 2001503384 T	13-03-2001
			NO 995751 A	28-01-2000
			NZ 333903 A	28-02-2000
			NZ 501429 A	30-11-2001
			PL 332867 A1	25-10-1999
			PL 337124 A1	31-07-2000
			SK 9199 A3	12-07-1999
			SK 163099 A3	12-06-2000
			TR 200000107 T2	21-07-2000
			WO 9803502 A1	29-01-1998
			WO 9854170 A1	03-12-1998
			US 6281230 B1	28-08-2001
			US 6316471 B1	13-11-2001
			US 6335349 B1	01-01-2002
WO 9803502	A	29-01-1998	US 5635517 A	03-06-1997
			US 5798368 A	25-08-1998
			AU 715779 B2	10-02-2000
			AU 3899897 A	10-02-1998
			CA 2261762 A1	29-01-1998
			CN 1239959 A	29-12-1999
			CZ 9900202 A3	16-06-1999
			EP 0925294 A1	30-06-1999
			FI 990101 A	19-03-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
WO 9803502	A	JP 2001503384 T	13-03-2001	
		NZ 333903 A	28-02-2000	
		PL 332867 A1	25-10-1999	
		SK 9199 A3	12-07-1999	
		WO 9803502 A1	29-01-1998	
		US 6281230 B1	28-08-2001	
		US 6316471 B1	13-11-2001	
		US 6335349 B1	01-01-2002	
		US 2002045643 A1	18-04-2002	
		AU 741982 B2	13-12-2001	
		AU 7701298 A	30-12-1998	
		CN 1258293 T	28-06-2000	
		EP 0984955 A1	15-03-2000	
		FI 992490 A	27-01-2000	
		HU 0003217 A2	28-06-2001	
		HU 9903929 A2	28-05-2000	
		JP 2002501536 T	15-01-2002	
		NO 995751 A	28-01-2000	
		NZ 501429 A	30-11-2001	
		PL 337124 A1	31-07-2000	
		SK 163099 A3	12-06-2000	
		TR 200000107 T2	21-07-2000	
		WO 9854170 A1	03-12-1998	
US 5635517	A	03-06-1997	AU 715779 B2	10-02-2000
			AU 3899897 A	10-02-1998
			CA 2261762 A1	29-01-1998
			CN 1239959 A	29-12-1999
			CZ 9900202 A3	16-06-1999
			EP 0925294 A1	30-06-1999
			FI 990101 A	19-03-1999
			JP 2001503384 T	13-03-2001
			NZ 333903 A	28-02-2000
			PL 332867 A1	25-10-1999
			SK 9199 A3	12-07-1999
			WO 9803502 A1	29-01-1998
			US 6281230 B1	28-08-2001
			US 6316471 B1	13-11-2001
			US 6335349 B1	01-01-2002
			US 2002045643 A1	18-04-2002